Doubling diet fat on sugar ratio in children with mitochondrial OXPHOS disorders: Effects of a randomized trial on resting energy expenditure, diet induced thermogenesis and body composition

Laurent Beghin, Stephanie Coopman, Manuel Schiff, Joseph Vamecq, Karine Mention-Mulliez, Regis Hankard, Jean-Marie Cuisset, Helene Ogier, Frederic Gottrand, Dries Dobbelaere

► To cite this version:

Laurent Beghin, Stephanie Coopman, Manuel Schiff, Joseph Vamecq, Karine Mention-Mulliez, et al.. Doubling diet fat on sugar ratio in children with mitochondrial OXPHOS disorders: Effects of a randomized trial on resting energy expenditure, diet induced thermogenesis and body composition. Clinical Nutrition, 2016, 35 (6), pp.1414-1422. 10.1016/j.clnu.2016.03.015 . hal-02177005

HAL Id: hal-02177005

<https://hal.univ-lille.fr/hal-02177005>

Submitted on 8 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Doubling diet fat on sugar ratio in children with mitochondrial OXPHOS disorders: Effects of a randomized trial on resting energy expenditure, diet induced thermogenesis and body composition

Laurent Béghin ^{a,b,*}, Stéphanie Coopman ^a, Manuel Schiff ^c, Joseph Vamecq ^d, Karine Mention-Mulliez ^e, Régis Hankard ^{f,g}, Jean-Marie Cuisset ^h, Hélène Ogier ^c, Frédéric Gottrand ^{F a,b}, Dries Dobbelaere ^e.

^a Centre d'Investigation Clinique, CIC-1403–Inserm–CH&U, Lille University Hospital F-59037 Lille France.

^b LIRIC- Lille Inflammation Research International Center/UMR U995 Inserm, Lille France

^c Reference center for Inherited Metabolic Diseases, Robert Debré University Hospital, Paris, France

^d Inserm, Department of Biochemistry and Molecular Biology, HMNO, CBP, CHRU Lille and and RADEME EA 7364, Lille Nord of France University, F-59037 Lille, France

^e Reference center for Inherited Metabolic Diseases in child and adulthood, Lille University Children's Hospital Jeanne de Flandre, and RADEME EA 7364, Lille Nord of France University, F-59037 Lille, France

^f Inserm U 1069, F Rabelais University, Tours, F-37000, France

^g CIC-1426- Inserm-CH&U of Robert Debré University Hospital, F-75935 Paris, France

^h Pediatric Neurology Unit Lille University Hospital, F-59037 Lille, France

Short title: Doubling diet fat on sugar ratio in mitochondriopathies

* Corresponding author : Dr Laurent BEGHIN, LIRIC-UMR-995-Inserm and CIC-1403-Inserm-CH&U de Lille

Centre d'Investigation Clinique, Antenne pédiatrique, Hôpital Jeanne de Flandre, CHRU de Lille, F-59037 LILLE Cédex, Tél : +33 3 20 44 60 58, Fax : +33 3 20 44 66 87,

E-mail addresses: laurent.beghin@chru-lille.fr (L. Béghin), Stephanie.coopman@chru-lille.fr (S. Coopman), manuel.schiff@aphp.fr (M. Schiff), joseph.vamecq@inserm.fr (J. Vamecq), karine.mention@chru-lille.fr (K. Mention-Mulliez), regis.hankard@univ-tours.fr (R. Hankard), (J.M. Cuisset), (H. Ogier), Frederic.gottrand@chru-lille.fr (F. Gottrand), dries.dobbelaere@chru-lille.fr (D. Dobbelaere)

Abbreviations:

ANOVA : Analysis Of Variance.

BD : Basal Diet

CD : Challenging Diet

CONSORT : Consolidated Standards of Reporting Trials

DIT : Diet Induced Thermogenesis

MOD : Mitochondrial OXPHOS Disorder

OXPHOS : Oxidative Phosphorylation

REE : Resting Energy Expenditure

SUMMARY

Background and aims: Mitochondrial OXPHOS disorders (MODs) affect one or several complexes of respiratory chain oxidative phosphorylation. An increased fat/low-carbohydrate ratio of the diet was recommended for treating MODs without, however, evaluating its potential benefits through changes in the respective contributions of cell pathways (glycolysis, fatty acid oxidation) initiating energy production. Therefore, the objective of the present work was to compare Resting Energy Expenditure (REE) under basal diet (BD) and challenging diet (CD) in which fat on sugar content ratio was doubled. Diet-induced thermogenesis (DIT) and body compositions were also compared. Energetic vs regulatory aspects of increasing fat contribution to total nutritional energy input were essentially addressed through measures primarily aiming at modifying total fat amounts and not the types of fats in designed diets.

Methods: In this randomized cross-over study, BD contained 10% proteins/30% lipids/60% carbohydrates (fat on sugar ratio = 0.5) and was the imposed diet at baseline. CD contained 10% proteins/45% lipids/45% carbohydrates (fat on sugar ratio = 1). Main and second evaluation criteria measured by indirect calorimetry (QUARK RMR®, Cosmed, Pavona; Italy) were REE and DIT, respectively. Thirty four MOD patients were included; 22 (mean age 13.2 ± 4.7 years, 50% female; BMI 16.9 ± 4.2 kg/m²) were evaluated for REE, and 12 (mean age 13.8 ± 4.8 years, 60% female; BMI 17.4 ± 4.6 kg/m²) also for DIT. OXPHOS complex deficiency repartition in 22 analysed patients was 55% for complex I, 9% for complex III, 27% for complex IV and 9% for other proteins.

Results: Neither carry-over nor period effects were detected ($p=0.878$; ANOVA for repeated measures). REE was similar between BD vs CD (1148.8 ± 301.7 vs 1156.1 ± 278.8 kcal/day ; $p=0.942$) as well as DIT (peak DIT 260 vs 265 kcal/day ; $p=0.842$) and body composition (21.9 ± 13.0 vs 21.6 ± 13.3 % of fat mass ; $p=0.810$).

Conclusion: Doubling diet fat on sugar ratio does not appear to improve, per se, energetic status and body composition of patients with MODs.

Keywords: Mitochondrial disorders or diseases, Energy expenditure, Diet induced thermogenesis, Body composition.

1. Introduction

Mitochondrial OXPHOS disorders (MODs) encompass a group of rare genetic disorders affecting respiratory chain oxidative phosphorylation (OXPHOS) and resulting from mitochondrial or nuclear DNA defects [1,2]. The OXPHOS system plays a key role in transferring energy from macronutrients to ATP through a sequence of coordinated reactions by which macronutrients are oxidized. OXPHOS comprises five major protein-lipid enzyme complexes which are located in mitochondrial inner membrane and work locally in synergy with the labile electron acceptors ubiquinone (an amphiphilic isoprenoid compound) and cytochrome *c* (a 12 kDa peptide). Electrons generated during glucose, fatty acids and amino acid oxidations are channeled across complexes I and II to be transferred sequentially to coenzyme Q10, complex III, and complex IV. Complexes I, III, and IV use part of the energy produced by the electron transfer flux to extrude protons across the mitochondrial inner membrane, from mitochondrial matrix to mitochondrial intermembrane space. The resulting proton gradient is consumed with energy utilization and production by complex V which condenses adenosine diphosphate and inorganic phosphate into adenosine trisphosphate (ATP) [3,4]. These overall mechanisms and players for ATP production are impaired in disorders affecting OXPHOS, jeopardizing various cell functions notably those which require a high energy supply. [5]. Currently, no satisfactory curative treatment for MODs exists. Only symptomatic treatments of MODs are available [2,7,8]. Interestingly, dietary modulations have been considered as regards to the key-role of mitochondria in energy production from macronutrients [9,10].

Increasing fat vs carbohydrate content in the diet has been recommended for the treatment of MOD. Several reasons justify this recommendation. (i) Mitochondrial oxidation of NADH is thought to be diminished in MOD [5,11] patients especially those affected with respiratory chain Complex I (CI) deficiencies, it might be by-passed by FADH₂, an alternative carrier of reducing equivalents, the electrons of which can enter the respiratory chain distal to complex I. (ii) Supply in FADH₂ to the mitochondria can be increased (relative to NADH) by increasing the amount of triacylglycerols and fatty acids in the diet. On the basis of stoichiometry of mitochondrial oxidations, fatty acids yield a ratio of FADH to NADH of 0.5, whereas glucose yields a much lower ratio of 0.2. In addition, fatty acids in contrast to glucose are endowed with uncoupling properties towards OXPHOS [12]. Though uncoupling OXPHOS lowers net ATP production by mitochondria, it virtually accelerates mitochondrial respiration through increased

105 electron transfer rates, a feature which improves mitochondrial oxidations regulated by the
106 NADH/NAD⁺ and FADH₂/FAD ratios.

107 In the wake of this rationale, we hypothesized that increasing fat to glucose ratio within an
108 isoenergetic and isonitrogenous diet should improve energy substrate oxidation in MOD patients.
109 As a corollary, this dietary measure should actually increase resting energy expenditure (REE), an
110 assumption also based on *in vitro* data on human cell [13,14] and animal studies [15], and *in vivo*
111 data obtained in complex I deficiency mitochondrial myopathy patients [16–19]. The aim of the
112 present study was, therefore, to study REE under basal diet (referred throughout the manuscript
113 to as the basal diet (BD)) challenged by a diet in which fat on carbohydrate content ratio was
114 doubled (referred to as the challenging diet (CD)) as a potential treatment in children with a MOD.
115 In these children, diet induced thermogenesis (DIT) and body composition were also measured to
116 get a better account for whole body energetic balances under the two diets.

117

118 **2. Materials and methods**

119

120 *2.1. Trial design and participants*

121 This 3-year prospective study (2008-2011) included MOD patients from the reference
122 centers for Inherited Metabolic Diseases of Lille and Paris (Jeanne de Flandre Lille University
123 Hospital ; Robert Debré Paris University Hospital). Inclusion criteria were : (i) Child or adult
124 between 5 to 21 years with MOD [20], (ii) MOD subjects given a stable balanced diet having the
125 characteristics of basal diet (BD : 60 % carbohydrates, 30 % lipids, 10 % proteins) for at least one
126 month. Subjects were excluded from the study when they required hospitalization or presented
127 with any acute condition (such as infection) known to interfere with energy metabolism
128 assessment.

129 As shown in flow CONSORT diagram [21] on (Figure 1), from 36 MOD subjects included over
130 a period of four years, 22 accepted compliance to the diets and examinations, and were analysed
131 for principal evaluation criterion (*ie* REE in Group 2), and from these 22, only 12 accepted to
132 undergo the secondary evaluation criterion (*ie* DIT in Group 3).

133 This study was a randomized cross-over study (Figure 2). Method used to generate
134 sequence was a simple randomisation by one-to-one allocation ratio. Subject randomisation was
135 performed using sealed envelopes. REE were measured at four time points (*ie* Visit 1: baseline,
136 Visit 2: after one month of basal or challenging diet ; Visit 3: after wash-out period consisted of
137 one month of BD ; and Visit: 4 after one month of basal or challenging diet; see Figure 2). Neither

138 carry-over effect, nor period effect was detected ($p = 0.878$; ANOVA for repeated measures). All
139 assessments were performed in two Clinical Investigation Centers (CIC-1403-Inserm-CH&U of Lille;
140 CIC-9202-Inserm-AP-HP of Paris). Before each assessment visit, all subjects arrived by car to the
141 Clinical Investigation Center at 8:00 h, being fasted from 20:00 h on the previous day. Weight and
142 height were first measured, and then compliance with the imposed diet was evaluated.

143

144 2.2. Nutrition protocol management

145 Intervention of the trial consisted of a CD diet period of one month (*ie* 10 % proteins, 45 %
146 lipids, 45 % carbohydrates). CD and BD diets were isoenergetic and isonitrogenous diets.

147 Each patient was randomly assigned to the basal diet (BD) for the control period *vs* the
148 challenging diet (CD) for the intervention period. Each one-month period was thereafter inversed
149 according to a cross-over trial. BD consisted of 10 % proteins, 30 % lipids, 60 % carbohydrates. To
150 reach this BD, research dieticians used a specific booklet communicated to subjects and/or their
151 parents to lower fat intake. CD consisted of 45 % lipids, 45 % carbohydrates ; 10 % proteins.

152 DIT was performed after a calibrated breakfast corresponding to 20% of recommended
153 daily energy intake according to Martin *et al's* formula [22]. According to the cross-over design,
154 breakfast had a BD in control period and a CD in intervention period. Basal content breakfast
155 consisted of bread, skimmed milk and jam. Challenging breakfast consisted of bread, full-cream
156 milk and chocolate paste. After breakfast, meals were re-weighted and diet energy supplies were
157 analysed by the difference between served and returned meals [23].

158 The assessment of energy intake and diet composition was performed using dietary record
159 technique previously reported [24]. The technique used a specific chart report form including how
160 to record the foods consumed, using size instruments such as pictures of graduated bowls, cups,
161 dishes, number of spoons. Each file corresponding to one day was separated in four parts:
162 breakfast, lunch, dinner and snack. Lunch and dinner were detailed as starter, main course, and
163 desert [25]. Both oral and written instructions were given to each patient and/or parents on how
164 to keep accurate records using the size instruments, and the parents assisted the children in
165 recording, identifying and quantifying the foods consumed. Subjects detailed daily food intakes on
166 the specific chart report form during 7 d [26]. The same project related dietician reviewed each
167 specific chart report. Quantification of meals and drinks were checked by the same trained
168 dietician, the parents and the child together. The dietician was well trained to identify both miss
169 and over-reporting and coherency of food records. The portion sizes were estimated using a three
170 dimensional portion size instrument with French current meal food photographs corresponding to

171 an exact BILNUT software (version 6 : Paris, France). This method has been shown to give unbiased
172 records of energy intake in lean subjects up to 9 y old and showed that children aged 8-15 y were
173 able to estimate food quantity to within = 10 % of the amount really eaten, suggesting that
174 children could quantify their food intake with reasonable accuracy [27].

175

176 2.3. Indirect calorimetry assessment of energy expenditure

177 The subject rested recumbent on a hospital bed watching a videotape for 15 min. Same
178 quiet cartoons or videos (depending on age range of the subject) were used for all the subjects
179 [28]. Oxygen consumption (Vo_2 in mL/min) and carbon dioxide production (Vco_2 in mL/min) were
180 determined by using the open-circuit indirect calorimeter validated QUARK® RMR (Cosmed,
181 Pavona di Albano, Italy). The validity and accuracy measurements were guaranteed by yearly
182 calibration of the flow settings by the manufacturer of the calorimeter and guaranty a high
183 accuracy of QUARK® RMR between study centres [29]. Calibration of zero, span and delay
184 alignment of the O_2 and CO_2 gas analyzers was performed daily before each test using a certified
185 calibration gas.

186 The QUARK® RMR was equipped with a canopy hood for spontaneously breathing subjects.
187 With this device, flow rate was directly measured with a digital turbine flowmeter. Accuracy of the
188 flowmeter was 2%. Ventilatory rate was regulated directly by the QUARK® RMR. During each test,
189 the readings were controlled and eventually compensated by means of periodic automatic room
190 air calibrations. Response times of O_2 and CO_2 sensors were less than 120 ms. The O_2 analyzer was
191 a paramagnetic sensor, which had a measuring range from 0 to 30% in the canopy mode. Accuracy
192 of the O_2 sensor was 0.02 %. The CO_2 analyzer was an infrared digital sensor with a measuring
193 range standing from 0 to 10%. Accuracy of the CO_2 sensor was 0.02%. The flow meter which
194 detects the ventilation rate of the canopy was a bidirectional turbine with an 18 mm diameter.
195 The ventilation range was from 0 to 80 L/min. Accuracy of the flowmeter was 2%. QUARK® RMR
196 software utilized the Weir equation to assess Resting Energy Expenditure (REE) [30].

197 DIT was measured at two occasions (Visit 2 and Visit 4) after REE measurement. DIT was
198 derived from postprandial energy expenditure (PEE) as described by Dabbech M *et al* [31]. PEE
199 assessment started immediately after a calibrated breakfast (see above for energetic ratio). DIT
200 was computed by subtracting REE from PEE (*ie* $\text{DIT} = \text{PEE} - \text{REE}$) according to Dabbech M *et al* [31].

201

202

203

2.4. Assessment of body composition by Dual X-ray Absorptiometry

The DXA (Dual X-Ray Absorptiometry) instrument was a Lunar DPX-IQ “pencilbeam” (Lunar Radiation Corporation, Madison, WI, USA). Whole-body scan time was 560–590 s, and the radiation dose was less than 0.03 mrem ($\frac{1}{4}$ 0.3 mSv). Total body image acquisition and analysis were obtained by following strictly manufacturer’s specified instructions. The use of two photon energies reduced errors due to irregular soft-body-mass and body contours. Two photon energies discriminated two substances in a given system to a high accuracy. When more than two substances, the discrimination depended on the number of additional substances, their attenuation characteristics, and their relative contents [32]. FFM (fat-free mass), FM (fat mass) and weight were calculated using the software provided by the manufacturer, and was adapted for the subject age.

2.5. Regulatory requirement

The protocol was approved by French appropriate regulatory authorities (Direction Générale de la Santé; DGS 2007-A0018-50) and ethical review committee (Comité de Protection des Personnes Nord-Ouest IV ; CPP 07/32). Study protocol was also declared to clinical trial.gov register (NCT02385565). Participation in the study was voluntary and written informed consent was obtained from both parents and children. Written informed consent was obtained by the subject when patient was aged over 18 years.

2.6. Statistical analysis

Required sample size was computed using previously published data from complex I deficiency (CID) mitochondrial myopathy patients [16,17] with REE = 1120 ± 93 Kcal/min in basal condition (Glucose infusion). REE obtained in hyperlipid MOD condition (Triacylglycerol infusion) were 5 % higher at 1175 ± 93 Kcal/min. We hypothesised a 10% difference of REE between basal and challenging diets. According to the cross-over study design of the study and 80% statistical power, the sample size was defined at 20 patients to detect a significant difference between diets.

Statistical analyses were performed using SAS V.9 (Cary, USA). According to a Shapiro-Wilk test, all variables had a normal distribution. In this context, parametric test χ^2 for ratio, and to a cross-over trial design: period effect was tested by ANOVA analysis for repeated measures, carry over effect was tested by Student *t* test. Difference between two diets was tested by Student *t* test when number of subjects was above 15 [33,34]. When number of subjects was under 15, we

236 used non-parametric tests: *ie* Wilcoxon for paired analyses and Mann-Whitney for non-paired
237 analyses.

238

239

240

241 3. Results

242

243 3.1. Anthropometric, biochemical defects and main clinics of enrolled patients

244 Anthropometric characteristics of patients from each of the 3 groups defined in the flow
245 CONSORT diagram illustrated at Figure 1 are presented on Table 1. No statistical differences were
246 observed between included and analysed subjects for REE, and DIT measurements. Table 2 further
247 summarizes main clinical and biochemical results of respiratory chain activities in skeletal muscle
248 and/or cultured skin fibroblasts. The most frequent defect was complex I deficiency at 55% (n=12).
249 Table 3 presents a minimum set for anthropometric data description of patients with complex I
250 deficiency and those with other defects. There was no difference between groups excepted for Z-
251 score BMI which was significantly lower in the group of patients with defects other than complex I
252 deficiency.

253

254 3.2. Study of patients undergoing REE measurements

255 3.2.1. Food intakes

256 Food intakes in fat, carbohydrates and proteins were determined during the cross-over
257 periods in the 22 patients undergoing REE measurements (Table 4).

258 3.2.2. Metabolic REE measurements and anthropometrics

259 As mentioned in the experimental section, REE were measured at four time points (see
260 Figure 2). No carry-over effect and no period effect was detected for the present study ($p = 0.878$;
261 ANOVA for repeated measures).

262 Table 5 presents full metabolic data at baseline and after basal vs challenging diet
263 conditions. Measured REE were close to the predictive REE. No significant differences were
264 observed between the basal and challenging periods for any parameters (*ie* VO_2 , VCO_2 , REE, RQ)
265 even when these parameters were adjusted by body composition parameters (*ie* weight, fat mass,
266 fat free mass, muscle weight). Patient body composition parameters did not also differ between
267 basal and challenging diet periods for whole analyzed population (table 6). There were no
268 differences between basal and challenging diet periods for measured REE and anthropometrics in
269 patients with complex I deficiency nor in the group of patients affected by others defects. (Tables
270 7 and 8)

271

272

273

274
275
276
277
278
279
280
281
282
283
284
285
286
287

288

289

290

291

292
293
294
295
296
297
298
299
300
301
302
303

3.3. Study of patients undergoing DIT measurements

3.3.1. Food intakes

In a subset of 12 subjects, it was possible to measure DIT. Table 9 presents respective carbohydrate, lipid and protein contents of basal and challenging breakfasts. According to these data, designed diet conditions (*ie* basal vs challenging) were in practice respected

3.3.2. DIT measurements

Figure 3 gives time-courses of DIT over a postprandial period of 250 *min*. Maximum DIT was obtained at 100 ± 27 *min* after basal breakfast and 122 ± 29 *min* after challenging breakfast ($p = 0.861$). No significant differences in DIT optima, areas under the curve, time durations of postprandial periods were observed between basal vs challenging diet conditions. Similarly, RQ during DIT measurements were not significantly different between basal and challenging diet conditions (Figure 4).

4. Discussion

Because of the well-established ability of fats to induce mitochondrial biogenesis, we aimed to test efficacy of an increased fat on carbohydrate ratio of the diet to manage nutrition in MODs in order to improve energy expenditure metabolism [15–18]. Our study was designed to determine to what extent and how REE, and secondarily DIT and body composition, may be impacted by doubling fat vs carbohydrate contribution to nutritional energy intake. The collected data do not confirm our primary hypothesis of a higher REE in challenging diet conditions. Therefore, they do not *a priori* corroborate previous benefits suggested for high-fat diet in MODs in a specific subset of patients using a specific type of fat enrichment. Our study present several strengths (*i*) the cross-over design that is perfectly adapted to the aims of our study and the rarity and heterogeneity of MODs, without period and carry-over effect, (*ii*) an adequate sample size, (*iii*) a very well controlled diet, (*iv*) a well characterized population and (*v*) the use of gold standard methodology based on previously validated tools (*ie* indirect calorimetry, body composition

analysis). Duration of the cross-over periods of one month was considered to be sufficient for a high stability of energy metabolism and changes in REE and DIT. The absence of differences of REE between basal and challenging diets observed in our study might be explained either by the wide genetic heterogeneity usually observed in subjects with a mitochondrial disease [35], and masking the power/homogeneity effect or by no impact of the challenging diet on REE, DIT and body composition. In this respect, majority of previous trials have been performed on complex I MOD [15,16,36,37] and, until now, trials mixing patients with distinct complex deficiencies were lacking. The present study overcomes this gap and allows comparison with previous studies, some extra statistical analyses were performed on REE and body composition in the subset of complex I MOD patients included in our study. These extra analyses did not detect any difference in REE and body composition in this subset of patients (Tables 7 and 8). This result does not reject the heterogeneity of our population as a cause mentioned above for a lack of benefit of challenging diet in MODs as even in Complex I defect, there is a huge genetic variability [38]. Concerning durations of the cross-over period, one month was an adapted period to observe any changes in REE and DIT in view of one week as a known successful period to normalize complex I activity in malnourished patients [37]. We did not observe any difference in DIT (*ie* energy expenditure and RQ) as Flatt observed in healthy subject [39,40]. The last hypothesis of the lack of impact of CD on energy metabolism despite a higher amount of energy intake at breakfast (+ 37 % see Table 9) might result from the fact that the CD tested was at a 45% instead of a 50% dietary fat content, though no effect of CD was also observed by De Meer K et al [41]. An important feature also is that our CD was primarily aimed at doubling fat vs carbohydrate contribution to nutritional energy intake and not to enrich fat qualitatively with a specific type of fats. Using a specific type of fat Roef *et al* [16] found that triacylglycerol infusion improved some metabolic parameters (*ie* VCO₂, respiratory quotient) during physical exercise in a sample of patients with deficient complex I. The significant effect observed in their study may be accounted for by several factors :

(i) a specific effect of triacylglycerol intravenous infusion which is more energetic efficient than oral intake because of a high rate of systemic delivery and contents in medium-chain triglycerides (MCT). MCT are more energetic efficient than long-chain triglycerides because of their faster handling by metabolic pathways. This advantage is used in ketogenic diets where high fat ratios force the body to use fats instead of carbohydrates [42]. The underlying energetic change in oxidative

phosphorylation (OXPHOS) lies in fatty acids yielding a ratio of FADH to NADH of 0.5, whereas glucose yields a much lower ratio of 0.2

(ii) measurements performed during a physical exercise occur during sustained substrate oxidation, allowing a better detection of differences in energetic metabolism,

(iii) a specific population response (only 4 subjects in this study). The absence of response in our population even if we analyzed a subgroup of complex I-deficient patients might be attributed to the lack of mechanisms covered by parenteral route, MCT and physical exercise which are all conditions absent from our experimental protocols. This does not preclude that the ketogenic diet does not improve mitochondrial oxidations as a source of metabolic energy, but this feature unfortunately failed to be detected by our study. In this respect, the lower fat enrichment of our CD vs classic ketogenic diet (43.6 % of fat in our CD vs 90% in ketogenic diet) is another aspect to take into account.

As MODs may exhibit a wide range of symptoms including developmental delay, seizures, vision and hearing impairments, autonomous nervous dysfunction, gastrointestinal signs, endocrine disturbances and failure to thrive, a special interest of these symptoms were checked during the study period. No adverse reactions, adverse events of special interest, or serious adverse events were observed excepted some fatigue in 6 patients from 24 randomized patients only during basal diet period (not significant). As a whole, doubling fat vs carbohydrate contribution to nutritional energy intake does not improve, *per se*, energetic status and body composition of patients with MODs but does not also impact negatively patients. However, undernutrition observed in defects other than complex I could suppose a higher defect in weight energy balance in these patients vs complex I-deficient patients.

5. Conclusion

As a whole, this study does not provide evidence that doubling fat vs carbohydrate ratio of the diet increases whole-body energy expenditure in patients with MOD. We were not able to

365 demonstrate a lowering of the metabolic challenge induced by substrate oxidation as a suggested
366 potential benefit in CD for mitochondrial diseases [5]. Our study suggests that underlying
367 mechanisms for a potential beneficial effect for CD does not a priori involve modulations of REE,
368 nor DIT. The wide heterogeneity of the studied population and/or the amount/quality of lipids in
369 the diet might play a role as well in the generated data and subsequent conclusions

370

371

372 **Conflict of Interest Statement & Statement of Authorship.**

373 Frédéric Gottrand has received consulting fees from Numico Clinical Nutrition, lecture fees
374 from SMS and grant support from Danone Research. The remaining authors state no conflict of
375 interest.

376

377

378

379

380 **Acknowledgments**

381 We thank parents and patients who have taken part in the study. Pascale HINCKER & Sarah
382 KAMMENEY for diet composition management. We thank Christelle GUIMBER study nurse from
383 CIC for paramedical/logistical assistance. We thank Brigitte PORAS, José CANOMORALES, Stéphane
384 DESCHILDER, Nathalie DUQUESNOY, Jeanne-Marie DESCAMPS-BEN, Valérie HELD, Isabelle
385 CHARDON, Laurence CRINON for DXA acquisition and Dr Georges LION for DXA validation and
386 interpretation. We thank Muriel BEUVRY and Anne GAUTREAU (CIC-PT-1403-CH&U-Inserm de
387 Lille, France) for typing this manuscript.

388

389 **Funding**

390 Lille University Hospital was the trial sponsor of this study. This study was founded by a
391 grant from French ministry of health (PHRC interregional N°2003/1901).

392

393

394

395

396

397 **References**

398
399 [1] Vafai SB, Mootha VK. Mitochondrial disorders as windows into an ancient organelle. *Nature* 2012;
400 491:374-383.

401 [2] Koopman WJ, Willems PH, Smeitink JA. Monogenic mitochondrial disorders. *N Engl J Med* 2012; 366:
402 1132-1141.

403 [3] Di Donato S. Disorders related to mitochondrial membranes: pathology of the respiratory chain and
404 neurodegeneration. *J Inherit Metab Dis* 2000;23:247-263.

405 [4] Roede JR, Go YM, Jones DP. Redox equivalents and mitochondrial bioenergetics. *Methods Mol Biol*
406 2012;810:249-280.

407 [5] Munnich A, Rotig A, Chretien D, Cormier V, Bourgeron T, Bonnefont JP, Saudubray JM, Rustin P.
408 Clinical presentation of mitochondrial disorders in childhood. *J Inherit Metab Dis* 1996 ;19:
409 521-527.

410 [6] Thorburn DR. Mitochondrial disorders: prevalence, myths and advances. *J Inherit Metab Dis* 2004;
411 27:349-362.

412 [7] Schiff M, Benit P, Coulibaly A, Loublier S, El-Khoury R, Rustin P. Mitochondrial response to controlled
413 nutrition in health and disease. *Nutr Rev* 2011 69: 65-75.

414 [8] Schiff M, Benit P, Jacobs HT, Vockley J, Rustin P. Therapies in inborn errors of oxidative metabolism.
415 *Trends Endocrinol Metab* 2012;23:488-495.

416 [9] Passarella S, Atlante A, Valenti D, de BL. The role of mitochondrial transport in energy metabolism.
417 *Mitochondrion* 2003;2:319-343.

418 [10] Wallace DC, Fan W, Procaccio V. Mitochondrial energetics and therapeutics. *Annu Rev Pathol* 2010;
419 5:297-348.

420 [11] Gwynne J. The role of nutrition in mitochondrial an metabolic diseases. A primary care physician's
421 guide 2000;15-16.

422 [12] Mokhova EN, Khailova LS. Involvement of mitochondrial inner membrane anion carriers in the
423 uncoupling effect of fatty acids. *Biochemistry (Mosc)* 2005;70:159-63

424 [13] Doctor RB, Bacallao R, Mandel LJ. Method for recovering ATP content and mitochondrial function
425 after chemical anoxia in renal cell cultures. *Am J Physiol* 1994;266:C1803-C1811.

426 [14] Santra S, Gilkerson RW, Davidson M, Schon EA. Ketogenic treatment reduces deleted mitochondrial
427 DNAs in cultured human cells. *Ann Neurol* 2004;56:662-669.

428 [15] Schiff M, Benit P, El-Khoury R, Schlemmer D, Benoist JF, Rustin P. Mouse studies to shape clinical
429 trials for mitochondrial diseases: high fat diet in Harlequin mice. *PLoS One* 2011;6:e28823.
430 10.1371/journal.

431 [16] Roef MJ, de Meer K, Reijngoud DJ, Straver HW, de Barse M, Kalhan SC, Berger R. Triacylglycerol
432 infusion improves exercise endurance in patients with mitochondrial myopathy due to
433 complex I deficiency. *Am J Clin Nutr* 2002;75: 237-244.

434 [17] Roef MJ, de Meer K, Reijngoud DJ, Straver HW, de Barse M, Kalhan SC, Berger R. Triacylglycerol
435 infusion does not improve hyperlactemia in resting patients with mitochondrial myopathy
436 due to complex I deficiency. *Am J Clin Nutr* 2002;75: 228-236.

- 437 [18] Roef MJ, Reijngoud DJ, Jeneson JA, Berger R, de MK. Resting oxygen consumption and in vivo ADP are
438 increased in myopathy due to complex I deficiency. *Neurology* 2002;58: 1088-1093.
- 439 [19] Panetta J, Smith LJ, Boneh A. Effect of high-dose vitamins, coenzyme Q and high-fat diet in paediatric
440 patients with mitochondrial diseases. *J Inherit Metab Dis* 2004;27: 487-498.
- 441 [20] Bernier FP, Boneh A, Dennett X, Chow CW, Cleary MA, Thorburn DR. Diagnostic criteria for
442 respiratory chain disorders in adults and children. *Neurology* 2002 ;59: 1406-1411.
- 443 [21] Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: Updated guidelines for reporting parallel
444 group randomised trials. *J Pharmacol Pharmacother* 2011 ;1: 100-107.
- 445 [22] Martin A. Apports nutritionnels conseillés pour la population française. TEC et DOC Lavoisier 2000;
446 23-43.
- 447 [23] Belko AZ, Barbieri TF, Wong EC. Effect of energy and protein intake and exercise intensity on the
448 thermic effect of food. *Am J Clin Nutr* 1986 ; 43: 863-869.
- 449 [24] Barnard JA, Tapsell LC, Davies PS, Brenninger VL, Storlien LH. Relationship of high energy expenditure
450 and variation in dietary intake with reporting accuracy on 7 day food records and diet
451 histories in a group of healthy adult volunteers. *Eur J Clin Nutr* 2002;56: 358-367.
- 452 [25] Informatic Center for Quality of foods. Pictures of different meals 1993 ; 1-63.
- 453 [26] Welch AA, McTaggart A, Mulligan AA, Luben R, Walker N, Khaw KT, Day NE, Bingham SA. DINER (Data
454 Into Nutrients for Epidemiological Research) - a new data-entry program for nutritional
455 analysis in the EPIC-Norfolk cohort and the 7-day diary method. *Public Health Nutr* 2001;4:
456 1253-1265.
- 457 [27] Chattaway FW, Happold FC, Happold AM. Nutrition of school-children in Leeds, winter, 1943, and
458 summer, 1944. *Br Med J* 1946;1: 429.
- 459 [28] Dougherty FE. Metabolic testing in mitochondrial disease. *Semin Neurol* 200 ;21: 303-308.
- 460 [29] Blond E, Maitrepierre C, Normand S, Sothier M, Roth H. A new indirect calorimeter is accurate and
461 reliable for measuring basal energy expenditure, thermic effect of food and substrate
462 oxidation in obese and healthy subjects. *e-journal of Clinical nutrition* 2010 ; .
- 463 [30] Weir JB. New methods for calculating metabolic rate with special reference to protein metabolism. *J*
464 *Physiol* 1949;109: 1-9.
- 465 [31] Dabbech M, Boulrier A, Apfelbaum M, Aubert R. Thermic effect of meal and fat mass in lean and obese
466 men. *Nutrition Research* 1996; 16:1133-1141.
- 467 [32] Goran MI, Driscoll P, Johnson R, Nagy TR, Hunter G. Cross-calibration of body-composition techniques
468 against dual-energy X-ray absorptiometry in young children. *Am J Clin Nutr* 1996;63:299-305.
- 469 [33] Putt M, Chinchilli VM. A mixed effects model for the analysis of repeated measures cross-over
470 studies. *Stat Med* 1999 18:3037-3058.
- 471 [34] Chen X, Wei L. A comparison of recent methods for the analysis of small-sample cross-over studies.
472 *Stat Med* 2003;22:2821-2833.
- 473 [35] Benit P, El-Khoury R, Schiff M, Sainsard-Chanet A, Rustin P. Genetic background influences
474 mitochondrial function: modeling mitochondrial disease for therapeutic development. *Trends*
475 *Mol Med* 2010; 16:210-217.

476 [36] Kirby DM, Crawford M, Cleary MA, Dahl HH, Dennett X, Thorburn DR (1999) Respiratory chain
477 complex I deficiency: an underdiagnosed energy generation disorder. *Neurology* 52:1255-
478 1264.

479 [37] Briet F, Twomey C, Jeejeebhoy KN. Effect of feeding malnourished patients for 1 mo on mitochondrial
480 complex I activity and nutritional assessment measurements. *Am J Clin Nutr* 2004;79:787-
481 794.

482 [38] Koene S, Rodenburg RJ, van der Knaap MS, Willemsen MA, Sperl W, Laugel V, Ostergaard E,
483 Tarnopolsky M, Martin MA, Nesbitt V, Fletcher J, Edvardson S, Procaccio V, Slama A, van den
484 Heuvel LP, Smeitink JA. Natural disease course and genotype-phenotype correlations in
485 Complex I deficiency caused by nuclear gene defects: what we learned from 130 cases. *J*
486 *Inherit Metab Dis* 2012;35:737-747.

487 [39] Flatt JP, Ravussin E, Acheson KJ, Jequier E. Effects of dietary fat on postprandial substrate oxidation
488 and on carbohydrate and fat balances. *J Clin Invest* 198 ;76:1019-1024.

489 [40] Acheson KJ, Schutz Y, Bessard T, Ravussin E, Jequier E, Flatt JP. Nutritional influences on lipogenesis
490 and thermogenesis after a carbohydrate meal. *Am J Physiol* 1984;246: E62-E70.

491 [41] de Meer K, Roef MJ, de Klerk JB, Bakker HD, Smit GP, Poll-The BT. Increasing fat in the diet does not
492 improve muscle performance in patients with mitochondrial myopathy due to complex I
493 deficiency. *J Inherit Metab Dis* 2005;28: 95-98.

494 [42] Branco AF, Ferreira A, Simoes RF, Magalhaes-Novais S, Zehowski C, Cope E, Silva AM, Pereira D, Sardao
495 VA, Cunha-Oliviera T. Ketogenic diets : from cancer to mitochondrial diseases and beyond.
496 *Eur J Clin Nutr* 2016 (*in press*)

497

Table 1

Anthropometric characteristics of total included subjects and those undergoing REE and DIT measurements. Results are expressed as mean values $\pm$ SD.

	Group 1 <i>Total included subjects</i>	Group 2 <i>Analysed subjects for REE</i>	Group 3 <i>Analysed subjects for DIT</i>	p-value <i>(group 1 vs 2)</i>	p-value <i>(group 1 vs 3)</i>
n	36	22	12	NA	NA
Female (%)	48	50	60	0.837 [#]	0.916 [∅]
Age : Year	13.2 $\pm$ 4.7	13.2 $\pm$ 4.6	13.8 $\pm$ 4.8	1.000 [§]	0.823 [§]
Weight: kg	36.3 $\pm$ 15.8	36.8 $\pm$ 15.9	40.2 $\pm$ 18.4	0.928 [§]	0.765 [§]
Height: cm	143.8 $\pm$ 22.1	143.4 $\pm$ 20.2	148 $\pm$ 21.3	0.918 [§]	0.811 [§]
BMI: Kg/m²	17.0 $\pm$ 4.1	16.9 $\pm$ 4.2	17.4 $\pm$ 4.6	0.924 [§]	0.875 [§]
Z-score BMI	-0.96 $\pm$ 1.6	-0.94 $\pm$ 1.68	-0.75 $\pm$ 1.59	0.878 [§]	0.749 [§]
Fat mass: %	23.1 $\pm$ 12.9	22.22 $\pm$ 12.3	22.3 $\pm$ 11.5	0.888 [§]	0.902 [§]
[∑] FFM: Kg	28.6 $\pm$ 11.7	23.3 $\pm$ 11.5	32.9 $\pm$ 12.2	0.817 [§]	0.396 [§]
[∑] Muscle: Kg	26.9 $\pm$ 10.3	22.1 $\pm$ 12.4	30.8 $\pm$ 9.4	0.902 [§]	0.753 [§]

BMI = Body Mass Index

NA = Not Applicable

NS = Not Significant

FFM = Fat Free Mass measured by [∑]DEXA

[#] = χ^2 test : [∅] = χ^2 test with Yates correction

[§] = Student *t* test

Table 2

Repartition of respiratory chain complex deficiencies in the group of REE analysed subjects along with their main clinical signs (n=22; group 2). Mitochondrial respiratory chain enzyme activities were assessed by biochemical measurements on skeletal muscle biopsy or on cultured skin fibroblasts; diagnoses of MELAS and MNGIE were made on the basis of molecular studies.

Complex deficiency	n	Neurological signs	Muscular signs	Neuro-sensitive signs
I	12*	7	1	0
III	2**	2	2	1
IV	6	6	0	0
MELAS ¹	1	1	0	0
MNGIE ²	1	1	0	0
TOTAL	22	16	3	1

¹ MELAS = Myoclonic Epilepsy Lactate Acidosis and Stroke like episodes with mutation 3243A>G (mitochondrial DNA).

² MNGIE = MyoNeuroGastro Intestinal Encephalopathy with mutation C130T-G128A.

* 3 of 12 patients are simultaneously deficient for complexes I, II and III.

** 2 patients are simultaneously deficient for complexes III and IV.

Table 3

Comparison between anthropometric characteristics of respiratory chain-deficient subjects with and without complex I deficiencies in the group analysed for REE and body composition. Results are expressed as mean values $\pm$ SD.

	Patients with complex I deficiency	Patients with other defects	p-value
n	12	10	NA
Female (%)	61	56	0.395 $\varnothing$
Age : Year	12.5 $\pm$ 4.5	14.3 $\pm$ 4.9	0.693 $\S$
Weight : kg	39.9 $\pm$ 15.9	33.5 $\pm$ 15.4	0.376 $\S$
Height : cm	142.6 $\pm$ 18.3	144.4 $\pm$ 23.1	0.927 $\S$
Z-score BMI	-0.06 $\pm$ 1.6	-2.0 $\pm$ 1.11	0.010$\S$

BMI = Body Mass Index

NA = Not Applicable

FFM = Fat Free Mass measured by 2 DEXA

$\varnothing$ = χ^2 test with Yates correction

$\S$ = U-Mann Whitney test

Table 4.

Food parameters during cross over periods for REE analysed subjects (n=22 group 2). Results are expressed as mean values $\pm$ SD.

	Cross-over periods			
	Baseline ^a	BD	CD	p-value (BD <i>vs</i> CD)
Energy Intake: Kcal/d	1722.2 $\pm$ 366.1	1729.6 $\pm$ 401.6	1950.0 $\pm$ 593.5	0.089 [§]
Carbohydrates: %	55.0 $\pm$ 4.0	53.8 $\pm$ 4.9	43.1 $\pm$ 3.1	<0.001 [§]
Lipids: %	29.3 $\pm$ 3.6	31.1 $\pm$ 2.9	43.6 $\pm$ 2.5	<0.001 [§]
Proteins: %	15.1 $\pm$ 3.4	15.1 $\pm$ 3.3	13.3 $\pm$ 2.3	0.24 [§]

^a Baseline is during a basal diet period

^b significant

BD = Basal diet : CD = Challenging diet

§ = Paired student *t* test

Table 5.

Metabolic parameters at rest of analysed subjects (n= 22 group 2) at baseline, after basal diet or after challenging diet. Results are expressed as mean values $\pm$ SD.

	Cross-over periods			
	Baseline	BD	CD	p-value (BD vs CD)
VO₂				
VO ₂ ml/min	210.5 $\pm$ 42.5	205.6 $\pm$ 54.2	204.7 $\pm$ 43.8	0.904 [§]
VO ₂ ml/min/kg of body weight	4.5 $\pm$ 2.1	4.8 $\pm$ 2.0	4.8 $\pm$ 2.1	0.899 [§]
VO ₂ ml/min/kg of FFM	5.4 $\pm$ 2.3	6.1 $\pm$ 2.3	6.9 $\pm$ 4.2	0.775 [§]
VO ₂ ml/min/kg of muscle	5.8 $\pm$ 2.2	6.5 $\pm$ 2.3	6.5 $\pm$ 2.3	0.939 [§]
VCO₂				
VCO ₂ ml/min	243.2 $\pm$ 55.5	235 $\pm$ 60.1	240.7 $\pm$ 51.9	0.842 [§]
VCO ₂ ml/min/kg of body weight	5.3 $\pm$ 2.3	4.5 $\pm$ 1.9	4.4 $\pm$ 2.2	0.810 [§]
VCO ₂ ml/min/kg of FFM	6.4 $\pm$ 2.4	5.6 $\pm$ 2.0	6.2 $\pm$ 3.8	0.436 [§]
VCO ₂ ml/min/kg of muscle	6.9 $\pm$ 2.3	6.0 $\pm$ 2.1	5.9 $\pm$ 2.4	0.332 [§]
REE				
REE Kcal/d	1229.6 $\pm$ 303.8	1148.8 $\pm$ 301.7	1156.1 $\pm$ 278.8	0.942 [§]
² pREE Kcal/d	1192.4 $\pm$ 232.4	1191.0 $\pm$ 246.7	1200 $\pm$ 245.2	0.934 [§]
REE Kcal/d/kg of body weight	38.1 $\pm$ 13.7	34.7 $\pm$ 11.8	35.7 $\pm$ 15.1	0.720 [§]
REE Kcal/d/kg of FFM	46.4 $\pm$ 13.7	43.3 $\pm$ 12.3	50.2 $\pm$ 28.9	0.624 [§]
REE Kcal/d/kg of muscle	49.1 $\pm$ 13.0	49.9 $\pm$ 12.4	48.1 $\pm$ 16.2	0.834 [§]
RQ				
mRQ	0.86 $\pm$ 0.10	0.87 $\pm$ 0.08	0.85 $\pm$ 0.10	0.834 [§]
pRQ	0.87 $\pm$ 0.11	0.88 $\pm$ 0.10	0.84 $\pm$ 0.10	0.729 [§]

^a Baseline is during a basal diet period

^b significant

BD = Basal diet; CD = Challenging diet

REE = Resting Energy Expenditure

pREE = Predictive Resting Energy Expenditure

FFM = Fat Free Mass

mRQ = Measured Respiratory Quotient (measured VO₂/measured VCO₂)

pRQ = Predictive Respiratory Quotient (theoretically calculated from the nutritional contents of diets in each class of energetic substrates [carbohydrates, lipids and proteins])

§ = Paired student *t* test

§ = Paired student *t* test

²Predictive REE calculated by Harris and Benedict equation

Table 6.

Body composition parameters of subjects (n= 22) undergoing REE evaluations at baseline, after one month of basal diet and after one month of challenging diet. Results are expressed as mean values $\pm$ SD.

	Cross-over periods			p-value (BD <i>vs</i> CD)
	Baseline	BD	CD	
Weight : Kg	38.8 $\pm$ 15.9	36.6 $\pm$ 13.7	38.5 $\pm$ 18.0	0.698 [§]
Fat mass : %	22.1 $\pm$ 12.3	21.9 $\pm$ 13.0	21.6 $\pm$ 13.3	0.810 [§]
FFM : Kg	29.3 $\pm$ 11.5	29.1 $\pm$ 12.2	26.8 $\pm$ 11.0	0.716 [§]
Muscle : Kg	26.9 $\pm$ 10.3	26.6 $\pm$ 11.1	26.5 $\pm$ 11.0	0.810 [§]

FFM = Fat Free Mass

§ = Paired student *t* test

^a Baseline is during a basal diet period

^b significant

BD = Basal diet : CD = Challenging diet

Table 7.

Minimum set of metabolic and anthropometrics data in complex I deficient subjects (n= 12) undergoing REE evaluations at baseline, after one month of basal diet and after one month of challenging diet. Results are expressed as mean values $\pm$ SD.

	Cross-over periods			p-value (BD vs CD)
	Baseline	BD	CD	
REE <i>Kcal/d</i>	1229.6 $\pm$ 258.2	1139.5 $\pm$ 250.3	1158.2 $\pm$ 255.5	0.695 §
RQ	0.75 $\pm$ 0.12	0.88 $\pm$ 0.14	0.82 $\pm$ 0.08	0.366 §
Weight: <i>Kg</i>	39.9 $\pm$ 15.9	40.2 $\pm$ 16.2	40.6 $\pm$ 16.1	0.948 §
Fat mass: %	27.9 $\pm$ 14.5	25.1 $\pm$ 15.9	26.1 $\pm$ 16.5	0.790 §
FFM: <i>Kg</i>	29.8 $\pm$ 8.9	28.9 $\pm$ 9.8	28.3 $\pm$ 9.8	0.657 §
Muscle: <i>Kg</i>	26.9 $\pm$ 8.2	26.5 $\pm$ 9.3	25.8 $\pm$ 9.1	0.929 §

FFM = Fat Free Mass

§ = Wilcoxon test

^a Baseline is during a basal diet period

^b significant

BD = Basal diet; CD = Challenging diet

Table 8.

Minimum set of metabolic and anthropometrics data of subjects with a respiratory chain defect other than complex I (n= 10) and undergoing REE evaluations at baseline, after one month of basal diet and after one month of challenging diet. Results are expressed as mean values $\pm$ SD.

	Cross-over periods			p-value (BD vs CD)
	Baseline	BD	CD	
REE <i>Kcal/d</i>	1195.1 $\pm$ 361.2	1166.5 $\pm$ 378.5	1157.0 $\pm$ 316.3	0.875 §
RQ	0.80 $\pm$ 0.15	0.81 $\pm$ 0.16	0.89 $\pm$ 0.12	0.555 §
Weight: <i>Kg</i>	33.5 $\pm$ 15.4	34.5 $\pm$ 15.5	34.5 $\pm$ 15.4	0.948 §
Fat mass: %	15.8 $\pm$ 4.6	17.5 $\pm$ 6.4	16.6 $\pm$ 6.1	0.109 §
FFM: <i>Kg</i>	28.7 $\pm$ 14.3	29.2 $\pm$ 15.5	25.2 $\pm$ 12.6	0.248 §
Muscle: <i>Kg</i>	26.7 $\pm$ 12.7	26.8 $\pm$ 13.7	27.3 $\pm$ 13.2	0.185 §

FFM = Fat Free Mass

§ = Wilcoxon test

^a Baseline is during a basal diet period

^b significant

BD = Basal diet; CD = Challenging diet

Table 9.

Food parameters during cross over periods for DIT assessment (n = 12 ; group 3). Results are expressed as mean values $\pm$ SD.

	Cross-over periods		p-value (BD vs CD)
	BD	CD	
Energy Intake*: <i>Kcal</i>	161.0 $\pm$ 12.3	219.9 $\pm$ 13.0	0.809 [§]
Carbohydrates: %	61.1 $\pm$ 6.6	45.5 $\pm$ 12.2	0.403 [§]
Lipids: %	28.5 $\pm$ 10.3	42.3 $\pm$ 11.1	0.878 [§]
Proteins: %	10.4 $\pm$ 6.4	12.2 $\pm$ 5.9	0.911 [§]

*Energy intake at breakfast for DIT measurement

§ = Wilcoxon test

BD = Basal diet

CD = Challenging diet

Figure 1 : Flow CONSORT diagram of screening, inclusion, randomisation and analysis rate in the study.

Figure 2 : Randomized cross-over design of the study.

Figure 3 : Evolution of diet induced thermogenesis (only group 3 ; n=12)

Figure 4 : Evolution of RQ during diet induced thermogenesis assessment (only group 3 ; n=12)

Figure 1 : Flow CONSORT diagram of screening, inclusion, randomisation and analysis rate in the study.

BD = Basal Diet
CD = Challenging Diet

Figure 2 : Randomized cross-over design of the study.

Figure 3 : Evolution of diet induced thermogenesis (only Group 3 ; n=12)

Figure 4 : Evolution of RQ during diet induced thermogenesis assessment (only group 3 ; n=12)