



HAL
open science

Outcome of EGFR-mutated NSCLC patients with MET-driven resistance to EGFR tyrosine kinase inhibitors

Simon Baldacci, Julien Mazieres, Pascale Tomasini, Nicolas Girard, Florian Guisier, Clarisse Audigier-Valette, Isabelle Monnet, Marie Wislez, Maurice Perol, Pascal Do, et al.

► **To cite this version:**

Simon Baldacci, Julien Mazieres, Pascale Tomasini, Nicolas Girard, Florian Guisier, et al.. Outcome of EGFR-mutated NSCLC patients with MET-driven resistance to EGFR tyrosine kinase inhibitors. *Oncotarget*, 2017, 8 (62), pp.105103-105114. 10.18632/oncotarget.21707 . hal-02543375

HAL Id: hal-02543375

<https://hal.univ-lille.fr/hal-02543375v1>

Submitted on 15 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Outcome of EGFR-mutated NSCLC patients with MET-driven resistance to EGFR tyrosine kinase inhibitors

Simon Baldacci^{1,15}, Julien Mazieres², Pascale Tomasini³, Nicolas Girard⁴, Florian Guisier⁵, Clarisse Audigier-Valette⁶, Isabelle Monnet⁷, Marie Wislez⁸, Maurice Pérol⁹, Pascal Dô¹⁰, Eric Dansin¹¹, Charlotte Leduc¹², Etienne Giroux Leprieur¹³, Denis Moro-Sibilot¹⁴, David Tulasne¹⁵, Zoulika Kherrouche^{1,15}, Julien Labreuche¹⁶ and Alexis B. Cortot^{1,15}

¹CHU Lille, Thoracic Oncology Department, Univ. Lille, Siric ONCOLille, Lille, France

²Toulouse University Hospital, Université Paul Sabatier, Toulouse, France

³Aix-Marseille University, Assistance Publique Hôpitaux de Marseille, Multidisciplinary Oncology & Therapeutic Innovations Department, Marseille, France

⁴Louis Pradel Hospital, Hospices Civils de Lyon, Lyon, France

⁵Rouen University Hospital, Thoracic oncology unit & Normandy University, IRIB, LITIS Lab, EA 4103 QuantIF team, Rouen, France

⁶Service de Pneumologie, Centre Hospitalier Sainte Musse, Toulon, France

⁷Centre Hospitalier Intercommunal de Créteil, Créteil, France

⁸APHP Hôpital Tenon, Paris, France

⁹Department of Medical Oncology, Centre Léon Bérard, Lyon, France

¹⁰Centre Régional de Lutte Contre le Cancer François Baclesse, Caen, France

¹¹Centre Oscar Lambret, Lille, France

¹²CHU Strasbourg, Strasbourg, France

¹³APHP – Hôpital Ambroise Paré, Boulogne-Billancourt, France

¹⁴Unité d'Oncologie Thoracique, Service de Pneumologie, CHU Grenoble-Alpes, La Tronche, France

¹⁵Univ. Lille, CNRS, Institut Pasteur de Lille, UMR 8161, M3T, Mechanisms of Tumorigenesis and Targeted Therapies, Lille, France

¹⁶EA 2694 University of Lille, Lille, France

Correspondence to: Alexis B. Cortot, **email:** alexis.cortot@chru-lille.fr

Keywords: non small cell lung cancer; EGFR; tyrosine kinase inhibitors; resistance; MET

Received: February 21, 2017

Accepted: August 04, 2017

Published: October 09, 2017

Copyright: Baldacci et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Background: Several mechanisms of acquired resistance to EGFR tyrosine kinase inhibitors (TKIs) in EGFR-mutated NSCLC have been described including the T790M mutation and *MET* amplification. Whereas T790M mutation confers prolonged survival and sensitivity to 3rd generation TKIs, data are lacking on clinical features and outcome of MET-driven resistant EGFR-mutated NSCLC patients.

Methods: Patients with metastatic EGFR-mutated NSCLC displaying high *MET* overexpression or *MET* amplification, detected on a biopsy performed after progression on EGFR TKI, were identified in 15 centers. Clinical and molecular data were retrospectively collected.

Results: Forty two patients were included. The median overall survival (OS), and the median post EGFR TKI progression overall survival (PPOS) were 36.2 months

[95%CI 27.3-66.5] and 18.5 months [95%CI 10.6-27.4] respectively. Nineteen out of 36 tumors tested for MET FISH had MET amplification. A T790M mutation was found in 11/41 (26.8%) patients. T790M-positive patients had a better OS than T790M-negative patients (p=0.0224). Nineteen patients received a MET TKI. Objective response was reported in 1 out of 12 evaluable patients treated with a MET inhibitor as a single agent and in 1 of 2 patients treated with a combination of MET and EGFR TKIs.

Conclusion: MET-driven resistance to EGFR TKI defines a specific pattern of resistance characterized by low objective response rate to MET inhibitors given alone and overlapping with T790M mutations. Further studies are warranted to define adequate therapeutic strategies for MET-driven resistance to EGFR TKI.

INTRODUCTION

EGFR mutations are found in 10% of non small cell lung cancer (NSCLC) in Caucasians and 40% in Asians [1]. Treatment of advanced *EGFR*-mutated NSCLC patients relies on EGFR tyrosine kinase inhibitors (TKIs), which demonstrated superiority over chemotherapy as first-line therapy [2–5]. However, despite initial efficacy, all the patients will eventually develop resistance to EGFR TKIs resulting in tumor progression [6]. The most frequent mechanism of resistance is the T790M mutation, a second *EGFR* mutation that can be successfully targeted with third generation EGFR TKIs which have been specifically designed to overcome T790M-driven resistance [7, 8]

Besides T790M mutation, bypass activation of other tyrosine kinase receptors including MET or HER2 is the second most common mechanism of resistance to EGFR TKI. *MET* amplification leads to overexpression and constitutive activation of the receptor, thus activating the PI3K pathway and bypassing EGFR [9]. *MET* amplification has been detected in 5 to 22% of patients with an acquired resistance to EGFR TKI [9–14]. *MET* amplification is highly associated with high overexpression of MET in NSCLC (p<0.001) [15, 16]. Moreover, high MET overexpression with a 3+ immunoscore (IHC3+) by immunohistochemistry was recently found in 27% of *EGFR* mutated NSCLC with acquired resistance to EGFR TKI [17].

Little data is available about clinical characteristics and outcome of *EGFR*-mutated NSCLC patients with MET-driven resistance to EGFR TKIs. Moreover, optimal treatment of these patients is still unknown. Preclinical data and case reports suggest that *MET*-amplified *EGFR*-mutated NSCLC are addicted to both *MET* and *EGFR* and that combination of MET and EGFR TKIs is required to overcome this mechanism of resistance [9, 18, 19].

Whereas available data and specific treatments emerge for T790M-driven resistance in *EGFR*-mutated NSCLC patients, MET-driven resistance is still an unexplored field [20–22]. In the present study, we report clinical features, outcome and treatment in a series of *EGFR*-mutated NSCLC patients with MET-driven resistance to EGFR TKI.

RESULTS

Clinico-pathological and molecular characteristics

Forty six patients with metastatic NSCLC displaying both *EGFR* mutation and MET overexpression or *MET* gene amplification were retrospectively identified in 15 centers. Four patients were excluded : 3 had only a biopsy performed before EGFR TKI initiation, and 1 had no data available on the treatments received. Forty two patients were included. Re-biopsies of these patients had been performed from May 2011 to May 2016.

Patient clinical characteristics are summarized in Table 1. All 42 patients were diagnosed with metastatic lung adenocarcinoma. Median age was 65.1 years (range 30-82.7). The majority of patients were women (66.7%) and never smokers (70.7%). Most of the *EGFR* mutations detected on the initial biopsy were exon 19 deletions or exon 21 L858R point mutations.

The most frequent site of rebiopsy was the lung and re-biopsies were performed, in 90.5% of the cases, after the Response Evaluation Criteria in Solid Tumor (RECIST) progression on EGFR TKI, which was given as first or second line treatment (Supplementary Table 1). The median time between EGFR TKI initiation and re-biopsy was 15.6 months (range 2.1-61.3).

Among the 42 patients included in the study, 36 tumor re-biopsy samples were tested for *MET* FISH and 19 (52.8%) were found *MET* amplified (Figure 1). MET IHC was performed on the re-biopsy of 36 patients and all displayed a high level of MET expression (IHC3+). Six patients had a MET FISH but no MET IHC on their re-biopsy and conversely 6 other patients had MET IHC and no interpretable MET FISH. No difference was found regarding the *MET* amplified status between patients with an EGFR exon 19 deletion or an EGFR exon 21 mutation on the initial biopsy (Supplementary Table 2). Re-biopsies of 34 patients were tested for the BRAF mutations and none harbored a mutation of this oncogene. Otherwise no histological transformation in small cell lung cancer was reported among the 42 patients of the study.

Table 1: Patient characteristics

	Overall population	<i>MET</i> amplification	<i>MET</i> overexpression no <i>MET</i> amplification	p	T790M+	T790M-	p
	n = 42	n = 19	n = 17		n = 11	n = 30	
Median age (years)	65,1 (30-82,7)*	64,6 (30-74,6)*	65,6 (38,4-82,7)*	0,73	56,9 (45,2-70,3)*	67,4 (30-82,7)*	0,038
Gender				0,35			0,28
Men	14 (33,3%)	5 (26,3%)	7 (41,2%)	2 (18,2%)	12 (40%)		
Women	28 (66,7%)	14 (73,7%)	10 (58,8%)	9 (81,8%)	18 (60%)		
Smoking status[†]				0,45			1
Never smoker	29 (70,7%)	12 (63,2%)	12 (75%)		8 (72,7%)	20 (69%)	
Former and current smoker	12 (29,3%)	7 (36,8%)	4 (25%)	3 (27,3%)	9 (31%)		
Stade IV	42 (100%)	19 (100%)	17 (100%)		11 (100%)	30 (100%)	
Adenocarcinoma	42 (100%)	19 (100%)	17 (100%)		11 (100%)	30 (100%)	
Initial EGFR mutation				0,59			0,55
Exon 19 deletion or Exon 21 L858R mutation	39 (92,9%)	18 (94,7%)	15 (88,2%)		11 (100%)	27 (90%)	
Other**	3 (7,1%)	1 (5,3%)	2 (11,8%)		0 (0%)	3 (10%)	

* : range ; ** : 1 exon 20 S768I mutation, 1 association of exon 21 L858R and K860I mutations, 1 association of exon 19 R761Y and exon 18 G719A mutations ; † : one missing data in total population ; TKI : tyrosine kinase inhibitor ; p : p value.

Clinical outcome and EGFR TKI treatment characteristics

Characteristics of initial EGFR TKI therapy are shown in Table 2. All patients received a first or second generation EGFR TKI. The overall response rate (ORR) was 82.1%, and the median progression free survival (PFS) was 11.1 months [95%CI 7.6-14.1]. In 73.2% of the cases, tumor progression involved a new lesion. The main site of new metastasis was the lung. 27 patients (66%) developed more than one progressive lesion at EGFR TKI resistance. The median post-progression overall survival (PPOS) and the median overall survival (OS) were respectively 18.5 months [95%CI 10.6-27.4] and 36.2 months [95%CI 27.3-66.5].

The characteristics of the EGFR TKI therapy were not significantly different according to the MET FISH status. No significant difference was found between patients with *MET* amplification and those with MET overexpression and no *MET* amplification in terms of OS (median OS : 42.8 vs. 36.2 months p = 0.69 ; Figure 2A), PPOS (median PPOS : 13.7 vs. 23.8 months p =0.77) and PFS on EGFR TKI (median PFS : 10.5 vs. 10.1 months p = 0.08). There were also no significant differences between the patients with EGFR exon 19 deletion on the initial biopsy and the patients with EGFR exon 21 mutation in terms of OS (median OS :

36.2 vs. 27.3 months p = 0.33), PPOS (median PPOS 18.5 vs. 12.6 months p = 0.91), and PFS on EGFR TKI (median PFS : 11.7 vs. 9.2 months p = 0.56). Of note only one patient with an EGFR exon 19 deletion received afatinib, and this drug was not used in patients with EGFR exon 21 mutation (Supplementary Table 2).

Impact of the T790M mutational status

Eleven among the 41 patients (26.8%) tested for the T790M mutation were T790M positive. In 9 patients, the T790M mutation was detected in the tumor re-biopsy displaying MET overexpression or *MET* amplification. In the remaining 2 patients, the T790M mutation was detected in circulating free DNA in one case and in a tumor biopsy obtained before the re-biopsy displaying MET overexpression and *MET* amplification in another case. Three patients had both *MET* amplification and T790M mutation in the same sample. The T790M-positive patients were significantly younger than T790M-negative patients and the time between EGFR TKI initiation and re-biopsy was significantly longer in T790M positive patients (Table 1 and Supplementary Table 1).

T790M-positive patients had a better OS (median OS 43.1 vs. 32.2 months, p=0.0224) than T790M-negative patients (Figure 2B). There was also a trend to

Table 2: EGFR TKI treatment characteristics

	Overall population	<i>MET</i> amplification	<i>MET</i> overexpression no <i>MET</i> amplification	p	T790M+	T790M-	p
	n = 42	n = 19	n = 17		n = 11	n = 30	
First EGFR TKI received				0,49			1
Erlotinib or Gefinitib	40 (95,2%)	17 (89,5%)	17 (100%)		11 (100%)	28 (93,3%)	
Afatinib	2 (4,76%)	2 (10,5%)	0 (0%)		0 (0%)	2 (6,7%)	
Line of the first EGFR TKI therapy				0,81			1
1	29 (69%)	13 (68,4%)	11 (64,7%)		8 (72,7%)	20 (66,7%)	
2	13 (31%)	6 (31,6%)	6 (35,3%)		3 (27,3%)	10 (33,3%)	
Response to the EGFR TKI †				1			0,17
Objective response	32 (82,1%)	14 (77,8%)	12 (80%)		10 (100%)	22 (78,6%)	
Stable disease and Progression	7 (17,9%)	4 (22,2%)	3 (20%)		0 (0%)	6 (21,4%)	
Median duration of TKI EGFR therapy (months)‡	13,1 (1,4-52,5)*	13,8 (4,1-21,6)*	10,2 (1,4-45,4)*	0,61	14,0 (7,7-25)*	10,7 (1,4-52,5)*	0,51
EGFR TKI progression involving a new metastasis ‡	30 (73,2%)	14 (77,8%)	12 (70,6%)	0,71	8 (80%)	22 (73,3%)	1
EGFR TKI progression involving more than one progressive lesion ‡	27 (65,8%)	12 (63,2%)	11 (68,8%)	1	10 (90,9%)	16 (55,2%)	0,065

* : range ; TKI :tyrosine kinase inhibitor ; p : p value ; † : 3 missing data ; ‡ : one missing data.




Figure 1: Flow chart of MET overexpression and MET amplification status on post EGFR TKI initiation sample. IHC : Immunohistochemistry ; FISH : Fluorescence In Situ Hybridization ; TKI : tyrosine kinase inhibitor.

a better PPOS in T790M-positive patients compared to the T790M-negative patients although it did not reach statistical significance (median 23.8 vs. 11.0 months, $p=0.075$, Supplementary Table 3).

Treatment with MET inhibitors and third generation EGFR TKI therapy

Nineteen patients received a MET inhibitor, mostly crizotinib, including 13 *MET*-amplified patients (Table 3). The MET inhibitor was given as a monotherapy in 15 patients and in combination with an EGFR TKI in 4 patients. The

MET inhibitor was used as 2nd or 3rd line in most of the cases. Objective response was reported in 1 out of 12 evaluable patients treated with a MET inhibitor as single agent and in 1 out of 2 evaluable patients treated with a combination of MET and EGFR inhibitors (Figure 3A and 3B). The MET inhibitor was stopped because of elevated liver enzymes in 2 patients and diarrhea in one patient. Two of these patients received the MET inhibitor in combination with an EGFR TKI. The median times between re-biopsy and MET inhibitor initiation and first EGFR TKI withdrawal and MET inhibitor initiation were respectively 2.3 months and 3.1 months.




Figure 2: Overall survival according to T790M status and MET amplification status. (A) Kaplan Meier estimates of overall survival in *MET* non amplified (blue) and *MET* amplified (red) patients. **(B)** Kaplan Meier estimates of overall survival in T790M-negative (blue) and T790M-positive (red) patients.

Ten patients received a 3rd generation EGFR TKI (Table 4) mostly osimertinib. Two partial responses were reported among the 5 T790M-positive evaluable patients who were treated following detection of MET overexpression or *MET* amplification. 3 patients were treated with 3rd generation EGFR TKI before they received a MET inhibitor.

DISCUSSION

Bypass activation of tyrosine kinase receptors is a well described mechanism of resistance to EGFR TKIs in *EGFR*-mutated NSCLC. MET-driven resistance has been reported in up to 22% of patients with acquired resistance to EGFR TKIs [9]. Still, although the biological basis of this mechanism of resistance has been extensively studied, there is very little data on the clinical characteristics and outcome of patients with MET-driven acquired resistance to EGFR TKIs. In this multicenter retrospective study, we report for the first time clinical features, response to

MET inhibitors and outcome of 42 metastatic *EGFR*-mutated NSCLC patients with *MET* amplification or MET overexpression, as assessed on a post-progression re-biopsy. Because of the very low number of patients with MET-driven resistance to EGFR TKIs, we performed a multicentric retrospective study. Therefore, we could not perform a central analysis of RECIST responses, *MET* FISH, and MET IHC analysis. The absence of *MET* FISH and MET IHC analysis on the initial biopsy is also challenging. Indeed, we cannot exclude that some patients might have a MET overexpression or a *MET* amplification before EGFR TKI therapy. Most patients with EGFR exon 19 deletions in our study received a first generation EGFR TKI and were not treated with afatinib. There were no specific clinical characteristics of the patients included in our study but we observed a low rate of objective response to MET inhibitors when used as a monotherapy and a substantial rate of concomitant T790M mutations, which were still partially associated




Figure 3: Duration of treatment with MET inhibitors. (A) Patients treated with a MET inhibitor as a single agent; (B) Patients treated with a MET inhibitor in combination with an EGFR TKI. ID : identification number, Red: progressive disease as best response according to RECIST criteria ; Orange : stable disease as best response according to RECIST criteria ; Green : partial response as best response according to RECIST criteria, White : no tumor response evaluation available ; Arrow indicates that the MET inhibitor therapy is still ongoing.

Table 3: MET inhibitor tumor response

ID	MET amplification	IHC MET 3+	T790M	MET inhibitor	EGFR TKI in combination therapy	Line	RECIST Response	MET inhibitor status	Duration of MET inhibitor (days)
1	-	+	-	crizotinib	gefitinib	2	-	stopped (toxicity)	10
2	+	+	-	crizotinib	-	5	PD	stopped (PD)	37
4	+		-	crizotinib	-	4	-	stopped (patient's decision)	2
5	+		+	crizotinib	-	5	SD	ongoing	107
6	+	+	-	crizotinib	-	2	SD	ongoing	15
7	+	+	-	crizotinib	-	3	PD	stopped (PD)	20
9	+	+	-	crizotinib	-	2	-	stopped (toxicity)	9
10	+	+	-	crizotinib	-	3	PD	stopped (PD)	59
12	-	+	-	other	other	2	SD	ongoing	49
14	+		-	crizotinib	-	2	SD	ongoing	42
18	+		+	crizotinib	-	4	SD	stopped (PD)	111
19	+	+		crizotinib	-	4	PD	stopped (PD)	55
20	-	+	-	other	-	6	SD	stopped (PD)	119
25	+		-	crizotinib	-	2	PR	ongoing	41
32	-	+	-	other	-	3	PD	stopped (PD)	22
37	+	+	-	other	other	3	PR	stopped (PD)	145
42	-	+	-	crizotinib	-	6	SD	stopped (PD)	60
43		+	-	crizotinib	gefitinib	4	-	stopped (toxicity)	28
45	+	+	+	crizotinib	-	4	-	ongoing	15

ID : identification number ; IHC : Immunochemistry ; PD : progressive disease ; SD : stable disease ; PR : partial response ; TKI : tyrosine kinase inhibitor ; RECIST : response evaluation criteria in solid tumor.

Table 4: Third generation EGFR TKI tumor response

ID	MET amplification	IHC MET 3+	T790M	Re-biopsy performed before 3 rd G EGFR TKI therapy	3 rd G EGFR TKI	Line	RECIST Response	3 rd G EGFR TKI status	Duration of 3 rd G EGFR TKI (days)
5	+		+	+	osimertinib	3	PD	stopped (PD)	48
16	-	+	+	+	other	2	PR	stopped (toxicity)	56
18	+		+	+	osimertinib	5	PD	stopped (PD)	80
20	-	+	-	+	osimertinib	5	PR	stopped (PD)	157
22		+	+	+	osimertinib	2	PR	ongoing	315
26	-	+	+	+	osimertinib	3	PD	stopped (PD)	84
27		+	+	+	osimertinib	4	-	ongoing	15
28	+	+	+	+	osimertinib	2	-	ongoing	18
45	+	+	+	-	osimertinib	3	SD	stopped (PD)	288
46	+	+	+	-	osimertinib	4	PR	ongoing	277

ID : identification number ; IHC : Immunochemistry ; TKI : tyrosine kinase inhibitor ; RECIST : response evaluation criteria in solid tumor ; 3rd G EGFR TKI : Third Generation EGFR TKI.

with efficacy of 3rd generation EGFR TKIs and favorable prognosis, even in association with MET overexpression or *MET* amplification.

In our study, *MET* amplification was defined using the criteria provided by Schildhaus et al. to define high-level *MET* amplification, i.e. as an average *MET* gene copy number (GCN) per cell ≥ 6 or a ratio *MET*/CEP7 ≥ 2 or the presence of *MET* clusters [15]. Indeed these criteria were already widely used across our pathological centers to define positive *MET* FISH. However, many other definitions of *MET* amplification based on FISH have been proposed based on various thresholds for *MET*/CEP7 ratio or mean *MET* GNC [21, 23–27]. Unlike the mean *MET* GNC, the ratio *MET*/CEP7 is thought to discriminate real amplification from polysomia. Recent data showed that a *MET*/CEP7 ratio >5 was able to discriminate lung adenocarcinoma with no other driver mutations and was associated with high objective response rate to crizotinib [27, 28]. However, the right definition of a positive FISH that would allow identification of *MET* gene amplification remains to be determined. The ability to detect *MET* amplification through next generation sequencing may favor routine screening and harmonization of the definition of *MET* amplification.

In our study, *EGFR* mutated NSCLC patients with either *MET* amplification or *MET* overexpression (*MET* IHC3+) were included. In lung adenocarcinoma, *MET* amplification is significantly associated with IHC3+ *MET* overexpression [16]. In our study, all the patients with a *MET* amplification who also underwent a *MET* IHC were scored 3+. *MET* overexpression, regardless *MET* amplification status, has been found to induce addiction to the *MET* pathway, and was recently found in 27.1% of *EGFR* mutated NSCLC with acquired resistance to EGFR TKI [29] [17]. Moreover, an ongoing clinical trial evaluating the efficacy of combining *MET* and EGFR TKIs in *EGFR*-mutated NSCLC patients with *MET*-driven resistance to EGFR TKIs includes patients with both *MET* amplification or *MET* overexpression (NCT01610336). In patients without *MET* amplification, the cause of *MET* overexpression at the time of re-biopsy may involve other molecular alterations including *MET* exon 14 splicing sites mutations which have been reported to be associated with *MET* overexpression [30] and may be involved in resistance to EGFR TKI [31]. Because of the unavailability of most of the tumor samples, we could not test for other genomic alterations.

In our cohort, metastatic *EGFR* mutated NSCLC with a *MET*-driven resistance to EGFR TKI did not display specific clinical features compared to those observed in previous studies. Median PPOS (18.5 months) was also in the range of what has been reported in previous studies focusing on patients with acquired resistance to EGFR TKIs (14.3–20 months) [13, 32, 33]. Moreover, although *MET* GCN alterations and *MET* overexpression have been associated with poor prognosis in resected NSCLC [34, 35], we did

not find poor outcome for the patients included in our study. These findings suggest that the prognostic impact of *MET* activation might depend on the stage of the disease and on the oncogenic environment. We observed only one objective response out of 12 patients treated with *MET* inhibitor monotherapy. In preclinical models, a double inhibition of EGFR and *MET* pathways was required to overcome *MET* driven resistance to EGFR TKI [9, 36]. Cases of *EGFR* mutated *MET* amplified NSCLC responding to combined *MET* and EGFR TKIs have been reported [18, 19]. This dual inhibition approach is currently evaluated in several phase I/II clinical trials in this setting using various EGFR (gefitinib, erlotinib, EGF816, osimertinib) and *MET* (capmatinib, volitinib, tepotinib) TKIs (NCT02468661, NCT01610336, NCT02335944, NCT02374645, NCT02143466, NCT01982955). Preliminary results from a phase II study, evaluating the combination of capmatinib and gefitinib in *EGFR* mutated NSCLC patients who progressed on EGFR TKI, reported an ORR of 19% in patients with *MET* IHC3+ or *MET* IHC $\geq 2+$ and *MET* GCN ≥ 5 and raised to 30% in the subgroup of patients with *MET* GCN ≥ 6 [37].

In our study, 11 patients were T790M positive. Three had both a *MET* amplification and a T790M mutation on the same sample. The coexistence of these mechanisms of resistance has been previously described [10]. Compared to the T790M negative, the T790M positive patients in our study had a better OS and a trend to a better PPOS. Several studies reported that, amongst patients with EGFR TKI acquired resistance, T790M-positive patients had a better PPOS than T790M-negative patients [13, 32, 38]. However, none of these studies evaluated the impact of concomitant *MET* amplification and overexpression. Of note, Gou *et al.* found that patients with a T790M mutation and *MET* overexpression had a worse PPOS than patients with only *MET* overexpression or the T790M mutation alone. This discrepancy with our results might be due to the small size of both cohorts, and variations in the characteristics of the study population (higher rate of never smokers and caucasians in our study). We cannot also exclude the influence of confounding factors such as the younger age and the use of 3rd generation EGFR TKI in T790M positive patients. Indeed in our study, 9 T790M-positive patients received a 3rd generation EGFR TKI and 3 partial responses were achieved. Very few data are available regarding the efficacy of 3rd generation TKIs in the context of multiple resistance mechanisms. Our results suggest that 3rd generation TKIs may retain activity against T790M-positive tumors in some patients, even in the presence of *MET* activation, which may be due to the spatial heterogeneity of resistance mechanisms rather than co-existence of 2 resistance mechanisms in the same tumor cells.

Finally, *MET*-driven resistance to EGFR TKI defines a specific pattern of resistance characterized by low objective response rate to *MET* TKIs given alone and overlapping with T790M mutations. Even when associated with *MET* dysregulation, the T790M mutation

was still associated with relative efficacy of 3rd generation EGFR TKIs and prolonged survival. Further studies are warranted to define adequate therapeutic strategies for MET-driven resistance to EGFR TKI.

MATERIALS AND METHODS

Patients

We constituted a multicenter retrospective observational cohort of patients identified among 15 French centers. Inclusion criteria were documented diagnosis of metastatic NSCLC, detection of an *EGFR* mutation on tumor sample at diagnosis, treatment with at least one EGFR TKI and detection of MET overexpression or *MET* amplification after the time of clinical or radiological progression on EGFR TKI. Post-progression re-biopsy and MET status assessment were routinely performed in participating centers during the study period. Clinical and pathological data were retrospectively collected in each center for all included patients. Best overall response, defined as the best response from the start of treatment until disease progression, was assessed by investigators from available follow-up exams in each center using Response Evaluation Criteria in Solid Tumor (RECIST) v1.1. Crizotinib, gefinitib, and osimertinib were respectively prescribed with the following doses and schedules : 250mg bid, 250mg daily, 80mg daily. The study was approved by a national ethic committee (CEPRO 2016-001).

Histological and molecular analyses

Histological and molecular analyses of formalin-fixed paraffin-embedded (FFPE) samples were prospectively assessed by local pathologists in accredited and quality controlled laboratories, as part of the routine procedure. MET immunohistochemistry and *MET* FISH analyses were realized according to locally certified and nationally approved procedures. MET overexpression was defined as a 3+ MET Immunoreactivity (≥ 50% of tumor cells showing high-intensity staining) on MET immunohistochemistry using MET monoclonal antibody (clone SP44 Ventana). *MET* amplification was defined by Fluorescence In Situ Hybridization (FISH) as a mean *MET* GNC per cell ≥ 6, or a Ratio MET/CEP7 ≥ 2, or the presence of MET clusters [15]. Patients were considered “T790M positive” if an *EGFR* T790M mutation was detected in post-progression circulating tumor DNA, on the re-biopsy on which MET overexpression or *MET* amplification was found, or on a tumor sample collected before the re-biopsy.

Statistical analyses

Categorical variables are expressed as frequencies and percentages. Quantitative variables are expressed

as medians (range). Normality of distribution was assessed graphically and by using the Shapiro–Wilk test. Bivariate analyses were realized to assess the sub-group comparability (*MET* amplified vs. *MET* non-amplified and T790M+ vs. T790M-). Chi-Squared tests or Fisher’s exact test (when expected cell frequency <5) were used to study the association between categorical variables and different groups. A Mann-Whitney U test was used to compare age, time between EGFR TKI initiation and rebiopsy, and duration of TKI EGFR therapy between groups. We estimated and compared overall survival (OS), post progression overall survival (PPOS) and progression free survival (PFS) between the study groups (*MET* amplified vs. *MET* non amplified and T790M+ vs. T790M-) using the Kaplan-Meier Method and log-rank test. OS was measured from the date of metastatic NSCLC diagnosis to the date of death from any cause or last follow-up. The PPOS was measured as the time from EGFR TKI RECIST progression to death from any cause. The PFS was defined as the time from treatment start to disease progression or death from any cause. Objective response rate (ORR) was defined as the percentage of patients with partial or complete response to the indicated treatment. Statistical testing was conducted at the 2-tailed α level of 0.05. Data were analyzed with SAS software version 9.3 (SAS Institute, Cary, NC).

Abbreviations

Non small cell lung cancer (NSCLC), Tyrosine kinase inhibitor (TKI), overall survival (OS), post progression overall survival (PPOS), progression free survival (PFS), objective response rate (ORR), response evaluation criteria in solid tumor (RECIST), gene copy number (GNC), immunohistochemistry (IHC), formalin-fixed paraffin-embedded (FFPE), Fluorescence In Situ Hybridization (FISH).

Authors contributions

Simon Baldacci : study design and conception, acquisition of data, analysis and interpretation of data, drafting manuscript

Julien Mazieres : acquisition of data, critical revision

Pascale Tomasini : acquisition of data, critical revision

Nicolas Girard : acquisition of data, critical revision

Florian Guisier : acquisition of data, critical revision

Clarisse Audigier-Valette : acquisition of data, critical revision

Isabelle Monnet : acquisition of data, critical revision

Marie Wislez : acquisition of data, critical revision

Maurice Pérol : acquisition of data, critical revision

Pascal Dô : acquisition of data, critical revision

Eric Dansin : acquisition of data, critical revision

Charlotte Leduc : acquisition of data, critical revision
Etienne Giroux Leprieur : acquisition of data, critical revision
Denis Moro-Sibilot : acquisition of data, critical revision
David Tulasne : critical revision
Zoulika Kherrouche : study design and conception, critical revision
Julien Labreuche : study design and conception, interpretation of data
Alexis B. Cortot : study design and conception, analysis and interpretation of data, drafting manuscript, critical revision

ACKNOWLEDGMENTS

This work was supported by grants from the SIRIC ONCOLille, the Fondation du Souffle, and the Fonds de Dotation Recherche en Santé Respiratoire. The authors are grateful to Ms Maeva Kyheng of the biostatistics unit of Lille University Hospital for her assistance regarding the statistical analysis, and Mr Eric Wasielewski for his support regarding the administrative procedure of the protocol approval.

CONFLICTS OF INTEREST

A B. Cortot reported conflicts of interest with : Astra-Zeneca, Boehringer-Ingelheim, Roche, Novartis, BMS, MSD, Pfizer

Simon Baldacci reported conflicts of interest with : Lilly, GSK, Pfizer, Roche

Nicolas Girard reported conflicts of interest with : Astra-Zeneca, Boehringer-Ingelheim, Roche, Novartis, BMS, MSD, Pfizer

Clarisse Audigier-Valette reported conflicts of interest with : Astra-Zeneca, Boehringer-Ingelheim, Roche, Novartis, BMS, MSD, Pfizer

Florian Guisier reported conflicts of interest with : Boehringer-Ingelheim, BMS

FUNDING

This work was supported by grants from the SIRIC ONCOLille, the Fondation du Souffle, and the Fond de Dotation Recherche en Santé Respiratoire.

REFERENCES

1. Barlesi F, Mazieres J, Merlio JP, Debieuvre D, Mosser J, Lena H, Ouafik L, Besse B, Rouquette I, Westeel V, Escande F, Monnet I, Lemoine A, et al. Routine molecular profiling of patients with advanced non-small-cell lung cancer: results of a 1-year nationwide programme of the French Cooperative Thoracic Intergroup (IFCT). *Lancet Lond Engl.* 2016; 387: 1415–26. [https://doi.org/10.1016/S0140-6736\(16\)00004-0](https://doi.org/10.1016/S0140-6736(16)00004-0).
2. Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, Sunpaweravong P, Han B, Margono B, Ichinose Y, Nishiwaki Y, Ohe Y, Yang JJ, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med.* 2009; 361: 947–57. <https://doi.org/10.1056/NEJMoa0810699>.
3. Maemondo M, Inoue A, Kobayashi K, Sugawara S, Oizumi S, Isobe H, Gemma A, Harada M, Yoshizawa H, Kinoshita I, Fujita Y, Okinaga S, Hirano H, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med.* 2010; 362: 2380–8. <https://doi.org/10.1056/NEJMoa0909530>.
4. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia Gomez R, Pallares C, Sanchez JM, Porta R, Cobo M, Garrido P, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol.* 2012; 13: 239–46. [https://doi.org/10.1016/S1470-2045\(11\)70393-X](https://doi.org/10.1016/S1470-2045(11)70393-X).
5. Sequist LV, Yang JCH, Yamamoto N, O'Byrne K, Hirsh V, Mok T, Geater SL, Orlov S, Tsai CM, Boyer M, Su WC, Bennouna J, Kato T, et al. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol.* 2013; 31: 3327–34. <https://doi.org/10.1200/JCO.2012.44.2806>.
6. Cortot AB, Jänne PA. Molecular mechanisms of resistance in epidermal growth factor receptor-mutant lung adenocarcinomas. *Eur Respir Rev.* 2014; 23: 356–66. <https://doi.org/10.1183/09059180.00004614>.
7. Zhou W, Ercan D, Chen L, Yun CH, Li D, Capelletti M, Cortot AB, Chiriac L, Iacob RE, Padera R, Engen JR, Wong KK, Eck MJ, et al. Novel mutant-selective EGFR kinase inhibitors against EGFR T790M. *Nature.* 2009; 462: 1070–4. <https://doi.org/10.1038/nature08622>.
8. Jänne PA, Yang JC, Kim DW, Planchard D, Ohe Y, Ramalingam SS, Ahn MJ, Kim SW, Su WC, Horn L, Haggstrom D, Felip E, Kim JH, et al. AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer. *N Engl J Med.* 2015; 372: 1689–99. <https://doi.org/10.1056/NEJMoa1411817>.
9. Engelman JA, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park JO, Lindeman N, Gale CM, Zhao X, Christensen J, Kosaka T, Holmes AJ, Rogers AM, et al. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science.* 2007; 316: 1039–43. <https://doi.org/10.1126/science.1141478>.
10. Bean J, Brennan C, Shih JY, Riely G, Viale A, Wang L, Chitale D, Motoi N, Szoke J, Broderick S, Balak M, Chang WC, Yu CJ, et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc Natl Acad*

- Sci U S A. 2007; 104: 20932–7. <https://doi.org/10.1073/pnas.0710370104>.
11. Arcila ME, Oxnard GR, Nafa K, Riely GJ, Solomon SB, Zakowski MF, Kris MG, Pao W, Miller VA, Ladanyi M. Rebiopsy of lung cancer patients with acquired resistance to EGFR inhibitors and enhanced detection of the T790M mutation using a locked nucleic acid-based assay. *Clin Cancer Res*. 2011; 17: 1169–80. <https://doi.org/10.1158/1078-0432.CCR-10-2277>.
 12. Sequist LV, Waltman BA, Dias Santagata D, Digumarthy S, Turke AB, Fidias P, Bergethon K, Shaw AT, Gettinger S, Cospers AK, Akhavanfard S, Heist RS, Temel J, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med*. 2011; 3: 75ra26. <https://doi.org/10.1126/scitranslmed.3002003>.
 13. Yu HA, Arcila ME, Rekhman N, Sima CS, Zakowski MF, Pao W, Kris MG, Miller VA, Ladanyi M, Riely GJ. Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin Cancer Res*. 2013; 19: 2240–7. <https://doi.org/10.1158/1078-0432.CCR-12-2246>.
 14. Noro R, Seike M, Zou F, Soeno C, Matsuda K, Sugano T, Nishijima N, Matsumoto M, Kitamura K, Kosaihiira S, Minegishi Y, Yoshimura A, Kubota K, et al. MET FISH-positive status predicts short progression-free survival and overall survival after gefitinib treatment in lung adenocarcinoma with EGFR mutation. *BMC Cancer*. 2015; 15: 31. <https://doi.org/10.1186/s12885-015-1019-1>.
 15. Schildhaus HU, Schultheis AM, Rüschoff J, Binot E, Merkelbach Bruse S, Fassunke J, Schulte W, Ko YD, Schlesinger A, Bos M, Gardizi M, Engel Riedel W, Brockmann M, et al. MET amplification status in therapy-naïve adeno- and squamous cell carcinomas of the lung. *Clin Cancer Res*. 2015; 21: 907–15. <https://doi.org/10.1158/1078-0432.CCR-14-0450>.
 16. Park S, Koh J, Kim DW, Kim M, Keam B, Kim TM, Jeon YK, Chung DH, Heo DS. MET amplification, protein expression, and mutations in pulmonary adenocarcinoma. *Lung Cancer Amst Neth*. 2015; 90: 381–7. <https://doi.org/10.1016/j.lungcan.2015.10.022>.
 17. Gou LY, Li AN, Yang JJ, Zhang XC, Su J, Yan HH, Xie Z, Lou NN, Liu SY, Dong ZY, Gao HF, Zhou Q, Zhong WZ, et al. The coexistence of MET over-expression and an EGFR T790M mutation is related to acquired resistance to EGFR tyrosine kinase inhibitors in advanced non-small cell lung cancer. *Oncotarget*. 2016; 7: 51311–51319. <https://doi.org/10.18632/oncotarget.9697>.
 18. Gainor JF, Niederst MJ, Lennerz JK, Dagogo-Jack I, Stevens S, Shaw AT, Sequist LV, Engelman JA. Dramatic Response to Combination Erlotinib and Crizotinib in a Patient with Advanced, EGFR-Mutant Lung Cancer Harboring De Novo MET Amplification. *J Thorac Oncol*. 2016; 11: e83–85. <https://doi.org/10.1016/j.jtho.2016.02.021>.
 19. Bahcall M, Sim T, Paweletz CP, Patel JD, Alden RS, Kuang Y, Sacher AG, Kim ND, Lydon C, Awad MM, Jaklitsch MT, Sholl LM, Jänne PA, et al. Acquired MET D1228V Mutation and Resistance to MET Inhibition in Lung Cancer. *Cancer Discov*. 2016; 6: 1335–1341. <https://doi.org/10.1158/2159-8290.CD-16-0686>.
 20. Tanaka A, Sueoka-Aragane N, Nakamura T, Takeda Y, Mitsuoka M, Yamasaki F, Hayashi S, Sueoka E, Kimura S. Co-existence of positive MET FISH status with EGFR mutations signifies poor prognosis in lung adenocarcinoma patients. *Lung Cancer Amst Neth*. 2012; 75: 89–94. <https://doi.org/10.1016/j.lungcan.2011.06.004>.
 21. Cappuzzo F, Marchetti A, Skokan M, Rossi E, Gajapathy S, Felicioni L, Del Gramastro M, Sciarrotta MG, Buttitta F, Incarbone M, Toschi L, Finocchiaro G, Destro A, et al. Increased MET gene copy number negatively affects survival of surgically resected non-small-cell lung cancer patients. *J Clin Oncol*. 2009; 27: 1667–74. <https://doi.org/10.1200/JCO.2008.19.1635>.
 22. Cappuzzo F, Jänne PA, Skokan M, Finocchiaro G, Rossi E, Ligorio C, Zucali PA, Terracciano L, Toschi L, Roncalli M, Destro A, Incarbone M, Alloisio M, et al. MET increased gene copy number and primary resistance to gefitinib therapy in non-small-cell lung cancer patients. *Ann Oncol*. 2009; 20: 298–304. <https://doi.org/10.1093/annonc/mdn635>.
 23. Kris MG, Johnson BE, Berry LD, Kwiatkowski DJ, Iafrate AJ, Wistuba II, Varella-Garcia M, Franklin WA, Aronson SL, Su PF, Shyr Y, Camidge DR, Sequist LV, et al. Using multiplexed assays of oncogenic drivers in lung cancers to select targeted drugs. *JAMA*. 2014; 311: 1998–2006. <https://doi.org/10.1001/jama.2014.3741>.
 24. Casadevall D, Gimeno J, Clavé S, Taus Á, Pijuan L, Arumí M, Lorenzo M, Menéndez S, Cañadas I, Albanell J, Serrano S, Espinet B, Salido M, et al. MET expression and copy number heterogeneity in nonsquamous non-small cell lung cancer (nsNSCLC). *Oncotarget*. 2015; 6: 16215–26. <https://doi.org/10.18632/oncotarget.3976>.
 25. Watermann I, Schmitt B, Stellmacher F, Müller J, Gaber R, Kugler C, Reinmuth N, Huber RM, Thomas M, Zabel P, Rabe KF, Jonigk D, Warth A, et al. Improved diagnostics targeting c-MET in non-small cell lung cancer: expression, amplification and activation? *Diagn Pathol*. 2015; 10: 130. <https://doi.org/10.1186/s13000-015-0362-5>.
 26. Beau-Faller M, Ruppert AM, Voegeli AC, Neuville A, Meyer N, Guerin E, Legrain M, Mennecier B, Wihlm JM, Massard G, Quoix E, Oudet P, Gaub MP. MET gene copy number in non-small cell lung cancer: molecular analysis in a targeted tyrosine kinase inhibitor naïve cohort. *J Thorac Oncol*. 2008; 3: 331–9. <https://doi.org/10.1097/JTO.0b013e318168d9d4>.
 27. Camidge DR, Ou SHI, Shapiro G, Otterson GA, Villaruz LC, Villalona-Calero MA, Iafrate AJ, Varella Garcia M, Dacic S, Cardarella S, Zhao W, Tye L, Stephenson P, et al. Efficacy and safety of crizotinib in patients with advanced

- c-MET-amplified non-small cell lung cancer (NSCLC). *J Clin Oncol*. 2014 [cited 2016 Nov 13]; 32:5s. Available from <http://meetinglibrary.asco.org/content/132030-144>.
28. Noonan SA, Berry L, Lu X, Gao D, Barón AE, Chesnut P, Sheren J, Aisner DL, Merrick D, Doebele RC, Varella-Garcia M, Camidge DR. Identifying the Appropriate FISH Criteria for Defining MET Copy Number-Driven Lung Adenocarcinoma through Oncogene Overlap Analysis. *J Thorac Oncol*. 2016; 11: 1293–304. <https://doi.org/10.1016/j.jtho.2016.04.033>.
 29. Kryeziu K, Pirker C, Engliger B, van Schoonhoven S, Spitzwieser M, Mohr T, Körner W, Weinmüller R, Tav K, Grillari J, Cichna-Marx M, Berger W, Heffeter P. Chronic arsenic trioxide exposure leads to enhanced aggressiveness via Met oncogene addiction in cancer cells. *Oncotarget*. 2016; 7: 27379. <https://doi.org/10.18632/oncotarget.8415>.
 30. Tong JH, Yeung SF, Chan AW, Chung LY, Chau SL, Lung RW, Tong CY, Chow C, Tin EK, Yu YH, Li H, Pan Y, Chak WP, et al. MET Amplification and Exon 14 Splice Site Mutation Define Unique Molecular Subgroups of Non-Small Cell Lung Carcinoma with Poor Prognosis. *Clin Cancer Res*. 2016; 22: 3048–56. <https://doi.org/10.1158/1078-0432.CCR-15-2061>.
 31. Lee J, Ou SH, Lee JM, Kim HC, Hong M, Kim SY, Jang J, Ahn S, Kang SY, Lee S, Kim ST, Kim B, Choi J, et al. Gastrointestinal malignancies harbor actionable MET exon 14 deletions. *Oncotarget*. 2015; 6: 28211–22. <https://doi.org/10.18632/oncotarget.4721>.
 32. Oxnard GR, Arcila ME, Sima CS, Riely GJ, Chmielecki J, Kris MG, Pao W, Ladanyi M, Miller VA. Acquired resistance to EGFR tyrosine kinase inhibitors in EGFR-mutant lung cancer: distinct natural history of patients with tumors harboring the T790M mutation. *Clin Cancer Res*. 2011; 17: 1616–22. <https://doi.org/10.1158/1078-0432.CCR-10-2692>.
 33. Sun JM, Ahn MJ, Choi YL, Ahn JS, Park K. Clinical implications of T790M mutation in patients with acquired resistance to EGFR tyrosine kinase inhibitors. *Lung Cancer Amst Neth*. 2013; 82: 294–8. <https://doi.org/10.1016/j.lungcan.2013.08.023>.
 34. Dimou A, Non L, Chae YK, Tester WJ, Syrigos KN. MET gene copy number predicts worse overall survival in patients with non-small cell lung cancer (NSCLC); a systematic review and meta-analysis. *PloS One*. 2014; 9: e107677. <https://doi.org/10.1371/journal.pone.0107677>.
 35. Guo B, Cen H, Tan X, Liu W, Ke Q. Prognostic value of MET gene copy number and protein expression in patients with surgically resected non-small cell lung cancer: a meta-analysis of published literatures. *PloS One*. 2014; 9: e99399. <https://doi.org/10.1371/journal.pone.0099399>.
 36. Turke AB, Zejnullahu K, Wu YL, Song Y, Dias-Santagata D, Lifshits E, Toschi L, Rogers A, Mok T, Sequist L, Lindeman NI, Murphy C, Akhavanfar S, et al. Preexistence and clonal selection of MET amplification in EGFR mutant NSCLC. *Cancer Cell*. 2010; 17: 77–88. <https://doi.org/10.1016/j.ccr.2009.11.022>.
 37. Wu YL, Kim DW, Felip E, Zhang L, Liu X, Zhou CC, Lee DH, Han JY, Krohn A, Lebouteiller R, Xu C, Squires M, Akimov M, et al. Phase (Ph) II safety and efficacy results of a single-arm phase II study of capmatinib (INC280) + gefitinib in patients (pts) with EGFR-mutated (mut), cMET-positive (cMET+) non-small cell lung cancer (NSCLC). *J Clin Oncol*. 2016 [cited 2016 Nov 16]; 34. Available from <http://meetinglibrary.asco.org/content/165978-176>.
 38. Hata A, Katakami N, Yoshioka H, Takeshita J, Tanaka K, Nanjo S, Fujita S, Kaji R, Imai Y, Monden K, Matsumoto T, Nagata K, Otsuka K, et al. Rebiopsy of non-small cell lung cancer patients with acquired resistance to epidermal growth factor receptor-tyrosine kinase inhibitor: Comparison between T790M mutation-positive and mutation-negative populations. *Cancer*. 2013; 119: 4325–32. <https://doi.org/10.1002/cncr.28364>.