



**HAL**  
open science

## A rapid and quantitative safranin-based fluorescent microscopy method to evaluate cell wall lignification

Fabien Baldacci-crep, Corentin Spriet, Laure Twyffels, Anne-Sophie Blervacq, Godfrey G. Neutelings, Marie Baucher, Simon Hawkins

### ► To cite this version:

Fabien Baldacci-crep, Corentin Spriet, Laure Twyffels, Anne-Sophie Blervacq, Godfrey G. Neutelings, et al.. A rapid and quantitative safranin-based fluorescent microscopy method to evaluate cell wall lignification. *Plant Journal*, 2020, *The Plant Journal*, 102 (5), pp.1074-1089. 10.1111/tpj.14675 . hal-03111641

**HAL Id: hal-03111641**

**<https://hal.univ-lille.fr/hal-03111641>**

Submitted on 15 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*the plant journal***A rapid and quantitative safranin-based fluorescent microscopy method to evaluate cell wall lignification**

Journal:	<i>The Plant Journal</i>
Manuscript ID	Draft
Manuscript Type:	Technical Advance
Biochemistry and Physiology:	
Cell Biology:	Cell wall
Genomics & Genetics:	
Plant Growth & Development:	
Plant interactions with other organisms:	
Plant Responses to Environment:	
Other (please specify):	

SCHOLARONE™  
Manuscripts

1  
2  
3 **A rapid and quantitative safranin-based fluorescent microscopy method to evaluate cell**  
4 **wall lignification**  
5  
6  
7  
8

9 Fabien Baldacci-Cresp<sup>1,2</sup>, Corentin Spriet<sup>1,3</sup>, Laure Twyffels<sup>4</sup>, Anne-Sophie Blervacq<sup>1</sup>, Godfrey  
10 Neutelings<sup>1</sup>, Marie Baucher<sup>2</sup> and Simon Hawkins<sup>1</sup>.  
11  
12  
13

14  
15 <sup>1</sup> Université de Lille, CNRS, UMR 8576 – UGSF - Unité de Glycobiologie Structurale et Fonctionnelle,  
16 F-59000 Lille, France.  
17  
18

19 <sup>2</sup> Université libre de Bruxelles, Laboratoire de Biotechnologie Végétale (LBV), B-6041 Gosselies,  
20 Belgium.  
21  
22

23 <sup>3</sup> TISBio, Université de Lille, CNRS, UMR 8576 – UGSF - Unité de Glycobiologie Structurale et  
24 Fonctionnelle, F-59000 Lille, France.  
25  
26

27 <sup>4</sup> Université libre de Bruxelles, Center for Microscopy and Molecular Imaging (CMMI), B-6041  
28 Gosselies, Belgium.  
29  
30  
31

32  
33  
34 Corresponding Author: Fabien Baldacci-Cresp & Simon Hawkins  
35

36 [fabien.baldacci-cresp@univ-lille.fr](mailto:fabien.baldacci-cresp@univ-lille.fr) & [simon.hawkins@univ-lille.fr](mailto:simon.hawkins@univ-lille.fr)  
37  
38  
39  
40

41 E-mail addresses: Fabien Baldacci-Cresp: [fabien.baldacci-cresp@univ-lille.fr](mailto:fabien.baldacci-cresp@univ-lille.fr); Corentin Spriet :  
42 [cspriet@univ-lille.fr](mailto:cspriet@univ-lille.fr); Laure Twyffels : [Laure.Twyffels@ulb.ac.be](mailto:Laure.Twyffels@ulb.ac.be); Anne-Sophie Blervacq : [anne-  
44 sophie.blervacq@univ-lille.fr](mailto:anne-<br/>43 sophie.blervacq@univ-lille.fr); Godfrey Neutelings : [godfrey.neutelings@univ-lille.fr](mailto:godfrey.neutelings@univ-lille.fr); Marie Baucher:  
45 [mbaucher@ulb.ac.be](mailto:mbaucher@ulb.ac.be); Simon Hawkins : [simon.hawkins@univ-lille.fr](mailto:simon.hawkins@univ-lille.fr)  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## SUMMARY

One of the main characteristics of plant cells is the presence of the cell wall located outside the plasma membrane. In particular cells, this wall can be reinforced by lignin, a polyphenolic polymer that plays a central role for vascular plants, conferring hydrophobicity to conducting tissues and mechanical support for upright growth. Lignin has been extensively studied by a range of different techniques including anatomical and morphological analyses using dyes to characterize the polymer localization *in situ*. With the constant improvement of imaging techniques, it is now possible to revisit old qualitative techniques and adapt them to obtain efficient, highly resolutive, quantitative, fast and safe methodologies. In this study, we revisit and exploit the potential of fluorescent microscopy coupled to safranin-O staining to develop a quantitative approach for lignin content determination. The developed approach is based on ratiometric emission measurements and the development of an ImageJ macro. To demonstrate the interest of our methodology compared to other commonly-used lignin reagents, we showed that safranin-O staining was used to evaluate and compare lignin contents in previously characterized *A. thaliana* lignin biosynthesis mutants. In addition, the analysis of lignin content and spatial distribution in the *Arabidopsis* laccase mutant also provided new biological insights into the effects of *laccase* gene down-regulation in different cell types. Our safranin-O based methodology, also validated for flax, maize and poplar, significantly improves and speed-up anatomical and developmental investigations of lignin, which we hope will contribute to new discoveries in many areas of cell wall plant research.

## SIGNIFICANCE STATEMENT (2 sentences 75 words max)

Lignin has long been the subject of extensive study as it plays a major role in plant growth and defense and is an important factor affecting the quality of plant biomass and plant-derived products. The development of an easy and quantitative confocal microscopy method to provide detailed spatial information about lignin content at the cell wall level will prove a valuable addition to the toolbox of lignin analytical methods for both fundamental and applied research.

## INTRODUCTION

A main characteristic of the plant cell is the presence of a cell wall. This wall allows plant cells to acquire a certain rigidity essential for the maintenance of a strong intracellular osmotic pressure that contributes to maintaining the plant stem upright. Depending upon cell types, this wall can be mainly composed of up to three layers. The first one is the middle lamella, a thin structure mainly composed of pectins and shared by two adjacent cells. This layer is the pedestal on which the two other structures – the primary and the secondary cell walls - can be developed. The primary wall is the first to be deposited and is mainly composed of cellulose, hemicelluloses, pectins and proteins (Carpita and Gibeaut, 1993). In certain tissues, plant cells acquire a secondary wall that is deposited between the primary cell wall and the plasma membrane. Like primary cell walls, secondary cell walls are rich in cellulose, but generally show higher hemicellulose and lower pectin content providing greater mechanical strength and reduced extensibility. This secondary cell wall, generally much thicker than the primary wall and made of several layers (S1, S2 and S3), is mainly observed in cells from vascular and supporting tissues. The formation of the secondary cell wall is, in most cases, associated with the deposition of lignin that confers hydrophobicity and rigidity to the cell wall (Boerjan et al., 2003). Lignin is a phenolic polymer composed of 3 main units, *p*-hydroxyphenyl (H), guaiacyl (G) and syringyl (S) units derived from *p*-coumaryl alcohol, coniferyl alcohol and sinapyl alcohol, respectively. The biosynthesis of lignins is initiated during the formation of the S1 secondary cell wall layer and newly formed lignin is first observed in the cell corners and middle lamella before spreading progressively, to the primary cell wall and different layers (S1, S2 and S3) of the secondary cell wall (Terashima and Fukushima, 1988). Given the major importance of lignin – both for the plant and economically - it is not surprising that the biosynthesis/polymerization and degradation of this polymer have been extensively studied for a very long time (Boerjan et al., 2003).

One of the major challenges of recent and future years is to improve our ability to characterize, both quantitatively and qualitatively, lignin in plant cell walls. In terms of quantitative analysis, three main types of wet chemistry methods are traditionally used: acetylbromide assays (Morrison, 1972b, Morrison, 1972a), the acidic method (Van Soest, 1963) or the Klason method (Johnson et al., 1961). If

1  
2  
3 the Klason method seems more suitable and efficient on woody species, the acetylbromide and the acidic  
4 methods are more suitable for herbaceous species. In these species, the acetyl bromide method is faster  
5 and presents a better recovery of lignin (Fukushima et al., 2015).  
6  
7  
8

9  
10 Various types of methods are also available for determining lignin chemical composition (i.e.  
11 determination of S, G and H contents and calculation of S/G ratios). There include chemical degradation  
12 techniques, such as thioacidolysis (Lapierre et al., 1986) and nitrobenzene oxidation (Monties, 1989),  
13 as well as derivatization methods and spectroscopic methods based on NMR (Mansfield et al., 2012) or  
14 using pyrolysis-GC/MS coupling (Wagner et al., 2007).  
15  
16  
17  
18  
19

20  
21 If these qualitative and quantitative approaches are powerful, they all present the same drawback: they  
22 are destructive methods that require grinding and/or extraction of cell wall polymers/lignins from the  
23 plant sample being analyzed. As a result, the data obtained represent an overall mean value for a whole  
24 organ/tissue and thus do not take into account the fact that lignin content and composition vary between  
25 different tissues, cells and cell wall layers. To get around this problem and to obtain information on  
26 lignin localization more rapidly, researchers have exploited different histochemical methods including  
27 basic fushin and auramine staining (epifluorescence observations) (Dharmawardhana et al., 1992,  
28 Pesquet et al., 2005) revealing global lignin deposition, as well as more 'specific' tests such as  
29 phloroglucinol staining (Weisner reagent) revealing lignin hydroxycinnamaldehyde functions (Vallet et  
30 al., 1996, Liljegren, 2010), and Maüle staining providing indications on the S/G ratio (Meshitsuka and  
31 Nakano, 1979). Another approach based on safranin/alcian blue staining, the FASGA method (Tolivia  
32 and Tolivia, 1987), coupled to an adapted image workflow (Legland et al., 2017) allows a quantification  
33 of the histology of maize based on lignin-containing tissues detection but does not permit a strict  
34 quantification of cell wall lignin content.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50  
51 Other approaches based on micro-vibrational spectroscopy (IR and Raman) were successfully adapted  
52 to plant biology and more particularly to *in situ* cell wall characterization (Gierlinger, 2014, Agarwal,  
53 2006, Agarwal and Ralph, 1997, Gierlinger, 2018). They allow scientists to obtain qualitative,  
54 quantitative and spatial data on lignin. However, microspectroscopy approaches are generally very  
55 difficult to calibrate in terms of acquisition analysis and are also time consuming for both machine use  
56  
57  
58  
59  
60

1  
2  
3 and data analyses. Overall, such methodology is not adapted for the rapid measurement of lignin content  
4  
5 in a tissue/sample compared to a reference and can therefore represent an inconvenience for functional  
6  
7 genomics approaches that can involve the analyses of large numbers of samples (e.g. mutant  
8  
9 populations, biological replicates).  
10

11  
12 In 2008, Bond and co-authors (Bond et al., 2008) demonstrated that the azo dye safranin-O, commonly  
13  
14 used with a counterstain to color lignified cell walls red in botanical samples examined by transmitted  
15  
16 light in the visible spectrum, also presents an interesting fluorescence characteristic. Using a range of  
17  
18 different plant samples made up of cells with high or low lignin levels these authors showed that the  
19  
20 safranin-O emission spectrum depends on lignin quantity – the higher the lignin content of the stained  
21  
22 tissue, the more intense the red part of the emission spectrum. While providing a novel way to distinguish  
23  
24 lignin-rich from cellulose-rich cell walls without the need for counterstaining, their approach remained  
25  
26 qualitative and did not allow for statistical comparison of lignin levels in different samples, We therefore  
27  
28 decided to develop a technique that would i) generate high definition spatial information about lignin  
29  
30 localization, ii) provide quantitative information that can be analyzed to support statistically significant  
31  
32 differences in lignin content between samples, and iii) be fast, easy and low-risk for the experimenter.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## RESULTS

### Safranin emission spectra depends on lignin content

Bond and co-authors (Bond et al., 2008) demonstrated that the safranin emission spectrum changes according to cell wall lignin content. Higher lignin amount induces a relative increase in the red part of the emission spectrum. This effect can be clearly seen in the safranin-O spectra (under 488 nm excitation) obtained from different cell walls (zones 1-4) in a stem cross-section from a 2-month-old flax plant (Fig. 1). We used flax as an appropriate model plant to develop the safranin-based methodology since the stem of this species contains two populations of cells with highly contrasted secondary cell wall lignification profiles: xylem cells with a highly lignified secondary walls and cortical bast fibers with poorly lignified secondary cell walls (Day et al., 2005). This allowed us to observe different levels of cell wall lignification within the same histological section as shown with the toluidine blue staining in Fig. 1a.

Analysis of the Safranin fluorescence emission spectra under 488 nm excitation of different flax stem cell types (Fig. 1b) indicated that the maximum peak intensity is 570 nm regardless of cell type analyzed. Spectral analysis shows that the more lignified the cell wall is, the higher the emission spectrum in the orange and red region (570 - 600 nm) (Fig. 1b). In contrast, the "green" part of the spectrum (520 - 560 nm) remains stable indicating that it is the orange/red part of the spectrum that discriminates between the different cell walls analyzed. Beyond 600 nm, the variability is lost.

Based on these observations, we decided to develop a ratiometric emission spectrum approach for evaluating cell wall lignification levels that would be robust, both within a histological section and between histological sections from different provenances (e.g. species, genotype, age).

### Method development: from sample preparation to image treatment.

We firstly developed a standardized methodology designed to limit time consumption and risk to the manipulator without altering result quality (Method S1 and Fig. S1). The aim of the first (facultative) step with EtOH alcohol is to fix and conserve samples (up to several weeks in 70 % EtOH) for later analysis as well as to limit the autofluorescence of pigments (e.g. chlorophyll) that can interfere with



1  
2  
3 safranin fluorescent emission. The time necessary for this step must be defined depending upon the  
4 considered sample but an overnight incubation is usually suitable for most samples. As described later,  
5 it is important to have flat cross-sections and, in this case, samples are sectioned with a vibroslicer as  
6 thinly as possible (30-100 $\mu$ m) to obtain the best morphology. However, since the methodology is based  
7 on epifluorescence the thickness of the cut will not unduly influence image acquisition. Next, a 10 min  
8 staining step in safranin-O alcoholic solution (0.2 % in 50 % EtOH) is performed at room temperature  
9 in the dark with shaking. This step is followed by a short rinse with 50% EtOH to remove safranin excess  
10 on the tube/plate, a 15 min rinse with 50% EtOH and 2 x 10 min H<sub>2</sub>O rinses under shaking in the dark  
11 to remove excess safranin. Samples are subsequently mounted in distilled water and analyzed with a  
12 confocal microscope.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 Before starting any comparative confocal analyses, it is important to obtain safranin spectra of lignified  
25 and non-lignified cell walls from the plant sample so as to determine invariable “green” channel and  
26 variable “red” channel values. Then, for each series of observations on a given sample, it is necessary  
27 to calibrate laser intensity and time acquisition. The more these two values are low, the less there will  
28 be a risk of safranin spectra shift as discussed later. Since our methodology is based on fluorescence  
29 quantification, the chosen laser intensity and time acquisition must provide enough fluorescence  
30 intensity per pixel whilst avoiding saturation. Finally, it is mandatory to select a flat area of the sample  
31 for observation.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

42 Spectrum acquisition provides images of the type presented in Fig. 2a, b and c. The corresponding image  
43 file is then treated with the developed ImageJ macro. This macro applies a region of interest (ROI)  
44 selection, a noise threshold subtraction, a 32-bit conversion and a median radius filter. Next, it defines  
45 the highest intensity value, applies a threshold and converts the background to NaN. At this moment  
46 there are two possibilities for the user: leave the automatic threshold selection, or manually correct this  
47 threshold so as to make it more coherent with your knowledge of your sample morphology. Finally, the  
48 ratio calculation is made and a ratiometric image is generated as presented in Fig 2c.  
49  
50  
51  
52  
53  
54  
55  
56

57 Default settings allow the user to obtain an image representation on a color scale from purple to red in  
58 a ratio range from 0 (purple) to 2 (red). In our context, black pixels correspond to NaN pixels without  
59  
60

1  
2  
3 ratiometric value (under the threshold, not considered in measurement even if selected). Purple pixels  
4  
5 correspond to pixels with ratiometric value close to 0. In the case where the default range does not lead  
6  
7 to useful image representation, it can be restricted or enlarged (Fig. S2). It is important to be careful at  
8  
9 this stage since a valid comparison of different images requires that they are all built using the same  
10  
11 range.  
12

13  
14 In addition to the ratiometric image, the macro also allows a quantitative representation. For this, the  
15  
16 user selects the ROI on the ratiometric image and makes a measurement with image J (Fig. 2d). The  
17  
18 same method is then applied to further ROIs allowing statistical analysis.  
19

### 20 21 Biological validation of the method in Arabidopsis

22  
23 In order to test the robustness of our technique and its capacity to generate biologically relevant  
24  
25 information on cell wall lignin content we applied it to different model plant species. We first evaluated  
26  
27 the ratiometric technique in different lignin mutants from *Arabidopsis thaliana*. The lignin biosynthetic  
28  
29 pathway has been extensively studied in this species over the last 20 years and modifications in lignin  
30  
31 content/structure have been well characterized in a number of different mutants making them ideal test  
32  
33 objects for our technique. Our results (Fig. 3) show that the ratiometric values of fiber cell walls in  
34  
35 *cinnamyl alcohol dehydrogenase (CAD)* and *caffeic acid O-methyl transferase (COMT)* (Fig. 3b, e)  
36  
37 mutants are not significantly different from those of WT plants (Fig. 3a, d). While both mutants contain  
38  
39 similar amounts of lignin to WT plants, *cad* mutants possess lignin with a higher aldehyde content and  
40  
41 *comt* mutants show an altered S/G lignin monomer ratio (Van Acker et al., 2013). The fact that the  
42  
43 ratiometric values remain unchanged suggests that our safranin ratiometric technique is insensitive to  
44  
45 differences in lignin structures. In contrast, similar analyses on the *f5h* mutant (Fig. 3c) revealed a  
46  
47 significant increase in the red/green ratiometric value indicating higher lignin content (9 %) in agreement  
48  
49 with previous lignin chemical analyses (Van Acker et al., 2013). Taken together, these results indicate  
50  
51 that the safranin ratiometric technique reflects cell wall lignin content but is not affected by lignin  
52  
53 composition making it a powerful *in situ* analytical method for lignin quantitative analysis.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 As another example we investigated the *lac4-2 lac17* double mutant previously characterized by Berthet  
4 and co-authors (Berthet et al., 2011). Destructive wet chemistry analysis (acetyl bromide) indicated that  
5 this mutant contains 40 % less lignin in floral stems as compared to WT plants and shows weaker  
6 staining of interfascicular fibers with the Weisner reagent. The application of our technique to floral  
7 stem cross-sections from this mutant enabled us to both confirm previous observations and obtain more  
8 detailed information about the effect of the mutation on the cell wall phenotype (Fig. 4). Firstly, mutant  
9 interfascicular fiber cell walls are thickened, but have less well-organized inner secondary cell walls in  
10 comparison with WT sections (Fig. 4a and b). Secondly, lignification appears to be restricted to the  
11 middle lamella and primary cell wall of mutants compared to WT fibers that also possess lignified  
12 secondary cell walls. In addition to these observations that confirm the results obtained by Berthet et al.,  
13 our method also allowed us to obtain quantification on the amount of lignin reduction in fiber cell walls.  
14 Examination of average ratiometric values (Fig. 4c) indicated that mutant fiber lignin content was  
15 reduced by about 50 % compared to WT cell walls. These values are higher than the reduction in Klason  
16 lignin values (30 – 40 %) determined by Berthet and co-workers but were obtained from analysis of the  
17 whole stem that contains both interfascicular fibers and fiber bundles.

18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
The differences between the values obtained by safranin and wet chemistry of whole stems could suggest  
that the reduction in cell wall lignin is higher in fibers compared to vascular bundles and we therefore  
decided to analyze cell walls in the latter region. The analysis of cell walls in vascular bundles (Fig. 4d  
- f) show that there is no significant difference in mean cell wall lignin levels between WT and mutant  
plants when overall mean values for the bundle are taken into account. This result suggests that the 30-  
40 % reduction in Klason lignin values reported by Berthet and coworkers is due to a proportionally  
greater reduction in fibers compared to vascular bundles and indicates that the *lac4-2 lac17* double  
mutation affects the lignification process in different ways depending upon the tissue considered.

The safranin ratiometric approach also allowed us to obtain new information about the effect of the  
mutation at the cell/cell wall layer level. For example, when the maximum (and not the mean) ratio  
values obtained from the lignified zones of WT fibers (middle lamella, primary cell wall, secondary cell  
wall) and mutant fibers (middle lamella, primary cell wall) are considered (Fig. 4c) the values indicate

1  
2  
3 that lignin levels in middle lamella and primary cell walls are comparable. Taken together these results  
4  
5 indicate that the *lac4-2 lac17* mutation primarily affects secondary cell wall lignification (but not middle  
6  
7 lamella/primary cell wall lignification) and is associated with increased secondary cell wall thickness -  
8  
9 potentially in an attempt to compensate the loss of rigidity due to reduced lignification. The ratiometric  
10  
11 approach also generates intriguing information about the effect of the mutation on the lignification  
12  
13 process in vessels. Indeed, our results would suggest that the down-regulation of the *Lac4* and *Lac17*  
14  
15 genes is not associated with a decrease in overall vascular bundle lignin content, but is, in fact, associated  
16  
17 with increased lignification in xylem vessel cell walls and decreased lignification in fiber cell walls  
18  
19 within the vascular bundle. Interestingly, mutant vascular bundles appear to contain a higher proportion  
20  
21 of xylem fibers with poorly-lignified walls thereby underlining the existence of cell specific lignification  
22  
23 programs.  
24  
25

26  
27 In conclusion, the use of our safranin-based imaging approach allowed us to obtain both quantitative  
28  
29 and precise spatial information about cell wall lignin content in a single experiment and on the same  
30  
31 sample thereby providing a considerable advantage over chemical analysis (quantitative data, but loss  
32  
33 of spatial information) or other histochemical reagents such as Weisner (spatial information, but no  
34  
35 quantitative data).  
36  
37

#### 38 Biological validation of the method in other plant species

39

40  
41 The applicability of our method was used to investigate cell wall lignin in the flax *lbfl* mutant that is  
42  
43 characterized by the presence of bast fibers with a secondary cell wall lignin content of around 17 %  
44  
45 compared to 5 % in WT plants (Chantreau et al., 2014). These values were obtained by destructive wet  
46  
47 chemistry (acetyl bromide) and mask a strong spatial heterogeneity. As shown in Figure 5, the safranin  
48  
49 ratio of bast fibers varies from 0 to 7 unit (safranin ratio mean of 1.2 unit corresponding to 2.4 times  
50  
51 more lignin signal than WT) in a first developmental stage of lignification in 4-week-old plants (Fig. 6b  
52  
53 and d) and from 0 to 15 (safranin ratio mean of 2.2; 4.4 times more lignin signal than WT) in a more  
54  
55 advanced developmental stage of lignification (fig. 5c and d).  
56  
57  
58  
59  
60

1  
2  
3 Lignin content was also evaluated in the cell walls of maize, a monocotyledon plant. On stem cross  
4 sections various tissues with different lignin levels were observed with a 4x objective (Fig. 6a). Non-  
5 lignified parenchyma cell walls appear in black i.e. under the detection threshold of our methodology.  
6  
7 Hypodermis cell walls are colored in blue indicating low lignin levels (safranin ratio = 0.53) in  
8 comparison with the more heavily lignified cell walls of the epidermis (safranin ratio = 0.93) and vascular  
9 bundle sheath that appear red (safranin ratio = 0.98) (Fig. 6a). This approach is complementary to an  
10 approach based on the bioinformatic treatment of FASGA stained stem sections that allows genotyping  
11 based on the surface of lignification (Legland et al., 2017). The safranin ratio methodology also allows  
12 a better resolution as it is independent of the section thickness that can become a limiting point when  
13 the cell density and therefore the opacity to the transmitted light are too important, as it occurs in the  
14 vicinity of large metaxylem vessels (Figure 6b). Sclerenchyma cell walls (red) are more lignified than  
15 vessels (blue to yellow) or phloem cell walls (purple, blue) (Figure 6b). Also, the development and the  
16 associated lignification appears polarized with 5 to 6 lignified cell layers on the protoxylem side against  
17 1 to 3 cell layers on the phloem side. Altogether, these results confirm that the safranin approach can be  
18 easily applied to both monocots and dicots and provides highly detailed ratiometric quantitative data on  
19 cell wall lignin levels in comparison with global wet chemistry analyses which are longer and more  
20 laborious to set up.

#### The finesse of analyzes beyond simple quantification.

21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42 The described safranin method is not only easy to use, but also provides a high resolution spatial  
43 information on cell wall lignin that can usually only be obtained by using more time-consuming and  
44 fastidious techniques such as light/SEM immunolocalisation or Raman spectroscopy (Gierlinger, 2018,  
45 Kiyoto et al., 2013, Day et al., 2005, Simon et al., 2018).

46  
47  
48  
49  
50  
51 An example of high-resolution using the safranin technique is illustrated in Figure 7 showing the results  
52 of an analysis of flax stem xylem. In contrast to other classical histological analyzes (eg. phloroglucinol-  
53 HCl) that suggest a uniform lignification of xylem cell walls (Day et al. 2005; Huis et al., 2012), our  
54 methodology indicates that periclinal cell walls often contain higher amounts of lignin than anticlinal  
55 cell walls (Fig. 7a). Similarly, examination of a pit between two fiber tracheids at 60x magnification  
56  
57  
58  
59  
60

1  
2  
3 (Fig. 7b) provides an image quality and detailed information that approaches that obtained by electron  
4 microscopy (Fromm et al., 2003). The obtained image reveals the localized high lignin levels in the  
5 vicinity of the pit as well as in the compound middle lamella between the two cells.  
6  
7  
8  
9

#### 10 Technical controls – limits to be respected.

11  
12 As in most imaging approaches, it is important to control different technical aspects that can affect the  
13 acquired signal and may lead to the generation of artifacts and false interpretations.  
14  
15  
16

17 A first parameter is related to sample thickness and the focal plane at which observations are made. The  
18 lower the focal plane (i.e. the deeper within the section that one goes), the more incident laser energy  
19 and the resulting emission energy will be decreased. The extent to which this occurs is likely to be  
20 different depending upon the wavelength considered. To characterize this phenomenon in our  
21 methodology, we analyzed green and red channel fluorescence intensities of fiber cell walls at different  
22 focal planes in a section of safranin-O stained poplar xylem (Fig. 8). Our results show that the  
23 fluorescence intensity of the two channels is identical for the first 9 optical sections (upper to lower) and  
24 increases from around 70% of maximum intensity in the first optical section to 100% at the 9<sup>th</sup> optical  
25 section. From the tenth optical section onwards, signal intensity decreases for the 2 channels but to a  
26 greater extent for the green channel compared to the red channel. These results indicate that the good  
27 benchmark for undertaking analysis is when the sample appears focused in the green channel.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 An additional requirement resulting from this phenomenon is that the sample prepared is flat. As shown  
42 in the Figure 8, an uneven sample when analyzed at a given z-depth will give a different result depending  
43 upon the tissue thickness crossed. In Figure 8b, at the surface of the sample section, only the xylem  
44 appears in blue (Fig. 8b, i) whereas deeper in the section, other tissues appear first in blue and then in  
45 yellow and the initial xylem which appears in yellow and then redder (Fig 8b, ii to iv).  
46  
47  
48  
49  
50  
51

52 A second parameter concerns the fluorophore's resistance to bleaching. Since the technique depends on  
53 a ratiometric approach, the absolute intensity level does not matter, and it is more important to check  
54 the ratiometric stability. To evaluate this parameter, we subjected images of a 3-month-old poplar stem  
55 cortical fiber stained by safranin-O to increasing (intensity and accumulation time per pixel) excitation  
56  
57  
58  
59  
60

1  
2  
3 (Fig. 9). For this, we exposed the sample at a laser intensity of 5% (Fig. 9a-c) and 100% (Figure 9d-f)  
4 on continuous acquisition (20s / acquisition - 512x512 pixels image). At 5% laser intensity (Fig. 9a-c),  
5 differences in the percentage red/green ratio between the first and the fifteenth passage were of the order  
6 of 4%, whereas this value increased to around 35 % when 100% laser intensity was used (Figure 9d-f,  
7 g) resulting in a very different visual image depending upon total (intensity and time) laser energy used  
8 (Fig. 9h). Such a result indicates that excessive excitation of safranin affects its spectral properties and  
9 provokes a photoinduced spectral shift in favor of green wavelengths. Analyses should therefore be  
10 performed at low (power and/or time) excitation levels in order to keep any potential changes in the  
11 red/green ratio to a minimum.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

## 25 DISCUSSION

26  
27  
28 Lignin is the third most abundant biopolymer on Earth and plays a vital role in many aspects of plant  
29 life (1). The presence and structure of this polymer also have an important impact on the quality of a  
30 wide range of different plant-based resources and is therefore of major economic importance. Together  
31 with other cell wall polymers, lignin also represents a major sink for atmospheric carbon and its  
32 biosynthesis therefore contributes to counteracting climate change (Bastin et al., 2019). It is therefore  
33 not surprising that scientists have developed a wide range of different techniques for studying this  
34 important polymer.  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Many different chemical and physical techniques are available to quantify and structurally characterize  
45 lignin in plant cell walls (Simon et al., 2018) and have been successfully used on both highly lignified  
46 tissues from woody plants and on more weakly lignified samples such as those found in herbaceous  
47 species.  
48  
49  
50  
51

52 However, many of these dosage methods are destructive and do not take into account the variability in  
53 lignin quantity and structure that occurs between different cell types and/or cell wall layers. As a result,  
54 the potential role of genes in the regulation of lignin biosynthesis and deposition at the cellular level is  
55 often ignored or simplified. This aspect is particularly important given recent reports demonstrating that  
56  
57  
58  
59  
60



1  
2  
3 the deposition of lignin in the cell wall is either under the spatial control of particular proteins localized  
4 in the cell wall (Yi Chou et al., 2018, Hosmani et al., 2013, Tobimatsu and Schuetz, 2019).

5  
6  
7  
8 In contrast to destructive quantitative techniques, a number of *in situ* non-destructive techniques  
9 including classic light microscopy histochemistry, UV microscopy, TEM immunolocalisation and  
10 Raman spectroscopy can also be used to analyse cell wall lignin (Simon et al., 2018).

11  
12  
13  
14  
15 Of these, it is perhaps the Maüle and Wiesner staining methods that are the most widely used light  
16 microscope approaches for analysing lignin *in situ* (Monties, 1989). The Wiesner reagent  
17 (phloroglucinol-HCl) colours lignified cell walls in red and the Maüle reaction brings information about  
18 the S/G ratio. Both of these methods are relatively rapid, easy to use and accessible to most labs that  
19 possess a standard light microscope. However, they do not provide quantitative data thereby preventing  
20 statistical comparisons of cell wall lignin levels, either within the same sample, or between different  
21 samples. In addition, care should also be taken with the Wiesner reagent as it only highlights the  
22 aldehyde groups present in the polymer. In this paper we developed a simple and rapid confocal  
23 fluorescence imaging approach based on safranin-O histochemistry and that is able to provide both  
24 detailed spatial information and quantitative data on cell wall lignin levels.

25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37 Safranin-O is a basic azo dye that binds to the acidic lignin polymer (it is believed to form pi-pi  
38 interactions with the polymer (Stockert et al., 1984). Using a range of plant samples characterized by  
39 the presence of cells showing different amounts of cell wall lignin, Bond and co-workers demonstrated  
40 that cell walls/cell wall regions containing high lignin levels (e.g. cell corners of pine wood) exhibited  
41 red fluorescence in safranin-O-stained samples observed by confocal microscopy, while non-lignified  
42 cell walls (cotton fibers, poplar wood G-layers) fluoresced green. This same study also demonstrated  
43 that the red fluorescence in safranin-O stained samples was spatially correlated with high lignin  
44 autofluorescence in non-stained samples confirming the correlation between lignin quantity and red  
45 fluorescence.

46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56 Since a number of different factors can affect the intensity of acquired signals, we firstly undertook a  
57 number of experiments to standardize sample preparation and analysis. Our results indicated that sample  
58  
59  
60



1  
2  
3 thickness, observation depth and laser intensity should all be carefully controlled to avoid the generation  
4 of artifacts during image acquisition and facilitate valid comparative studies. We also demonstrated that  
5 a simple fixation in 70 % EtOH either before or after section preparation was compatible with the  
6 technique thereby facilitating sample collection and processing.  
7  
8  
9

10  
11  
12 The development of a novel macro in the widely used ImageJ software enabled the generation of a  
13 ratiometric image and the subsequent quantification of relevant cell wall lignin levels in selected areas.  
14 It should be emphasized that the ratiometric images and derived quantitative data do not represent  
15 absolute values such as those obtained by Klason or acetyl bromide analyses (e.g. lignin level expressed  
16 as a % of dry cell wall residue). Moreover, since safranin fluorescence can be modulated by a number  
17 of different factors (dye concentration, incubation and washing time, sample thickness and observation  
18 depth, photobleaching) it is extremely difficult to establish absolute correlation curves. Nevertheless,  
19 the technique does provide quantitative values that allow statistically valid comparisons to be made –  
20 either between different cell walls/cell wall layers in the same sample, or indeed between cell walls/cell  
21 wall layers from different samples (e.g. mutant vs WT plants). This point was demonstrated by our  
22 analysis of the previously characterized *Arabidopsis f5h* lignin mutant that has increased lignin content.  
23 A comparison of safranin-O ratiometric images and statistical analysis of derived quantitative data  
24 indicated statistically significant differences in agreement with previous chemical analysis thereby  
25 confirming that the safranin-O method can be used to analyze cell wall lignin levels *in situ* (Van Acker  
26 et al., 2013). Similar analyses of *Arabidopsis cad* and *comt* mutants indicated no statistically significant  
27 differences in cell wall lignin levels also in agreement with previous chemical analysis (Van Acker et  
28 al., 2013). Although lignin content is unchanged in these two mutants, they do show important  
29 modifications in lignin structure (*cad*: increased G units, decreased S units, increased S-unit aldehydes;  
30 *comt*: almost total elimination of S units, severely decreased S/G ratio). The fact that these differences  
31 did not affect the ratiometric images strongly suggests that the safranin-O based method is insensitive  
32 to such changes in lignin structure. As such, it represents a powerful complement to Wiesner staining  
33 (sensitive to aldehyde content) and Mañile staining (sensitive to S unit content) for characterizing lignin.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Another important advantage of our method is that it enables quantification for individual cell walls/cell  
4 wall layers thereby enabling a much more precise understanding of the effects of mutations on the  
5 lignification process in plants. This point was demonstrated by our analysis of the *lac4-2 lac17* double  
6 mutant previously characterized by Berthet and co-workers (21). Comparison of ratiometric images  
7 from WT and mutant plants enabled us to observe and demonstrate the existence of statistically-  
8 significant changes in cell wall lignin levels at both the tissue (xylem vs interfascicular fiber) and cell  
9 (vessels vs xylem fiber) levels. Somewhat intriguingly, these analyses also suggested that vessel lignin  
10 levels were higher in mutants compared to WT plants.  
11  
12  
13  
14  
15  
16  
17  
18  
19

## 20 CONCLUSION

21  
22  
23 The rapidly increasing availability of relatively low-cost and powerful confocal microscopes is  
24 providing scientists with the possibility to revisit old histochemical methods and to develop more  
25 efficient, safer, faster and more reproducible approaches, with a better spatial resolution. In this paper  
26 we have presented the combination of safranin-O staining coupled to confocal microscopy and an  
27 automated image treatment with the free software ImageJ that allows researchers to obtain quantitative  
28 data on cell wall lignin content very precisely and to undertake statistically valid comparisons between  
29 different cells, either within the same section or between different samples. The toolbox (method plus  
30 provided Image J macro) we describe in this paper will create a field of opportunities for modern  
31 histology in lignin imaging. It can be used to improve and complete the existing classical staining  
32 procedures and other approaches and will contribute to a better understanding of how lignification is  
33 regulated at the cellular level.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## EXPERIMENTAL PROCEDURES

### Plant material and growth conditions

For this study, 4 model plants were used: *Arabidopsis thaliana*, *Linum usitatissimum* (flax), *Populus tremula* x *P. alba* (poplar) and *Zea mays* (maize). For *Arabidopsis thaliana*, wild-type (Col-0, N60000), and selected 'lignin' mutant lines (*comt1*, *cad6*, *f5h1* (Vanholme et al., 2012) and *lac4-2 lac17* double mutant) were used (Berthet et al., 2011). For *L. usitatissimum*, the Diane fiber variety (WT and *lbf1* mutant) was used (Chantreau et al., 2014). For *P. tremula* x *alba*, the 717-1B4 WT line (INRA clone), was used. For *Z. mays*, an indeterminate forage genotype was used. *A. thaliana*, *L. usitatissimum* and *Z. mays* were grown in soil using a 16h/8h photoperiod, 23°C/20°C temperature, 35-65% hygrometry in a greenhouse with natural light completed with sodium to maintain light intensity at 100  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ . *A. thaliana* plants were harvested at stage 6. *Populus* plants were grown in soil for 2.5 months (60 cm high) in a phytotron with a 16h/8h photoperiod, 24°C/21°C temperature, 35-65% hygrometry with white light (HPI Master Plus Philips, metal halid) to maintain light intensity at 120  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ .

### Instruments

Plant stem cross-sections - 80  $\mu\text{m}$  - were made using Microm HV640 and Leica S100 vibroslicers. All the image acquisitions were performed on a LSM710 Confocal (Zeiss) equipped with Argon laser and 541 nm laser and a Nikon A1R confocal equipped with Argon laser and 561 nm laser.

### Image J - Fiji macro step development

An Image J - Fiji macro was developed by successive automated steps with possible user intervention at a key point. The steps of the process are as follows : // split channels; // selection of region of interest and subtraction of background noise; // 32 bit conversion and apply median filter radius 1; // set the green image that has the highest intensity value; // apply a threshold and convert the background to NaN; // wait for user to define threshold; // calculate the ratio (plugin ratioplus) ; // copy aaa\_lut\_fret.lut in the lut field of the Fiji install and run; // Set ratio range; // Measures.

Macro downloading address: <https://nextcloud.univ-lille.fr/index.php/s/rKDBBEM3fznRa6Z>

## Method resume

The complete method is described in the supplementary file Methods S1. A resume scheme of the method is presented in the supplementary figure S1.

## ACKNOWLEDGEMENTS

We thank the TisBio facilities and the FRABio research federation for providing the technical environment conducive to achieving this work. We thank Fabienne Guillon for her critical reading of the manuscript. This study was funded by grants from the French Region Hauts de France and the EU (CPER/FEDER project ALIBIOTECH) to F.B.-C. and by the Fonds de la Recherche Scientifique (FRS-FNRS) research project T.0004.14 to F.B.-C and M.B. We thank Prof. Wout Boerjan for providing the *A.thaliana* lines used in this study. We sincerely thank all those who believed in this project, trusted us and supported us in our approach. Thanks to Dr. Nathalie Bessodes without whom this project could not have been possible. I dedicate this work to my daughters Laura and Rosalie.

## SUPPORTING INFORMATION LEGENDS

**Figure S1.** Presentation of the detailed safranin-based fluorescence method.

**Figure S2.** Illustration of differential settings for the ratiometric representation depending on the chosen color range without affecting ratiometric data for quantification. a) Color range from 0 to 6. b) Color range from 0 to 3. c) Color range from 0 to 1. Scale bar = 150 $\mu$ m.

## REFERENCES

- 1  
2  
3  
4  
5 AGARWAL, U. P. 2006. Raman imaging to investigate ultrastructure and composition of plant cell  
6 walls: distribution of lignin and cellulose in black spruce wood (*Picea mariana*). *Planta*, 224,  
7 1141-53.
- 8 AGARWAL, U. P. & RALPH, S. A. 1997. FT-Raman Spectroscopy of Wood: Identifying  
9 Contributions of Lignin and Carbohydrate Polymers in the Spectrum of Black Spruce (*Picea*  
10 *mariana*). *Applied Spectroscopy*, 51, 1648-1655.
- 11 BASTIN, J. F., FINEGOLD, Y., GARCIA, C., MOLLICONE, D., REZENDE, M., ROUTH, D.,  
12 ZOHNER, C. M. & CROWTHER, T. W. 2019. The global tree restoration potential. *Science*,  
13 365, 76-79.
- 14 BERTHET, S., DEMONT-CAULET, N., POLLET, B., BIDZINSKI, P., CEZARD, L., LE BRIS, P.,  
15 BORREGA, N., HERVE, J., BLONDET, E., BALZERGUE, S., LAPIERRE, C. &  
16 JOUANIN, L. 2011. Disruption of LACCASE4 and 17 results in tissue-specific alterations to  
17 lignification of *Arabidopsis thaliana* stems. *Plant Cell*, 23, 1124-37.
- 18 BOERJAN, W., RALPH, J. & BAUCHER, M. 2003. Lignin biosynthesis. *Annu Rev Plant Biol*, 54,  
19 519-46.
- 20 BOND, J., DONALDSON, L., HILL, S. & HITCHCOCK, K. 2008. Safranin fluorescent staining of  
21 wood cell walls. *Biotech Histochem*, 83, 161-71.
- 22 CARPITA, N. C. & GIBEAUT, D. M. 1993. Structural models of primary cell walls in flowering  
23 plants: consistency of molecular structure with the physical properties of the walls during  
24 growth. *Plant J*, 3, 1-30.
- 25 CHANTREAU, M., PORTELETTE, A., DAUWE, R., KIYOTO, S., CRONIER, D., MORREEL, K.,  
26 ARRIBAT, S., NEUTELINGS, G., CHABI, M., BOERJAN, W., YOSHINAGA, A.,  
27 MESNARD, F., GREC, S., CHABBERT, B. & HAWKINS, S. 2014. Ectopic lignification in  
28 the flax lignified bast fiber1 mutant stem is associated with tissue-specific modifications in  
29 gene expression and cell wall composition. *Plant Cell*, 26, 4462-82.
- 30 DAY, A., RUEL, K., NEUTELINGS, G., CRONIER, D., DAVID, H., HAWKINS, S. &  
31 CHABBERT, B. 2005. Lignification in the flax stem: evidence for an unusual lignin in bast  
32 fibers. *Planta*, 222, 234-45.
- 33 DHARMAWARDHANA, D. P., ELLIS, B. E. & CARLSON, J. E. 1992. Characterization of vascular  
34 lignification in *Arabidopsis-Thaliana*. *Canadian Journal of Botany*, 70, 2238-2244.
- 35 FROMM, J., ROCKEL, B., LAUTNER, S., WINDEISEN, E. & WANNER, G. 2003. Lignin  
36 distribution in wood cell walls determined by TEM and backscattered SEM techniques. *J*  
37 *Struct Biol*, 143, 77-84.
- 38 FUKUSHIMA, R. S., KERLEY, M. S., RAMOS, M. H., PORTER, J. H. & KALLENBACH, R. L.  
39 2015. Comparison of acetyl bromide lignin with acid detergent lignin and Klason lignin and  
40 correlation with in vitro forage degradability. *Animal Feed Science and Technology*, 201, 25-  
41 37.
- 42 GIERLINGER, N. 2014. Revealing changes in molecular composition of plant cell walls on the  
43 micron-level by Raman mapping and vertex component analysis (VCA). *Front Plant Sci*, 5,  
44 306.
- 45 GIERLINGER, N. 2018. New insights into plant cell walls by vibrational microspectroscopy. *Appl*  
46 *Spectrosc Rev*.
- 47 HOSMANI, P. S., KAMIYA, T., DANKU, J., NASEER, S., GELDNER, N., GUERINOT, M. L. &  
48 SALT, D. E. 2013. Dirigent domain-containing protein is part of the machinery required for  
49 formation of the lignin-based Casparian strip in the root. *Proc Natl Acad Sci U S A*, 110,  
50 14498-503.
- 51 JOHNSON, D. B., MOORE, W. E. & ZANK, L. C. 1961. The spectrophotometric determination of  
52 lignin in small wood samples. *TAPPI journal*, 44, 793-798.
- 53 KIYOTO, S., YOSHINAGA, A., TANAKA, N., WADA, M., KAMITAKAHARA, H. & TAKABE,  
54 K. 2013. Immunolocalization of 8-5' and 8-8' linked structures of lignin in cell walls of  
55 *Chamaecyparis obtusa* using monoclonal antibodies. *Planta*, 237, 705-15.
- 56 LAPIERRE, C., MONTIES, B. & ROLANDO, C. 1986. Thioacidolysis of Poplar Lignins:  
57 Identification of Monomeric Syringyl Products and Characterization of Guaiacyl-Syringyl  
58  
59  
60

- 1  
2  
3 Lignin Fractions. *Holzforchung - International Journal of the Biology, Chemistry, Physics*  
4 *and Technology of Wood*.
- 5 LEGLAND, D., EL-HAGE, F., MECHIN, V. & REYMOND, M. 2017. Histological quantification of  
6 maize stem sections from FASGA-stained images. *Plant Methods*, 13, 84.
- 7 LILJEGREN, S. 2010. Phloroglucinol stain for lignin. *Cold Spring Harb Protoc*, 2010, pdb prot4954.
- 8 MANSFIELD, S. D., KIM, H., LU, F. & RALPH, J. 2012. Whole plant cell wall characterization  
9 using solution-state 2D NMR. *Nat Protoc*, 7, 1579-89.
- 10 MESHITSUKA, G. & NAKANO, J. 1979. Studies on the mechanism of lignin color reaction. XIII.  
11 Maule color reaction. *J. Japanese Wood Res. Soc.*, 25, 588-594.
- 12 MONTIES, B. 1989. Lignins. In: DEY, P. M. & HARBORNE, J. B. (eds.) *Methods in plant*  
13 *biochemistry*. London ; San Diego: Academic Press.
- 14 MORRISON, I. M. 1972a. Improvements in the acetyl bromide technique to determine lignin and  
15 digestibility and its application to legumes. *Journal of the Science of Food and Agriculture*,  
16 23, 1463-1469.
- 17 MORRISON, I. M. 1972b. A semi-micro method for the determination of lignin and its use in  
18 predicting the digestibility of forage crops. *Journal of the Science of Food and Agriculture*, 23,  
19 455-463.
- 20 PESQUET, E., RANOCHA, P., LEGAY, S., DIGONNET, C., BARBIER, O., PICHON, M. &  
21 GOFFNER, D. 2005. Novel markers of xylogenesis in zinnia are differentially regulated by  
22 auxin and cytokinin. *Plant Physiol*, 139, 1821-39.
- 23 SIMON, C., LION, C., BIOT, C., GIERLINGER, N. & HAWKINS, S. 2018. Lignification and  
24 Advances in Lignin Imaging in Plant Cell Walls. *Annual Plant Reviews*, 1, 1-32.
- 25 STOCKERT, J. C., CANETE, M. & COLMAN, O. D. 1984. Histochemical mechanism for the  
26 orthochromatic staining and fluorescence reaction of lignified tissues. *Cell Mol Biol*, 30, 503-  
27 8.
- 28 TERASHIMA, N. & FUKUSHIMA, K. 1988. Heterogeneity in formation of lignin—XI: An  
29 autoradiographic study of the heterogeneous formation and structure of pine lignin. *Wood*  
30 *Science and Technology*, 22, 259-270.
- 31 TOBIMATSU, Y. & SCHUETZ, M. 2019. Lignin polymerization: how do plants manage the  
32 chemistry so well? *Curr Opin Biotechnol*, 56, 75-81.
- 33 TOLIVIA, D. & TOLIVIA, J. 1987. Fasca: A New Polychromatic Method for Simultaneous and  
34 Differential Staining of Plant Tissues. *Journal of Microscopy (Oxford)*, 148, 113-117.
- 35 VALLET, C., CHABBERT, B., CZANINSKI, Y. & MONTIES, B. 1996. Histochemistry of Lignin  
36 Deposition during Sclerenchyma Differentiation in Alfalfa Stems. *Annals of Botany*, 78, 625-  
37 632.
- 38 VAN ACKER, R., VANHOLME, R., STORME, V., MORTIMER, J. C., DUPREE, P. & BOERJAN,  
39 W. 2013. Lignin biosynthesis perturbations affect secondary cell wall composition and  
40 saccharification yield in *Arabidopsis thaliana*. *Biotechnol Biofuels*, 6, 46.
- 41 VAN SOEST, P. J. 1963. Use of detergents in the analysis of fibrous foods. II. A rapid method for the  
42 determination of fibre and lignin. *Journal - Association of Official Analytical Chemists*, 46,  
43 829-835.
- 44 VANHOLME, R., STORME, V., VANHOLME, B., SUNDIN, L., CHRISTENSEN, J. H.,  
45 GOEMINNE, G., HALPIN, C., ROHDE, A., MORREEL, K. & BOERJAN, W. 2012. A  
46 Systems Biology View of Responses to Lignin Biosynthesis Perturbations in *Arabidopsis*[W].  
47 *Plant Cell*, 24, 3506-29.
- 48 WAGNER, A., RALPH, J., AKIYAMA, T., FLINT, H., PHILLIPS, L., TORR, K.,  
49 NANAYAKKARA, B. & TE KIRI, L. 2007. Exploring lignification in conifers by silencing  
50 hydroxycinnamoyl-CoA:shikimate hydroxycinnamoyltransferase in *Pinus radiata*. *Proc Natl*  
51 *Acad Sci U S A*, 104, 11856-61.
- 52 YI CHOU, E., SCHUETZ, M., HOFFMANN, N., WATANABE, Y., SIBOUT, R. & SAMUELS, A.  
53 L. 2018. Distribution, mobility, and anchoring of lignin-related oxidative enzymes in  
54 *Arabidopsis* secondary cell walls. *J Exp Bot*, 69, 1849-1859.
- 55  
56  
57  
58  
59  
60



## FIGURE LEGENDS

**Figure 1.** Cell wall associated safranin spectra. a) Flax stem cross-section stained with toluidine blue showing tissue organization and region of interest (yellow dotted circles) used for safranin spectra acquisition, b) Safranin spectra (excitation 488nm) of 4 regions of interest from cells known to contain different amounts of lignin: “low”, (bast fiber cell wall, spectra 1, orange line; cambial cell wall, spectra 2, blue line), “high” (xylem cell walls, spectra 3 and 4, green and yellow lines). bf = bast fibers, c = cortex, cz = cambial zone, m = medulla, p = phloem, px = primary xylem, x = xylem. Scale bar = 100µm

**Figure 2.** Presentation of safranin-based fluorescence method. a) Acquisition of the green part of the safranin spectra (ex:488nm, em:540-560nm), b) Acquisition of the red part of the safranin spectra (ex:561nm, em:570-600nm), c) ratiometric image resulting from red and green image treatment using the Image J macro developed, d) Ratio quantification in different parts of the section. The Region Of Interest (ROI) 1 corresponds to the safranin ratio from hypolignified flax fibers. The ROI 2 corresponds to the safranin ratio from lignified flax xylem. The range color used for image representation goes from ratio 0 (purple) to ratio 2 (red). bf = bast fibers, c = cortex, cz = cambial zone, x = xylem. Scale bar = 100µm

**Figure 3.** Biological validation of the method in *Arabidopsis* monolignol biosynthetic pathway mutants. From a) to c) are represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a), *comt* mutant (b) and *f5h* mutant (c) from a first experiment. In d) and e) are respectively represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a), *cad* mutant (e) from a second independent experiment. f) Histograms of average safranin ratios for each analyzed line, values show no significant differences except \* = significant difference at  $P < 0.01$ . The color range used for the image representations goes from ratio 0 (purple) to ratio 1.5 (red). Scale bar = 50µm.

**Figure 4.** Biological validation of the method in *Arabidopsis lac4-2 lac17* mutant. From a) to b) are represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a) and *lac4-2 lac17* mutant (b). c) Histograms of average and maximum safranin ratios. From d)

to e) are represented safranin ratiometric images of xylem bundle from floral stem cross sections of Col-0 line (d) and *lac4-2 lac17* mutant (e). f) Histogram of xylem bundle average safranin ratios. f) Histogram of xylem vessels average safranin ratios. \* = significant difference at P=0.01. The color range used for the image representations goes from ratio 0 (purple) to ratio 2 (red). Scale bar = 50µm.

**Figure 5.** Biological validation of the method in other plant species: flax. From a) to c) are represented safranin ratiometric images of flax stem cross sections of WT (a) and LBF1 mutant of 4weeks old (b and c) plants. The color range used for the image representations varies from ratio 0 (purple) to ratio 2 (red). d) histogram of average safranin ratio and e) histogram of maximum safranin ratio for each analyzed line. bf = bast fibers, c = cortex, cz = cambial zone, x = xylem. Scale bar = 50µm

**Figure 6.** Biological validation of the method in other plant species: maize. a) Safranin ratiometric image of a stem cross-section of maize observed with a 4x objective. b) Safranin ratiometric image of a vascular bundle from a stem cross-section of maize observed with a 60x objective. c) Histograms of average intensity safranin ratios for epidermis, hypodermis and external vascular bundles. The range color used for image representation varies from ratio 0 (purple) to ratio 1.5 (red). bs = bundle sclerenchyma, e = epidermis, mx = metaxylem, pa = parenchyma, ph = phloem, px = protoxylem, vb = vascular bundle. Scale bar a) = 1mm and b) = 100µm.

**Figure 7.** Flax xylem cell walls: the finesse of the method. a) Safranin ratiometric images of xylem cross sections from 8-weeks-old flax stems observed with a 60x objective. b) Detail of a pit and anticlinal walls from a). Arrows show periclinal xylem cell walls. Scale bar = 15µm

**Figure 8.** Effect of depth and flatness on the safranin ratio. (a) Analysis of green and red channel fluorescence intensity in several Z-optical sections of poplar xylem. Graphical representation of the fluorescence intensity of both channels for 15 optical sections of 6.15µm. Green line corresponds to green channel. Red line corresponds to red channel. Grey area corresponds to the optimal section for analysis and comparison of several independent plant sections. (b) Analysis of *A. thaliana* floral stem cross section. From i) to iv) images of cell wall lignin in a vascular bundle obtained at 4 different optical



1  
2  
3 focal planes. For each optical section, a synthetic sketch illustrates the corresponding image along the  
4  
5 A-B section in term of color corresponding to safranin ratio depending on the depth. Scale bar = 50 $\mu$ m.  
6  
7

8 **Figure 9.** Effects of laser energy (intensity and time) on cell wall images and green/red ratio. The effects  
9  
10 of different 488nm excitation energies were evaluated on fiber poplar cross sections stained with  
11  
12 safranin O. From a) to c) analysis of safranin ratiometric image with a low excitation power (5%). From  
13  
14 d) to f) analysis of safranin ratiometric image with a high excitation power (100%). a) and d) correspond  
15  
16 to one optimal pass acquisition, b) and e) correspond to the image after the 7<sup>th</sup> pass and c) and f)  
17  
18 correspond to the image at the 15<sup>th</sup> pass. g) Quantitative analysis of red/green safranin ratios for 15 pass  
19  
20 acquisitions. White bars correspond to safranin ratio evolution at low laser intensity and black bars  
21  
22 correspond to safranin ratio evolution at high laser intensity. The 1<sup>st</sup> pass acquisition is normalized to  
23  
24 100% for both laser conditions. h) Safranin ratiometric image of poplar fibers including an insert  
25  
26 delimited by a dotted line representing the safranin ratiometric fibers after 15 pass acquisitions.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1.

Cell wall associated safranin spectra. a) Flax stem cross-section stained with toluidine blue showing tissue organization and region of interest (yellow dotted circles) used for safranin spectra acquisition, b) Safranin spectra (excitation 488nm) of 4 regions of interest from cells known to contain different amounts of lignin: "low", (bast fiber cell wall, spectra 1, orange line; cambial cell wall, spectra 2, blue line), "high" (xylem cell walls, spectra 3 and 4, green and yellow lines). bf = bast fibers, c = cortex, cz = cambial zone, m = medulla, p = phloem, px = primary xylem, x = xylem. Scale bar = 100 $\mu$ m

190x275mm (96 x 96 DPI)





Figure 2.

Presentation of safranin-based fluorescence method. a) Acquisition of the green part of the safranin spectra (ex:488nm, em:540-560nm), b) Acquisition of the red part of the safranin spectra (ex:561nm, em:570-600nm), c) ratiometric image resulting from red and green image treatment using the Image J macro developed, d) Ratio quantification in different parts of the section. The Region Of Interest (ROI) 1 corresponds to the safranin ratio from hypolignified flax fibers. The ROI 2 corresponds to the safranin ratio from lignified flax xylem. The range color used for image representation goes from ratio 0 (purple) to ratio 2 (red). bf = bast fibers, c = cortex, cz = cambial zone, x = xylem. Scale bar = 100 $\mu$ m


190x275mm (96 x 96 DPI)



**Figure 3.**

Biological validation of the method in Arabidopsis monoglignol biosynthetic pathway mutants. From a) to c) are represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a), comt mutant (b) and f5h mutant (c) from a first experiment. In d) and e) are respectively represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a), cad mutant (e) from a second independent experiment. f) Histograms of average safranin ratios for each analyzed line, values show no significant differences except \* = significant difference at  $P < 0.01$ . The color range used for the image representations goes from ratio 0 (purple) to ratio 1.5 (red). Scale bar =  $50\mu\text{m}$ .


190x275mm (96 x 96 DPI)



**Figure 4.**

Biological validation of the method in Arabidopsis lac4-2 lac17 mutant. From a) to b) are represented safranin ratiometric images of interfascicular fibers from floral stem cross sections of Col-0 line (a) and lac4-2 lac17 mutant (b). c) Histograms of average and maximum safranin ratios. From d) to e) are represented safranin ratiometric images of xylem bundle from floral stem cross sections of Col-0 line (d) and lac4-2 lac17 mutant (e). f) Histogram of xylem bundle average safranin ratios. g) Histogram of xylem vessels average safranin ratios. \* = significant difference at  $P=0.01$ . The color range used for the image representations goes from ratio 0 (purple) to ratio 2 (red). Scale bar = 50µm.


190x275mm (96 x 96 DPI)



**Figure 5.**

Biological validation of the method in other plant species: flax. From a) to c) are represented safranin ratiometric images of flax stem cross sections of WT (a) and LBF1 mutant of 4weeks old (b and c) plants. The color range used for the image representations varies from ratio 0 (purple) to ratio 2 (red). d) histogram of average safranin ratio and e) histogram of maximum safranin ratio for each analyzed line. bf = bast fibers, c = cortex, cz = cambial zone, x = xylem. Scale bar = 50 $\mu$ m

190x275mm (96 x 96 DPI)



**Figure 6.**

Biological validation of the method in other plant species: maize. a) Safranin ratiometric image of a stem cross-section of maize observed with a 4x objective. b) Safranin ratiometric image of a vascular bundle from a stem cross-section of maize observed with a 60x objective. c) Histograms of average intensity safranin ratios for epidermis, hypodermis and external vascular bundles. The range color used for image representation varies from ratio 0 (purple) to ratio 1.5 (red). bs = bundle sclerenchyma, e = epidermis, mx = metaxylem, pa = parenchyma, ph = phloem, px = protoxylem, vb = vascular bundle. Scale bar a) = 1mm and b) = 100 $\mu$ m.

190x275mm (96 x 96 DPI)




Figure 7.

Flax xylem cell walls: the finesse of the method. a) Safranin ratiometric images of xylem cross sections from 8-weeks-old flax stems observed with a 60x objective. b) Detail of a pit and anticlinal walls from a). Arrows show periclinal xylem cell walls. Scale bar = 15 $\mu$ m

190x275mm (96 x 96 DPI)






Figure 8.

Effect of depth and flatness on the safranin ratio. (a) Analysis of green and red channel fluorescence intensity in several Z-optical sections of poplar xylem. Graphical representation of the fluorescence intensity of both channels for 15 optical sections of  $6.15\mu\text{m}$ . Green line corresponds to green channel. Red line corresponds to red channel. Red line corresponds to red channel. Grey area corresponds to the optimal section for analysis and comparison of several independent plant sections. (b) Analysis of *A. thaliana* floral stem cross section. From i) to iv) images of cell wall lignin in a vascular bundle obtained at 4 different optical focal planes. For each optical section, a synthetic sketch illustrates the corresponding image along the A-B section in term of color corresponding to safranin ratio depending on the depth. Scale bar =  $50\mu\text{m}$ .

190x275mm (96 x 96 DPI)




Figure 9.

Effects of laser energy (intensity and time) on cell wall images and green/red ratio. The effects of different 488nm excitation energies were evaluated on fiber poplar cross sections stained with safranin O. From a) to c) analysis of safranin ratiometric image with a low excitation power (5%). From d) to f) analysis of safranin ratiometric image with a high excitation power (100%). a) and d) correspond to one optimal pass acquisition, b) and e) correspond to the image after the 7th pass and c) and f) correspond to the image at the 15th pass. g) Quantitative analysis of red/green safranin ratios for 15 pass acquisitions. White bars correspond to safranin ratio evolution at low laser intensity and black bars correspond to safranin ratio evolution at high laser intensity. The 1st pass acquisition is normalized to 100% for both laser conditions. h) Safranin ratiometric image of poplar fibers including an insert delimited by a dotted line representing the safranin ratiometric fibers after 15 pass acquisitions.

190x275mm (96 x 96 DPI)




Figure S1.

Presentation of the detailed safranin-based fluorescence method.

190x275mm (96 x 96 DPI)




Figure S2.

Illustration of differential settings for the ratiometric representation depending on the chosen color range without affecting ratiometric data for quantification. a) Color range from 0 to 6. b) Color range from 0 to 3. c) Color range from 0 to 1. Scale bar = 150 $\mu$ m.

190x275mm (96 x 96 DPI)

## Methods S1

**Baldacci-Cresp F., et al.**

### **A rapid and semi-quantitative safranin-based fluorescent microscopy method to evaluate cell wall lignification**

#### **CHEMICALS:**

The following chemicals were used to develop the protocol: Agarose (Sigma, CAS-No: 30525-89-4), Safranin-O (Sigma, CAS-No: 477-73-6), ethanol (Sigma, CAS-No: 64-17-5) and that's all.

#### **SAFRANIN-O SOLUTION:**

- **Ethanol** [final 50% (w/v)],
- **Safranin-O** [final 0.2% (w/v)]
- **Ultrapure water** to the final volume

*Mix the solution well for at least 15min until you get everything completely dissolved.*

#### **PREPARATION OF THE SOLUTION AND ITS STORAGE**

- Safranin-O is purchased as a powder and dissolved in ethanol 50% solution.
- To prepare the staining solutions, weigh the indicated amount of the corresponding dye powder, dissolve it in ethanol 50% solution by mixing the solution on the magnetic stirrer in a small flask for at least 15min.
- To store the solutions, wrap them in aluminium foil to avoid exposing them to light at 4°C.

**Note 1:** if possible prefer fresh safranin-O solution.

**Note 2:** for long term conservation, a 10x solution can be prepared, split in aliquots and keep several months in the dark at -20°C.

#### **SAMPLE TREATMENT**

- Collect samples, maximum 0.5 cm length.
- Place them in 70% EtOH overnight with shaking at 4°C to fix and remove pigments that can impact the fluorescence acquisition.
- Remove alcohol solution.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14
- Coat samples in moult with agarose solution (3 to 4%).
  - Perform section - ideally from 30 $\mu$ m to 100 $\mu$ m - on a vibroslicer and place them on EtOH 50% solution until staining. For this step and depending on your samples and the vibroslicer you must adapt section parameters (speed, frequency, amplitude section) to obtain the best morphology. Better the samples morphology is kept better will be the results.

15  
16  
17  
18  
19  
20  
21  
22  
23

**Note:** the same procedure can be envisaged with other embedding as paraffin or PEG to use microtome instead of vibroslicer to obtain better morphology. In this case the global procedure will be longer. However, the gain in morphological quality is justified only for fragile tissues.

### 24 25

#### SAFRANIN-O STAINING

- 26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41
- Prepare fresh 0.2% Safranin-O.
  - Place vibrosliced samples in safranin-O solution for 10min under shaking at RT at obscurity.
  - Remove all safranin-O solution.
  - Rinse one time in ethanol 50% solution for 10min under shaking at RT at obscurity.
  - Remove all the solution.
  - Rinse two time in ultrapure water for 15min under shaking at RT at obscurity.
  - Mount samples in water for imaging.

42  
43  
44  
45  
46

**Note 1:** For the staining and for rinsing it's important to have a big volume of solution compared to sample number and to incubate in container allowing good homogenization under shaking.

47  
48  
49

**Note 2:** Adapt the flushing time if necessary. Young lignified tissues or tissues with low lignin content don't support too high decolouring time.

50  
51  
52  
53

**Note 3:** Never mount your samples in glycerol or glycerol-containing mounting medium because safranin-O is highly soluble in glycerol.

### 54 55

#### CONFOCAL ANALYSIS

56  
57  
58  
59  
60

For the first analysis:

- Perform a comparative spectra analysis between lignified and non-lignified tissue. This step is essential to determine the peak of maximum intensity, the invariable area of the spectrum (green part, generally between 520nm and 560 nm) and the area of the spectrum sensitive to lignin quantity (yellow to red part, generally between the maximum intensity peak and 630 nm).
- For the spectral analysis use a 488nm excitation coupled with acquisition from 500nm to 700nm. Warning: limit 488 nm laser power and pixel time exposition to avoid spectra alteration as presented in this study.

#### Current analysis:

- To optimize results and limiting 488nm laser using to avoid spectral shift, the acquisition was developed with 2 independent track with two different laser line excitation.
  - Track 1- green channel** - ex: 488nm, em: 530-560 nm – limit 488 nm laser power (1-5% on recent laser).
  - Track 2- red channel** - ex: 561nm, em: 570-600 nm.
- Considering the semi-quantitative comparison between independent samples, settings of the reference sample must limit light range acquisition to 70% of the maximum tones (about 2900 tones for 12 bit image acquisition).
- Acquire sequentially both tracks.

**Note:** Acquisition parameters depend on the confocal microscope used. The important point is to well define the green and the red channel and adapt the parameters in function of.

#### **IMAGE TREATMENT**

- For the image treatment use the Fiji install completed with the developed macro available to this address: (<https://nextcloud.univ-lille.fr/index.php/s/rKDBBEM3fznRa6Z>).
- Open the Fiji app
- Open the macro file with Fiji.
- Open the image.
- Run the macro and follow the instructions.

The result is a ratiometric image (representing ratio range from 0 to 2).

1  
2  
3 **Note 1:** if the macro is completely automated, the user has the possibility to act on the threshold  
4 applied to image treatment. Let the automatic threshold choice except if you consider that the  
5 morphology doesn't correspond to your scientific knowledge of the model studied.  
6  
7

8 **Note 2:** Two parameters of the macro can be adjusted depending on the acquisition.  
9

- 10 - The "rooling" must be adapted according to image resolution and size of the  
11 analysed biological structure which vary with the objective used and the zoom used  
12 for the acquisition.  
13  
14  
15 - The ratio range representation is set from 0 to 2 (purple to red). If the variation is  
16 more or less important, the ratiometric range can be adjusted to optimize  
17 representation. In this case, the same range must be applied to all image included  
18 in the comparison. Modify the range don't modify raw pixel values.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60