



Copper Complexes as Anticancer Agents Targeting Topoisomerases I and II.

Caroline Molinaro, Alain Martoriati, Lydie Pelinski, Katia Cailliau

► To cite this version:

Caroline Molinaro, Alain Martoriati, Lydie Pelinski, Katia Cailliau. Copper Complexes as Anticancer Agents Targeting Topoisomerases I and II.. Cancers, 2020, Cancers, 12 (10), pp.2863. 10.3390/cancers12102863 . hal-03125689v2

HAL Id: hal-03125689

<https://hal.univ-lille.fr/hal-03125689v2>

Submitted on 16 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Review

Copper Complexes as Anticancer Agents Targeting Topoisomerases I and II

Caroline Molinaro ¹ , **Alain Martoriati** ¹ , **Lydie Pelinski** ² and **Katia Cailliau** ^{1,*} ¹ Univ. Lille, CNRS, UMR 8576-UGSF-Unité de Glycobiologie Structurale et Fonctionnelle, F-59000 Lille, France; caroline.molinaro@univ-lille.fr (C.M.); alain.martoriati@univ-lille.fr (A.M.)² Univ. Lille, CNRS, Centrale Lille, Univ. Artois, UMR 8181-UCCS-Unité de Catalyse et Chimie du Solide, F-59000 Lille, France; lydie.pelinski@univ-lille.fr

* Correspondence: katia.maggio@univ-lille.fr

Received: 3 September 2020; Accepted: 29 September 2020; Published: 5 October 2020



Simple Summary: Organometallics, such as copper compounds, are cancer chemotherapeutics used alone or in combination with other drugs. One small group of copper complexes exerts an effective inhibitory action on topoisomerases, which participate in the regulation of DNA topology. Copper complexes of topoisomerase inhibitors work by different molecular mechanisms that have repercussions on the cell cycle checkpoints and death effectors. The expansion of this family of highly active anticancer drugs and their use in combination with other emerging cancer therapies opens new avenues for the treatment of cancers.

Abstract: Organometallics, such as copper compounds, are cancer chemotherapeutics used alone or in combination with other drugs. One small group of copper complexes exerts an effective inhibitory action on topoisomerases, which participate in the regulation of DNA topology. Copper complexes inhibitors of topoisomerases 1 and 2 work by different molecular mechanisms, analyzed herein. They allow genesis of DNA breaks after the formation of a ternary complex, or act in a catalytic mode, often display DNA intercalative properties and ROS production, and sometimes display dual effects. These amplified actions have repercussions on the cell cycle checkpoints and death effectors. Copper complexes of topoisomerase inhibitors are analyzed in a broader synthetic view and in the context of cancer cell mutations. Finally, new emerging treatment aspects are depicted to encourage the expansion of this family of highly active anticancer drugs and to expand their use in clinical trials and future cancer therapy.

Keywords: copper complexes; topoisomerase inhibitor; DNA damage response; cell cycle; cell death; chemotherapy

1. Introduction

Chemotherapy is a systemic treatment proposed to patients suffering from cancer. It is often a complementary approach to surgery or radiotherapy. The discovery of platinum's inhibitory effect on tumor cell growth in the 1960s [1] was a milestone for anticancer drug application in medicine [2]. Platinum (II) sits at the center of the squared planar structure of cisplatin and is coordinated with two chlorides and two ammonia molecules in a cis configuration. Cisplatin and its derivative drugs (carboplatin of second generation and oxaliplatin of third generation) are used worldwide in clinical applications and several other platinum analogs (lobaplatin, nedaplatin, and heptaplatin) are approved in several countries (Figure 1) [3,4]. However, serious side effects including toxicities on the kidney, heart, ear, and liver, decrease in immunity, hemorrhage, and gastrointestinal disorders limit the use of platinum derivatives [5–7]. The appearance of drug resistances, issuing from acquired or intrinsic

multiple genetic and epigenetic changes, has also limited the clinical use of platinum-derived drugs [8]. Platinum-based treatment efficiency is challenged by cross-resistance and multiple changes including a decreased accumulation of the drug, a reduction in DNA–drug adducts, a modification in cell survival gene expression, an alteration of DNA damage repair mechanisms, modifications of transporters, protein trafficking, and altered cell metabolism [9–14].




Figure 1. Platinum (II) complexes.

To circumvent drug resistance, a possible approach consists of designing and developing new therapeutic metal-based anticancer drugs [15–21]. Several transition metals from the d-block of the periodic table (groups 3 to 12) and particularly essential trace metals [15,22,23], such as copper [24–29], are useful for the implementation of metal-based complexes in anticancer therapies. Copper plays central roles in various cellular processes being an essential micronutrient and an important cofactor for several metalloenzymes involved in mitochondrial metabolism (cytochrome c oxidase), or cellular radical detoxification against reactive oxygen species (ROS) (superoxide dismutase) [30]. Copper is essential for angiogenesis, proliferation, and migration of endothelial cells [31–33]. Elevated copper favors tumor growth and metastasis. It is detected in several brain [34], breast [35], colon, prostate [36], and lung [37] tumors and serves as an indicator of the course of the disease [38]. The differences in tumor cells' responses to copper compared to normal cells laid the foundation of copper complexes' (CuC) evolution as anticancer agents. Numerous developed CuC contain different sets of N, S, or O ligands and demonstrate high cytotoxicity and efficient antitumor activity [25]. Different mechanisms are involved in copper drugs' anticancer effect. They act as chelators, and interact with and sequester endogenous copper, reducing its availability for tumor growth and angiogenesis [39]. On the contrary, ionophores trigger intracellular copper accumulation, cytotoxicity, and activate apoptosis inhibitor factor (XIAP) [24,40–46]. Other CuC are proteasome inhibitors [47,48]. Several CuC are actually on clinical trials: a number of copper/disulfiram-based drug combinations for therapy and as diagnostic tools (metastatic breast cancer and germ cell tumor), several casiopeínas compounds and elesclomol (leukemia), and thiosemicarbazone-based copper complexes labeled with a radioactive isotope for positron emission tomography imaging of hypoxia (in head and neck cancers) [49].

The cisplatin DNA-targeting principle of action also conditioned the development of anticancer copper-based drugs [4,23,50]. Antitumor activities of copper-based drugs are based on the interactive properties of both copper and the ligand. Copper toxicity results from its redox capacities (Cu(I) and Cu(II) redox states' interconversion in oxidation–reduction cycles), the property to displace other ions from the enzyme binding sites, a high DNA binding affinity, and the ability to promote DNA breaks [28,51]. In most cases, copper modifies the backbone of the complexed ligand and grants better DNA affinity, specificity, and stability [52]. Copper derivatives can interact with DNA without the formation of covalent adducts. The noncovalent interactions with DNA include binding along with the major or the minor DNA grooves, intercalation, or electrostatic binding. Some copper-based drugs generate reactive oxygen species (ROS) that overwhelm cellular antioxidant defenses to produce

oxidative damages in the cytoplasm, mitochondria, and DNA [53]. An important class of CuC, actually on focus for chemotherapy, inhibits topoisomerases (Top) 1 and 2, resulting in severe DNA damages, cell cycle arrest, and death [40,54–57]. Chemotherapeutics that target Top as poisons convert a transient DNA-enzyme complex into lethal DNA breaks [58–62]. However, topoisomerase inhibitors' activity and their multifaceted binding modes to DNA, the effects, and the modulations they produce on the control of cancer cell division necessitate better understanding to optimize their efficiency.

This review focuses on CuC targeting human Top1 and Top2, the molecular mechanism of induced DNA damages, cell cycle arrest, programmed cell death responses, and emerging research strategies.

2. Copper Complexes as Topoisomerases Inhibitors

DNA topoisomerases have been molecular targets for anticancer agents since their discovery in 1971 [63]. Topoisomerases regulate DNA winding and play essential functions in DNA replication and transcription [59,64]. Topoisomerase 1 (Top1) creates transient single-DNA nicks, while topoisomerases 2 (Top2 α and Top2 β) produce transient double-stranded DNA breaks. Both nuclear Top1 and Top2 are important targets for cancer chemotherapy, and Top inhibitors are used in therapeutic protocols [65–67]. Top inhibitors are classified into two groups: poisons and catalytic inhibitors. Top poisons (or interfacial poisons) stabilize the reversible cleavage complex formed between Top and DNA and form a ternary complex. Top2 catalytic inhibitors can prevent DNA strands cleavage through inhibition of the ATPase activity (novobiocin, merbarone), by impeding ATP hydrolysis to block Top dissociation from the DNA (ICRF-193), or by DNA intercalation at the Top fixation site (aclarubicinet) see [68]. In all cases, inhibitors convert the indispensable nuclear Top enzyme into a killing tool.

Top inhibitors' activity increases upon complexation with copper ion. Top1, Top2, or Top1/2 inhibitors synthesized in the form of copper complexes (CuC) are mostly mononuclear Cu(II) complexes associated with a variety of ligands (Table 1). Different strategies are currently proposed to design and develop Top inhibitory agents based on ligands' properties [69]. If both Top1 and Top2 inhibitors CuC primarily target DNA by a direct interaction through intercalation or cleavage, their antiproliferative activity is reinforced by ROS production and other molecular targets (Table 1) [25,52].

Table 1. Copper complexes inhibitors of topoisomerases: targeted top isoforms, cancer cell lines responses, and molecular mechanisms are summarized. * Tests were realized in vitro with human Top1 or Top2 α/β unless specified. IC50: half-maximal inhibitory concentration. EC50: half-maximal effective concentration. GI50: half-average of growth inhibition.

Ligand Class of Cu-C	Compound Number	Targeted Top(s)	Inhibition of DNA Relaxation Total (μ M) (minimal (μ M))	Inhibition Mechanism	Cancer Cell Lines	IC50 (μ M)	Cell Cycle Arrest	Cell Death Type	Other Specificity	Reference Number
Oxindolimine	1	Top1	50 (25)	Fixation in the DNA Top1 binding site	Neuroblastoma SH-SY5Y Promonocytic U937		G2/M arrest	Apoptosis	ROS induction	[70–73]
Hydrazone with triphenylphosphonium	2	Top1	40	DNA Binding Enzyme complex formation	Lung A549 Prostatic PC-3	4.2 \pm 0.8 3.2 \pm 0.2				[74]
Plumbagin	3	Top1	1.56	DNA intercalation	Breast MCF-7 Colon HCT116 Hepatoma BEL7404 Hepatoma HepG2 Kidney 786-O Lung NCI-H460 Nasopharyngeal cancer CNE2	3.2 \pm 1.1 5.9 \pm 1.4 12.9 \pm 3.6 9.0 \pm 0.7 2.5 \pm 0.9 2.0 \pm 1.2 11.8 \pm 5.9				[75]
Phenanthroline with amino acids	4	Top1	50 (10)	DNA intercalation	Nasopharyngeal cancer HK1	2.2–5.2		Apoptosis		[76]
Pyrophosphate	5	Top1	500	DNA interaction	Ovarian A2780/AD	0.64 \pm 0.12				[77]
Heterobimetallic Cu(II)-Sn2(IV) phenanthroline	6	Top1	20	DNA intercalation cleavage	Breast Zr-75-1 Cervix SiHa Colon HCT15, SW620 Kidney 786-O, A498 Lung Hop-62, A569 Pancreatic MIA PaCa-2	<10 (GI50)				[78]
Analogs					Neuroblastoma SH-SY5Y	2–8		Apoptosis		[79]
Tridentate chiral Schiff base	7, 8	Top1	25 (15)	DNA binding major groove	Hepatoma HuH7 Hepatoma HepG2	25 6.2 \pm 10			ROS Cytokine TGF β mRNA upregulation	[81,82]

Table 1. Cont.

Ligand Class of Cu-C	Compound Number	Targeted Top(s)	Inhibition of DNA Relaxation Total (μM) (minimal (μM))	Inhibition Mechanism	Cancer Cell Lines	IC50 (μM)	Cell Cycle Arrest	Cell Death Type	Other Specificity	Reference Number
Salicylidene	9	Top1	(E. coli) *	DNA binding	Prostatic PC-3	7.3 ± 0.2			antimetastasis	[83]
				DNA cleavage	Breast MCF7	51.1 ± 1.6				[84]
					Colon HT29	16.6 ± 0.6				
					Hepatoma HepG2	2.3 ± 0.1				
					Lung A549	16.8 ± 1.0				
					Ovary A2780	14.6 ± 0.2				
					Prostatic LNCaP	25.4 ± 0.8				
Chalcone-derived Thiosemicarbazone	10	Top1	3 (0.75)	DNA binding DNA cleavage Religation inhibition	Breast MCF-7 Leukemia THP-1	0.16 ± 0.06 0.20 ± 0.06				[85]
Pyridyl-substituted tetrazolopyrimidie	11	Top1	(Molecular docking) *	DNA binding groove mode	Cervix HeLa Colon HCT-15 Lung A549	0.565 ± 0.01 0.358 0.733		Apoptosis	CDK receptor binding	[86]
Tetrazolopyrimidine Diimine		Top1	102 ± 1.1	DNA binding groove mode	Cervical HeLa Colon HCT-15 Lung A549	0.620 ± 0.0013 0.540 ± 0.00015 0.120 ± 0.002		Apoptosis	vEGF receptor binding	[87]
Piperazine	12	Top1	12.5 (5)	DNA binding minor groove					SOD mimic	[88]
Elesclomol	13	Top1	50	Poison	Erythroleukemic K562	0.0075		Apoptosis Necrosis Oxidative stress	Copper chelator Not a substrat for ABC transporters	[89]
Cu(SBCM)2	14	Top1	* (Molecular docking)	DNA intercalation DNA binding	Breast MCF7 Breast MDA-MB-231	27 18.7 ± 3.1	G2/M arrest	Apoptosis	p53 increase No ROS	[90] [91]
TSC and TSC CuC [92–97]										
Pyridine-TSC	15	Top2a	50 (10)		Breast MDA-MB-231 Breast MCF7	1.01 0.0558				[98]
			50	ATP hydrolysis inhibition						[99]
		Top2β	(5)	ATP hydrolysis inhibition						[100]

Table 1. *Cont.*

Ligand Class of Cu-C	Compound Number	Targeted Top(s)	Inhibition of DNA Relaxation Total (μ M) (minimal (μ M))	Inhibition Mechanism	Cancer Cell Lines	IC50 (μ M)	Cell Cycle Arrest	Cell Death Type	Other Specificity	Reference Number
Piperazine-TSC	16	Top2a	0.9 ± 0.7	Potentially catalytic	Breast MCF7 Breast SK-BR-3	4.7 ± 0.3 1.3 ± 0.3				[101,102] [99]
	17	Top2a	4 (2)		Breast MDA-MB-231 Breast MCF7	1.41 (EC50) 0.13 (EC50)				[103]
	17–18	Top2a	25 (10)	ATP hydrolysis inhibition + Poison	Breast HCC 70, HCC 1395, HCC 1500, and HCC 1806 Colon Caco-2, HCT-116 and HT-29	1 to 20 0.83 to 41.2				[104,105]
L- and D-Proline-TSC	19	Top2a	300		Ovarian carcinoma CH1	113 ± 16				[106]
Quinoline-TSC	20	Top2a	0.48	Potentially catalytic	Lymphoma U937	0.48-16.2				[107]
Naphthoquinone-TSC	21	Top2 α	1 mM		Breast MCF7	3.98 ± 1.01		No apoptosis		[108]
	22	Top2a	100 (5)	Poison	Breast MDA-MB-231 Colon HCT116 Keratinocyte HaCaT	1.45 ± 0.07 1.23 ± 0.27 0.65 ± 0.07	G2/M arrest	Apoptosis	DNA synthesis inhibition No ROS	[109]
Bis-TSC					Colon HCT116	Delayed mice xenograft				
Carbohydrazone	23	Top2 α	250 (25)	DNA binding major groove	Breast MCF7 Breast MDA-MB-231 Breast HCC 1937 Breast MX1 Breast MDA-MB-436	9.916 7.557 3.278 4.534 5.249		Apoptosis		[110]
					Breast MX-1	Reduced mice xenograft (83%)				

Table 1. *Cont.*

Ligand Class of Cu-C	Compound Number	Targeted Top(s)	Inhibition of DNA Relaxation Total (μ M) (minimal (μ M))	Inhibition Mechanism	Cancer Cell Lines	IC50 (μ M)	Cell Cycle Arrest	Cell Death Type	Other Specificity	Reference Number
Chromone	24	Top2a	25 (15)	DNA binding major groove	Breast MCF7 Breast Zr-75-1 Colon HT29 Cervix SiHa Kidney A498 Lung A549 Ovary A2780	18.6 (GI 50) 25.2 (GI 50) >80 (GI 50) 34.6 (GI 50) 73.3 (GI 50) 31.7 (GI 50) 17.4 (GI 50)				[111]
Quinolinone Schiff Base	25	Top2 α	9	No intercalation	Hepatic HepG2	17.9 \pm 3.8			DNA synthesis inhibition Slight substrate for ABC transporter	[112]
Bis-pyrazolyl Carboxylate	26	Dual Top1/Top2	(Molecular docking)*	ATP entry (potentially) DNA religation inhibition (potentially)	Hepatic HepG2	3.3 \pm 0.02		Apoptosis DNA replication ROS		[113]

2.1. CuC Top1 Inhibitors

All the structures of CuC Top1 inhibitors are reported in Figure 2 and the main characteristics in Table 1. Oxindolimine-Cu(II) Top1 inhibitors such as **1** are planar copper compounds [70] that do not permit enzyme-DNA complex formation [71–73]. Besides, they produce ROS [70]. Cu(II) derivative complexes of the hydrazone ligand with triphenylphosphonium moiety **2** can bind DNA and the Top enzyme [74]. Plumbagin-Cu(II) **3** selectively intercalates into DNA [75]. The latter compound [75] and the phenanthroline-Cu(II) complexes modulated by amino acids **4** [76] can induce cancer cell apoptosis via mitochondrial signaling. Copper pyrophosphate-bridged binuclear complex **5** interacts with DNA, and based on the redox chemistry of copper, induces significant oxidative stress in cancer cell lines [77].




Figure 2. Structure of Cu(II) complexes as Top1 inhibitors.

In the heterobimetallic Cu(II)-Sn₂(IV) (copper/tin) complex **6**, the planar phenanthroline heterocyclic ring approaches the Top–DNA complex Cu(II)-Sn₂(IV) toward the DNA cleavage site and forms a stable complex with Top1 [78,79]. Other Cu(II)-Sn₂(IV) analogs induce apoptosis [80]. Chiral monometallic or heterobimetallic complexes **7** and **8** with tridentate chiral Schiff base–ONO-ligand are DNA groove binders and produce ROS [81,82].

Salicylidene-Cu(II) derivative **9** of 2-[2-bromoethyliminomethyl] phenol [83,84] is a bifunctional drug that inhibits both cancer cell growth and metastasis.

Chalcone-derived thiosemicarbazone (TSC) Cu(II) complex **10** prevents the DNA cleavage step of the Top1 catalytic cycle and DNA relegation [85].

Tetrazolo[1,5-*a*]pyrimidine-based Cu(II) complexes **11** have a square planar geometry, and despite their high capability to inhibit Top1, interact with CDK for **11** [86] and VEGF receptors for an analog of **11** [87]. Binuclear Cu(II) dipeptide piperazine-bridged complex **12** recognizes specific sequences in the DNA, oxidatively cleaves DNA, and displays superoxide dismutase (SOD) activity [88].

Derived from elesclomol (in clinical trials: phase 3 against melanoma and randomized phases 2 and 3 for the treatment of a variety of other cancers), the elesclomol-Cu(II) complex **13** inhibits Top1 and induces apoptosis in cancer cells [89].

As recently studied, Cu(II)(SBCM)₂ **14** derived from S-benzylidithiocarbazole and 3-acetylcoumarin intercalates into DNA, induces ROS production, and has an antiproliferative activity in breast cancer lines [90,91].

2.2. CuC Top2 α Inhibitors

Due to its cell cycle phase dependence and its high expression in proliferating cells, the Top2 α isoform is primarily targeted by copper complexes (CuC), whereas Top2 β remains unchanged during the course of the cell cycle [66]. Another reason to limit the clinical application of Top2 β inhibitors is the strong unwanted side effects produced (secondary leukemia, myelodysplastic syndrome (MDS), and cardiac toxicity [92,93]).

The main characteristics and structures of CuC Top2 inhibitors are reported in Figure 3 and Table 1. Several α -(N)-heterocyclic thiosemicarbazone (TSC) CuC [94,95] present a greater inhibitory effect on Top2 α than corresponding TSC ligands alone [96,97] due to a square planar structure around the Cu(II) ion. A specific subset of pyridine-TSC CuC 15 inhibits Top2 α [98] acting as ATP hydrolysis inhibitors in a non-competitive mode [94,99,100]. Another pyridine-TSC CuC inhibits Top2 β [100]. Molecular modeling supports the binding of the complexes near but outside the ATP binding pocket in communication with the DNA cleavage/ligation site of Top2. Piperazine-TSCs based CuC 16 inhibit Top2 α [101,102] by a strong interaction with the ATP-binding pocket residues [99] without ROS production [102]. Thiazole-TSC CuC 17 and 18 are Top2 α catalytic inhibitors [103, 104] or poisons [105]. The highly water-soluble proline-TSC CuC series 19 inhibit Top2 α and cell proliferation [106]. Quinoline-TSC CuC 20 interact with the DNA phosphate group preventing relegation. The presence of two methyl groups on the terminal nitrogen is responsible for high activity and confers a cationic nature responsible for easier passive access into the cell [107].





Figure 3. Structure of Cu(II) complexes as Top2 inhibitors.

Non-heterocycle naphthoquinone-TSC CuC 21 [108] and bis-TSC CuC 22 [109] are Top2 α inhibitors acting as poisons [109]; they induce apoptosis in various human cancer cell lines and delay colorectal growth of carcinoma xenografts in mice [109]. Carbohydrazone CuC 23 [110] is a Top2 α inhibitor that binds DNA, induces apoptosis, and reduces mice xenograft (83% after a treatment of 2 mg/kg). Chiral chromone Cu(II)/Zn(II) 24 [111] revealed catalytic inhibition of Top2 α with DNA binding in the major groove. Quinolinone CuC 25 [112] inhibit Top2 α and DNA synthesis without DNA intercalation and are only minimized PGP (P-glycoprotein efflux transporter) substrates.

2.3. CuC Dual Top1/Top2 α Inhibitors

Heteroleptic Cu(I) complexes of the bis-pyrazolyl carboxylate ligand with auxiliary phosphine **26** (Figure 4) may inhibit Top1 by blocking the reagation step and inhibit Top2 α by preventing ATP hydrolysis, as proposed by molecular docking analysis. They also perturb DNA replication, generate ROS, and induce apoptosis [113].




bis-pyrazolyl-Cu(I) complex, **26**

Figure 4. Structure of Cu(I) complex as a Top1/2 α dual inhibitor.

3. Cell Cycle Regulation by Copper Complexes and Top Inhibitors

CuC inhibitors targeting Top1 [72,90] or Top2 [109] as DNA-damaging drugs or poisons arrest cancer cells in G2/M (Table 1). This common G2/M arrest involves the activation of two different cell cycle pathways: the DNA damage response (DDR) and the decatenation checkpoint.

Both Top1 and Top2 CuC inhibitors produce DNA damages. Top2 poisons prevent DNA reagation and stabilize an enzyme–DNA complex with the double-stranded cleaved DNA [114]. Top1 poisons induce single-stranded DNA breaks and associated signaling cascades. The collision between the Top1 cleavage complexes and the DNA replication forks ends up generating double-strand breaks [115] (Figure 5A). Top1- and Top2-induced DNA breaks trigger a DDR executed by ATM-, ATR-, and DNA-PK-related kinases, and an arrest of the cell cycle machinery [116–118]. ATM- and ATR-dependent phosphorylations of p53, Chk1, and Chk2 regulate the G1/S, S, or G2/M cell cycle checkpoints. Chk1 and Chk2 inhibit Cdc25 phosphatases (A,B,C) required for Cdks activation. Phosphorylated and ubiquitinated Cdc25A (Ser123) is degraded, leading to the absence of activation of the Cdk2/Cyclin E and the Cdk4/cyclin D complexes and followed by an arrest in G1/S. Phosphorylated Cdc25C (Ser216) binds to 14-3-3, prevents Cdk1/Cyclin B (MPF) activation, and induces a G2/M arrest (Figure 5A). Cdc25B inactivation also results in a G2 arrest [119,120]. The DNA damage-induced cell cycle arrest in G1 is dependent on p53 phosphorylation by ATM (Ser15) and Chk2 (Ser20) but arrest in S and G2 phases is p53-independent [121–124]. Phosphorylated p53 dissociates from MDM2 and activates the transcription of Cdk inhibitor p21WAF1 [125,126]. In several CuC (Top1 DNA binding CuC inhibitors [72,82,88] and a dual Top1/2 inhibitor with heteroleptic CuC [113]), Cu(II) exhibits a high redox potential and reinforces DDR activation by ROS production. ROS are also involved in a G2/M arrest through the decrease in Cdc25C [127] and Cdc25A levels [128], the activation of Chk1 [129] and Chk2 [130], and genomic instability through induced-DNA damages [131] (Figure 5A).

**Figure 5. Cont.**




Figure 5. Molecular mechanisms and treatment perspectives for copper complexes (CuC) drugs. (A) Molecular checkpoints and networks involved in DNA damage (red), cell cycle regulation (yellow), and death response (violet) triggered by topoisomerase inhibitors (poison and catalytic), including CuC of topoisomerase inhibitors. (B) Treatment perspectives alone or in association with other chemotherapeutics (see text for more details).

By contrast to poisons, Top2 catalytic inhibitors do not form cleavable complexes. They function by enzymatic activity deprivation and cell cycle arrest in G2 through a decatenation checkpoint distinct from the DNA damage checkpoint. To delay the mitotic entry, an insufficient decatenation engages molecular components from the DDR and the spindle assembly checkpoint (SAC) (Rad9a, ATR, and BRCA1), SUMOylation and phosphorylation of Top2, the p38 and the MAPK pathways, and several decatenation checkpoint effectors but not p53 [66,132–136] (Figure 5A).

Cell cycle checkpoint effectors arrest DNA-damaged cells and induce their death providing that cell cycle regulatory networks are effective. Cell cycle checkpoint effectors integrity influences responses to Top2 inhibitors [137]. Besides, cancer disease is associated with multiple overexpression and mutations [138] in Cdc25 [139,140] and p53 [141,142], to a loss of Cdk inhibitors expression and/or overexpression of cell cycle-regulated protein [143,144], Top deregulation, and multidrug resistance [145–147]. Moreover, cell cycle variation of Top2 α is regulated by post-translational modifications that represent potential targets. These alterations include ubiquitination by Cdk-1 [148], sumoylation [149], phosphorylation by polo-like kinase 1, Cdc7 [150], protein kinase C, Ca/calmodulin-dependent kinase II, and casein kinase [151], and the association with 14-3-3 [152]. Rewiring cellular pathways leading to cell death is a challenge that requires targeting specific molecular checkpoint effectors [153]. For example, a mutated p53 pathway arrests the cell cycle but avoids DDR-induced cell death [154]. Some anticancer therapeutic strategies (e.g., Chk1/2 pathways targeting drugs associated with DNA-damaging drugs) can force cancer cells to bypass S and G2/M arrest, enter mitosis with damaged DNA, and finally undergo a mitotic catastrophe and death [155]. ATR inhibition is another strategy to overcome the resistance of BRCA-deficient cancers [156].

4. Programmed Cell Death Engaged by Copper Complexes and Top Inhibitors

Multiple stress factors ranging from various cell damages, ATP levels, and specific pathways (e.g., caspases) determine the type of cell death [157]. Most Top1 CuC inhibitors that interact with DNA [70,76,79,86,87,90], Top1 poison [89], Top2 α CuC poison [109], or dual Top1/Top2 inhibitor [113] trigger apoptotic programmed cell death. Genetic damages and oxidative stress activate an intrinsic mitochondrial response [158]. Pro-apoptotic members of the Bcl-2 family (Bid, Noxa, Puma, BAX, BAK) neutralize the anti-apoptotic members (Bcl-2, Bcl-xL, and Mcl-1), disrupt the mitochondrial outer membrane, and allow cytoplasmic cytochrome-c release. The binding of cytochrome c to the apoptotic protease activating factor-1 (Apaf-1), ATP, and the pro-caspase-9 create the apoptosome protein complex. Pro-caspase 9 is cleaved into its active caspase-9 form, which in turn cleaves pro-caspase-3 into caspase-3 effector, and the downstream executor caspase-7. SMAC (second mitochondria-derived activator of caspases), and Omi/HtrA2 (high-temperature requirement protein A2) are simultaneously released from mitochondria and deactivate the IAPs factors (inhibitors of apoptosis proteins). p53, activated by the DNA damage, contributes to apoptosis through the translation of several pro-apoptotic members of the Bcl-2 family (Bid, Puma) that inhibit the pro-survival action of Bcl-2 on BAX (Figure 5A). Most cancer cells evade apoptosis through caspase inhibition, upregulation of Bcl-2 (in more than 50% of all types of cancers), and loss of BAX/BAK and become resistant to anticancer drugs [159].

A Top1 DNA-damaging CuC inhibitor induces necrotic cell death. To facilitate cell destruction, necrosis is activated by ROS or ATP metabolic stresses in crosstalk with apoptosis [160]. When the intracellular energy/ATP level is low, the apoptotic cell death is converted into necrosis [161] (Figure 5A). However, necrosis releases pro-inflammatory and tumor-promoting cytokine HMGB1 [162] into the extracellular space reported to stimulate inflammation and angiogenesis, and promote tumor progression [163].

Apoptosis and necrosis often co-exist with another cell death with controversial pro-death and pro-survival functions: autophagy [164]. Up to the current study, no CuC Top inhibitors are involved in autophagic or necroptotic programmed cell death (Table 1). However, some CuC trigger stress-mediated protective autophagy in response to ROS that impedes apoptosis and creates survival

of malignant cells [165]. Moreover, topoisomerase inhibition-induced autophagy is associated with cancer resistance [166].

5. Future Strategies for Copper Complexes as Top Inhibitors in Cancer Cell Treatments

The development of new effective anticancer drugs is a major research area against the continuing increase in cancers worldwide. Top inhibitors used in chemotherapy are limited in number [61,167,168]. Top1 inhibitors' camptothecin derivatives used are irinotecan (colorectal [169], pancreatic (in combination) [170], and small cell lung cancers (in clinical trials and in combination) [171,172]), and topotecan (ovarian [173,174], cervical [175], and small cell lung cancers [176]). Top2 anticancer drugs commonly used are from the anthracycline group such as doxorubicin (acute leukemia [177], lymphomas [178], sarcomas [179,180], and solid tumors [181]), epirubicin (breast cancer [182]), valrubicin (bladder cancer [183]), and idarubicin (acute myeloid leukemia [184]), from the anthracenedione classes: mitoxantron and pixantron (lymphoma, [185–187]), and from the epipodopodophyllotoxins group such as etoposide (testicular [188] and small cell lung cancers [189]) and teniposide (brain [190] and small cell lung [191] cancers, acute lymphocytic leukemia [192]). Only a few numbers of Top1 inhibitors are in clinical trials including the promising indenoisoquinoline derivatives LMP400 (Indotecan), LMP776 (Indimitecan) (phase I), and LMP744 examined in a phase I study on lymphoma in dogs [193]. In addition to better stability, and milder side effects, they can escape ABC transporter efflux and the drug resistance mechanism, as Elesclomol-CuC Top complexes 13 [89] or Quinolinone-CuC 25 [112]. Perspectives to use CuC of Top inhibitors in clinical trials are summarized in Figure 5B. Development and optimization in CuC of Top inhibitors imply structure modifications that must encompass several specific strategies [194], such as scaffold hopping [195], pharmacophore hybridization [196], bioisosteric replacement [197], and conformational restrictions. Generally, a rigidification of the ligand heterocycle structure with a copper metal [78] provides a planar configuration that facilitates DNA intercalation and Top-DNA ternary complex formation compared to the molecular backbone alone.

Top inhibitors in clinical use and particularly Top poison display unwanted drawbacks, such as cumulative cardiotoxicity in long-term protocols, secondary malignancies, and drug resistance [198]. A therapeutic option would be to use preferentially catalytic Top agents that disturb the catalytic cycle without the formation of a ternary complex. CuC Top catalytic inhibitors, listed in Table 1, exhibit high antitumor effects on cancer cell lines and for some compounds on tumor growth in animal models, compared to their respective ligands (see Table 1). They constitute a reservoir of anticancer drugs. For example, TSC-based CuC Top2 inhibitors (Figure 3) [98,102,103,105,107] have demonstrated strong inhibition of tumor growth compared to TSC derivatives currently used in cancer chemotherapies [199].

Considering that cancer is a multigenetic and multifactorial disease that recruits numerous molecular effectors, monotherapies (based on Top inhibitors) do not provide the optimal curative effects. Combination therapy with a few numbers of therapeutics against two or more biotargets is the base of promising treatments such as the association of a Top 2 inhibitor (vosaroxin) with a DNA methyltransferase inhibitor (decitabine) in AML [200,201]. Inhibitors of Top1 and Top2, currently developed, also exert their effect against other cancer-related targets [202]. Dual Top inhibitors, e.g., Top 1/2 [203], Top2/microtubule [204], or Top2/histone deacetylase [205], may exert improved efficacy. Besides, Top1 inhibitors are nonspecific RNA polymerase inhibitors. An RNA Pol1-mediated ribosomal RNA gene increase is involved in cancer progression, through the control of cellular checkpoints and chromatin structure and is, therefore, an interesting co-target [206]. CuC dual Top inhibitors display a high antiproliferative activity. Particularly, some CuC and non-CuC are dual inhibitors of Top1 and superoxide dismutase agonist [88,207,208] or Cdk receptor, like VEGF inhibitors, involved in cancer cells proliferation [86,87,209,210] (Figure 5B). Another strategy to improve therapies is the association of a CuC with a TDP1/2 (tyrosyl-DNA-phosphodiesterase 1/2) inhibitor. TDP1/2 are enzymes responsible for the reparation of DNA breaks induced by topoisomerase poisons [57,211,212]. TDP1/2 inhibitors are capable of improving cancer cells' sensitivity to these poisons [213].

Autophagy, an essential mechanism for cell integrity and survival, is stimulated in cancer cells under several chemotherapeutic drugs and acts as an unwanted protective system towards tumor cells. Association of specific autophagic inhibitors with Cu-C treatment (disulfiram) in non-small cell lung cancer [214] has proven to be a novel efficient strategy to enhance apoptosis in cancer therapy.

Immunogenic cell death is an important mechanism used in chemotherapy. Association of CuC with immune checkpoint therapies is certainly a new avenue in cancer treatment. CuC and non-CuC Top inhibitors induce DNA damages and are linked to adaptive and innate immunities [215]. Top poisons promote immunogenicity in various ways [216]. Top1 poison camptothecin enhances the adaptive immune response [217]. Top inhibitors also increase chromosomal instability and mutations accumulated by cancer cells [59,218]. Consequently, due to their high number of mutations, tumors display more neoantigens presented at their surface by the major histocompatibility complex class I (MHCI) and recruit lymphocytes T harboring TCR (T cell receptor) and CD8 co-receptor (adaptive immunity). This response is counterbalanced by the overexpression of immune checkpoint modulators, such as the immune-suppressive ligand PD-L1 (programmed death-ligand 1) targeted in immune therapies [219] (Figure 5B). DNA-damaging agents such as Top2 poison anthracycline also interfere with the innate immune response. They enhance the malignant formation of cytosolic bircatenated DNA fragments that activate the cyclic GMP-AMP synthase-stimulator of the interferon (IFN) gene pathway (cGAS-STING) and initiate innate anti-cancer immunity. cGAS-STING agonist serves as a sensitizer in immunotherapies [220]. Top1-DNA covalent cleavage complex enables cGAS-mediated cytoplasmic chromatin recognition and immune checkpoint response [221] (Figure 5B). Top2 inhibitors teniposide and doxorubicin potentiate the therapeutic immune checkpoint blockade therapies based on anti-PD-1 (programmed cell death 1) in multiple types of mouse tumor models [222,223]. Besides, ROS produced by Top inhibitors alter the molecular pattern recognized as immunogenic structures and enhance apoptosis [224] (Figure 5B).

As DDR gene mutations exist in a large range of tumor types, the determination of tumor-specific mutations is another accurate strategy to generate chemotypes with beneficial efficacies superior to adverse effects [225,226]. In each tumor, the signaling components of the DDR exhibit numerous defects that result in a unique mutational signature [227]. Cancer cells with defects in their homologous recombination mechanism are more sensitive to Top2 inhibitory therapies that generate DNA double-strand breaks [228]. Moreover, the prediction of anticancer treatments determined by the clinical stage and the pathological features of the tumor does not always ascertain a cancer death response. Cellular biomarkers that may predict sensitivity or resistance to therapy based on DNA damage induced by Top inhibitors would be useful. Insights into the Top2 regulatory mechanisms have identified genetic markers to allow the prediction of an overcome treatment with a Top inhibitor. γ-H2AX is a DNA-damaged marker, recruited on DNA breaks after Top poison action, currently evaluated [229]. Schlaffen is also a promising marker for an accurate response to Top1 and Top2 inhibitors, especially for colon and ovarian adenocarcinomas [56,230] (Figure 5B).

Recently, cancer cells were targeted specifically by a Top2 inhibitor, etoposide, attached to a single-stranded oligonucleotide with a complementary sequence to a DNA cleavage hotspot corresponding to a translocated region only present in promyelocytic leukemia cells [231].

Finally, to overcome toxicity to normal cells, Top drugs could be attached to vehicles. Top2 inhibitors delivery has been optimized using liposomes [232], micelles [233], or functionalized nanoparticles [234] (Figure 5B).

Topoisomerases are present in mitochondria where they participate in mitochondrial DNA replication and transcription. Mitochondrial Top1 isoform (Top1mt) is involved in the metabolism of cancer cells providing energy to tumors surrounded by a nutrient-low microenvironment. Exposures to a Top1 inhibitor (lamellarin D) or Top2 inhibitors (doxorubicin or fluoroquinolones) exert mitochondrial toxicity [235]. However, the loss of Top1mt in liver cancers correlates with increased survival of hepatocellular carcinoma patients, showing that co-targeting Top1mt in addition to nuclear topoisomerases is another option for anticancer therapies [236].

6. Conclusions

In a multifactorial disease such as cancer, Top inhibitors are efficient anticancer compounds used in monotherapy or polypharmacological strategies. They certainly have to target closely related modulators of the cellular checkpoints' networks. CuC Top inhibitors are particularly adapted to fulfill this role. A perspective in anticancer strategy is to increase and to enlarge this family of highly active anticancer drugs.

Author Contributions: Writing—original draft preparation: C.M., K.C., A.M. and L.P. All authors have read and agreed to the published version of the manuscript.

Funding: C.M. is a recipient from a doctoral fellowship from the French ministry. The scientific supports were provided by the “Centre National de la Recherche Scientifique”, “Université de Lille”, and the “Ligue Contre le Cancer”.

Acknowledgments: The authors are thankful to the Research Federation FRABio (FR 3688, FRABio, Biochimie Structurale et Fonctionnelle des Assemblages Biomoléculaires) and to E. Germain (Inserm U1003) for reading the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Rosenberg, B.; VanCamp, L.; Trosko, J.E.; Mansour, V.H. Platinum compounds: A new class of potent antitumour agents. *Nature* **1969**, *222*, 385–386. [[CrossRef](#)] [[PubMed](#)]
- Alderden, R.A.; Hall, M.D.; Hambley, T.W. The discovery and development of cisplatin. *J. Chem. Educ.* **2006**, *83*, 728–734. [[CrossRef](#)]
- Dilrubia, S.; Kalayda, G.V. Platinum-based drugs: Past, present and future. *Cancer Chemother. Pharm.* **2016**, *77*, 1103–1124. [[CrossRef](#)] [[PubMed](#)]
- Bergamo, A.; Dyson, P.J.; Sava, G. The mechanism of tumour cell death by metal-based anticancer drugs is not only a matter of DNA interactions. *Coord. Chem. Rev.* **2018**, *360*, 17–33. [[CrossRef](#)]
- Dasari, S.; Tchounwou, P.B. Cisplatin in cancer therapy: Molecular mechanisms of action. *Eur. J. Pharm.* **2014**, *740*, 364–378. [[CrossRef](#)]
- Manohar, S.; Leung, N. Cisplatin nephrotoxicity: A review of the literature. *J. Nephrol.* **2018**, *31*, 15–25. [[CrossRef](#)]
- Herradón, E.; González, C.; Uranga, J.A.; Abalo, R.; Martín, M.I.; López-Miranda, V. characterization of cardiovascular alterations induced by different chronic cisplatin treatments. *Front. Pharm.* **2017**, *8*, 196–211. [[CrossRef](#)]
- Shen, D.W.; Pouliot, L.M.; Hall, M.D.; Gottesman, M.M. Cisplatin resistance: A cellular self-defense mechanism resulting from multiple epigenetic and genetic changes. *Pharm. Rev.* **2012**, *64*, 706–721. [[CrossRef](#)]
- Chen, S.H.; Chang, J.Y. New insights into mechanisms of cisplatin resistance: From tumor cell to microenvironment. *Int. J. Mol. Sci.* **2019**, *20*, 4136. [[CrossRef](#)]
- Obrist, F.; Michels, J.; Durand, S.; Chery, A.; Pol, J.; Levesque, S.; Joseph, A.; Astesana, V.; Pietrocola, F.; Wu, G.S.; et al. Metabolic vulnerability of cisplatin-resistant cancers. *EMBO J.* **2018**, *37*, e98597. [[CrossRef](#)]
- Amable, L. Cisplatin resistance and opportunities for precision medicine. *Pharm. Res.* **2016**, *106*, 27–36. [[CrossRef](#)] [[PubMed](#)]
- Galluzzi, L.; Senovilla, L.; Vitale, I.; Michels, J.; Martins, I.; Kepp, O.; Castedo, M.; Kroemer, G. Molecular mechanisms of cisplatin resistance. *Oncogene* **2012**, *31*, 1869–1883. [[CrossRef](#)] [[PubMed](#)]
- Housman, G.; Byler, S.; Heerboth, S.; Lapinska, K.; Longacre, M.; Snyder, N.; Sarkar, S. Drug resistance in cancer: An overview. *Cancers* **2014**, *6*, 1769. [[CrossRef](#)] [[PubMed](#)]
- Martinho, N.; Santos, T.; Florindo, H.F.; Silva, L.C. Cisplatin-membrane interactions and their influence on platinum complexes activity and toxicity. *Front. Physiol.* **2019**, *9*, 1898–1913. [[CrossRef](#)]
- Zeng, L.; Gupta, P.; Chen, Y.; Wang, E.; Ji, L.; Chao, H.; Chen, Z.-S. The development of anticancer ruthenium(ii) complexes: From single molecule compounds to nanomaterials. *Chem. Soc. Rev.* **2017**, *46*, 5771–5804. [[CrossRef](#)]

16. Zhang, P.; Sadler, P.J. Advances in the design of organometallic anticancer complexes. *J. Org. Chem.* **2017**, *83*, 5–14. [[CrossRef](#)]
17. Jaouen, G.; Vessières, A.; Top, S. Ferrocifen type anti cancer drugs. *Chem. Soc. Rev.* **2015**, *44*, 8802–8817. [[CrossRef](#)]
18. Gianferrara, T.; Bratsos, I.; Alessio, E. A categorization of metal anticancer compounds based on their mode of action. *Dalton Trans.* **2009**, *37*, 7588–7598. [[CrossRef](#)]
19. Hartinger, C.G.; Dyson, P.J. Bioorganometallic chemistry from teaching paradigms to medicinal applications. *Chem. Soc. Rev.* **2009**, *38*, 391–401. [[CrossRef](#)]
20. Wambang, N.; Schifano-Faux, N.; Martoriati, A.; Henry, N.; Baldeyrou, B.; Bal-Mahieu, C.; Bousquet, T.; Pellegrini, S.; Meignan, S.; Cailliau, K.; et al. Synthesis, structure, and antiproliferative activity of ruthenium(II) arene complexes of indenoisoquinoline derivatives. *Organometallics* **2016**, *35*, 2868–2872. [[CrossRef](#)]
21. Wambang, N.; Schifano-Faux, N.; Aillerie, A.; Baldeyrou, B.; Jacquet, C.; Bal-Mahieu, C.; Bousquet, T.; Pellegrini, S.; Ndifon, T.P.; Meignan, S.; et al. Synthesis and biological activity of ferrocenyl indeno[1,2-c]isoquinolines as topoisomerase II inhibitors. *Bioorg. Med. Chem.* **2016**, *24*, 651–660. [[CrossRef](#)] [[PubMed](#)]
22. Komeda, S.; Casini, A. Next-generation anticancer metallodrug. *Curr. Top. Med. Chem.* **2012**, *12*, 219–235. [[CrossRef](#)] [[PubMed](#)]
23. Mejía, C.; Ortega-Rosales, S.; Ruiz-Azuara, L. Mechanism of action of anticancer metallodrugs. In *Biomedical Applications of Metals*; Rai, M., Ingle, A., Medici, S., Eds.; Springer: Berlin/Heidelberg, Germany, 2018; Volume 10, pp. 213–234.
24. Denoyer, D.; Clatworthy, S.A.S.; Cater, M.A. Copper complexes in cancer therapy. *Met. Ions Life Sci.* **2018**, *18*, 469–506.
25. Santini, C.; Pellei, M.; Gandin, V.; Porchia, M.; Tisato, F.; Marzano, C. Advances in copper complexes as anticancer agents. *Chem. Rev.* **2014**, *114*, 815–862. [[CrossRef](#)] [[PubMed](#)]
26. Jungwirth, U.; Kowol, C.R.; Keppler, B.K.; Hartinger, C.G.; Berger, W.; Heffeter, P. Anticancer activity of metal complexes: Involvement of redox processes. *Antioxid. Redox Signal.* **2011**, *15*, 1085–1127. [[CrossRef](#)] [[PubMed](#)]
27. Tardito, S.; Marchiò, L. Copper compounds in anticancer strategies. *Curr. Med. Chem.* **2009**, *16*, 1325–1348. [[CrossRef](#)] [[PubMed](#)]
28. Marzano, C.; Pellei, M.; Tisato, F.; Santini, C. Copper complexes as anticancer agents. *Anticancer Agents Med. Chem.* **2009**, *9*, 185–211. [[CrossRef](#)] [[PubMed](#)]
29. Kellett, A.; Molphy, Z.; McKee, V.; Slator, C. Recent advances in anticancer copper compounds. In *Metal-Based Anticancer Agents*; Vessieres, I.A., Meier-Menches, S.M., Casini, A., Eds.; Royal Society of Chemistry, RSC Metallobiology: London, UK, 2019; Volume 14, pp. 91–119.
30. Hordjewska, A.; Popiółek, L.; Kocot, J. The many “faces” of copper in medicine and treatment. *Biometals* **2014**, *27*, 611–621. [[CrossRef](#)] [[PubMed](#)]
31. Urso, E.; Maffia, M. Behind the link between copper and angiogenesis: Established mechanisms and an overview on the role of vascular copper transport systems. *J. Vasc. Res.* **2015**, *52*, 172–196. [[CrossRef](#)] [[PubMed](#)]
32. Lowndes, S.A.; Harris, A.L. The role of copper in tumour angiogenesis. *J. Mammary Gland. Biol. Neoplasia* **2005**, *10*, 299–310. [[CrossRef](#)] [[PubMed](#)]
33. Hu, G.F. Copper stimulates proliferation of human endothelial cells under culture. *J. Cell. Biochem.* **1998**, *69*, 326–335. [[CrossRef](#)]
34. Yoshida, D.; Ikeda, Y.; Nakazawa, S. Quantitative analysis of copper, zinc and copper/zinc ratio in selected human brain tumors. *J. Neurooncol.* **1993**, *16*, 109–115. [[CrossRef](#)] [[PubMed](#)]
35. Geraki, K.; Farquharson, M.J.; Bradley, D.A. Concentrations of Fe, Cu and Zn in breast tissue: A synchrotron XRF study. *Phys. Med. Biol.* **2002**, *47*, 2327–2339. [[CrossRef](#)] [[PubMed](#)]
36. Nayak, S.B.; Bhat, V.R.; Upadhyay, D.; Udupa, S.L. Copper and ceruloplasmin status in serum of prostate and colon cancer patients. *Indian J. Physiol. Pharm.* **2003**, *47*, 108–110.
37. Díez, M.; Arroyo, M.; Cerdà, F.J.; Muñoz, M.; Martin, M.A.; Balibrea, J.L. Serum and tissue trace metal levels in lung cancer. *Oncology* **1989**, *46*, 230–234. [[CrossRef](#)]

38. Kaifa, G.D.; Saouli, Z.; Diamantidis, M.D.; Kontoninas, Z.; Voulgaridou, V.; Raptaki, M.; Arampatzis, S.; Chatzidimitriou, M.; Perifanis, V. Copper levels in patients with hematological malignancies. *Eur. J. Intern. Med.* **2012**, *23*, 738–741. [CrossRef]
39. Baldari, S.; Di Rocco, G.; Toietta, G. Current biomedical use of copper chelation therapy. *Int. J. Mol. Sci.* **2020**, *21*, 1069. [CrossRef]
40. Denoyer, D.; Masaldan, S.; La Fontaine, S.; Cater, M.A. Targeting copper in cancer therapy: ‘Copper That Cancer’. *Metallooms* **2015**, *7*, 1459–1476. [CrossRef]
41. Cater, M.A.; Pearson, H.B.; Wolyniec, K.; Klaver, P.; Bilandzic, M.; Paterson, B.M.; Bush, A.I.; Humbert, P.O.; La Fontaine, S.; Donnelly, P.S.; et al. Increasing intracellular bioavailable copper selectively targets prostate cancer cells. *ACS Chem. Biol.* **2013**, *8*, 1621–1631. [CrossRef]
42. Cater, M.A.; Haupt, Y. Clioquinol induces cytoplasmic clearance of the X-linked inhibitor of apoptosis protein (XIAP): Therapeutic indication for prostate cancer. *Biochem. J.* **2011**, *436*, 481–491. [CrossRef]
43. Cherian, V.T.; Wang, Y.; Muthu, M.; Jamal, S.; Chen, D.; Yang, H.; Polin, L.A.; Tarca, A.L.; Pass, H.I.; Dou, Q.P.; et al. Disulfiram suppresses growth of the malignant pleural mesothelioma cells in part by inducing apoptosis. *PLoS ONE* **2014**, *9*, e93711. [CrossRef] [PubMed]
44. Duan, L.; Shen, H.; Zhao, G.; Yang, R.; Cai, X.; Zhang, L.; Jin, C.; Huang, Y. Inhibitory effect of Disulfiram/copper complex on non-small cell lung cancer cells. *Biochem. Biophys. Res. Commun.* **2014**, *446*, 1010–1016. [CrossRef] [PubMed]
45. Jivan, R.; Damelin, L.H.; Birkhead, M.; Rousseau, A.L.; Veale, R.B.; Mavri-Damelin, D. Disulfiram/copper-disulfiram damages multiple protein degradation and turnover pathways and cytotoxicity is enhanced by metformin in oesophageal squamous cell carcinoma cell lines. *J. Cell. Biochem.* **2015**, *116*, 2334–2343. [CrossRef] [PubMed]
46. Safi, R.; Nelson, E.R.; Chitreni, S.K.; Franz, K.J.; George, D.J.; Zalutsky, M.R.; McDonnell, D.P. Copper signaling axis as a target for prostate cancer therapeutics. *Cancer Res.* **2014**, *74*, 5819–5831. [CrossRef]
47. Wang, F.; Jiao, P.; Qi, M.; Frezza, M.; Dou, Q.P.; Yan, B. Turning tumor-promoting copper into an anti-cancer weapon via high-throughput chemistry. *Curr. Med. Chem.* **2010**, *17*, 2685–2698. [CrossRef]
48. Zhang, Z.; Wang, H.; Yan, M.; Wang, H.; Zhang, C. Novel copper complexes as potential proteasome inhibitors for cancer treatment. *Mol. Med. Rep.* **2017**, *15*, 3–11. [CrossRef]
49. Krasnovskaya, O.; Naumov, A.; Guk, D.; Gorelkin, P.; Erofeev, A.; Beloglazkina, E.; Majouga, A. Copper Coordination Compounds as Biologically Active Agents. *Int. J. Mol. Sci.* **2020**, *21*, 3965. [CrossRef]
50. Brissos, R.F.; Caubet, A.; Gamez, P. Possible DNA-interacting pathways for metal-based compounds exemplified with copper coordination compounds. *Eur. J. Inorg. Chem.* **2015**, *16*, 2633–2645. [CrossRef]
51. Kagawa, T.F.; Geierstanger, B.H.; Wang, A.H.J.; Ho, P.S. Covalent modification of guanine bases in double-stranded DNA. *J. Biol. Chem.* **1991**, *266*, 20175–20184.
52. Ceramella, J.; Mariconda, A.; Iacopetta, D.; Saturnino, C.; Barbarossa, A.; Caruso, A.; Rosano, C.; Sinicropi, M.S.; Longo, P. From coins to cancer therapy: Gold, silver and copper complexes targeting human topoisomerases. *Bioorg. Med. Chem. Lett.* **2020**, *30*, 126905–126916. [CrossRef]
53. Shobha Devi, C.; Thulasiram, B.; Aerva, R.R.; Nagababu, P. Recent advances in copper intercalators as anticancer agents. *J. Fluoresc.* **2018**, *28*, 1195–1205. [CrossRef] [PubMed]
54. Liang, X.; Wu, Q.; Luan, S.; Yin, Z.; He, C.; Yin, L.; Zou, Y.; Yuan, Z.; Li, L.; Song, X.; et al. A comprehensive review of topoisomerase inhibitors as anticancer agents in the past decade. *Eur. J. Med. Chem.* **2019**, *171*, 129–168. [CrossRef] [PubMed]
55. Cuya, S.M.; Bjornsti, M.A.; van Waardenburg, R.C.A.M. DNA topoisomerase-targeting chemotherapeutics: What’s new? *Cancer Chemother. Pharmacol.* **2017**, *80*, 1–14. [CrossRef] [PubMed]
56. Thomas, A.; Pommier, Y. Targeting topoisomerase I in the era of precision medicine. *Clin. Cancer Res.* **2019**, *25*, 6581–6589. [CrossRef] [PubMed]
57. Pommier, Y. Drugging topoisomerase: Lessons and challenges. *ACS Chem. Biol.* **2013**, *8*, 82–95. [CrossRef] [PubMed]
58. Bjornsti, M.A.; Kaufmann, S.H. Topoisomerase and cancer chemotherapy: Recent advances and unanswered questions. *F1000 Res.* **2019**, *8*. [CrossRef]
59. Pommier, Y.; Sun, Y.; Huang, S.N.; Nitiss, J.L. Roles of eukaryotic topoisomerases in transcription, replication and genomic stability. *Nat. Rev. Mol. Cell. Biol.* **2016**, *17*, 703–721. [CrossRef] [PubMed]

60. Pommier, Y.; Leo, E.; Zhang, H.; Marchand, C. DNA topoisomerases and their poisoning by anticancer and antibacterial drugs. *Chem. Biol.* **2010**, *17*, 421–433. [[CrossRef](#)]
61. Hevener, K.; Verstak, T.A.; Lutat, K.E.; Riggsbee, D.L.; Mooney, J.W. Recent developments in topoisomerase-targeted cancer chemotherapy. *Acta Pharm. Sin. B* **2018**, *8*, 844–861. [[CrossRef](#)]
62. Xu, Y.; Her, C. Inhibition of Topoisomerase (DNA) I (TOP1): DNA damage repair and anticancer therapy. *Biomolecules* **2015**, *5*, 1652. [[CrossRef](#)]
63. Wang, J.C. Interaction between DNA and an Escherichia coli protein omega. *J. Mol. Biol.* **1971**, *55*, 523–533. [[CrossRef](#)]
64. Madabhushi, R. The roles of DNA topoisomerase II β in transcription. *Int. J. Mol. Sci.* **2018**, *19*, 1917. [[CrossRef](#)] [[PubMed](#)]
65. Sakasai, R.; Iwabuchi, K. The distinctive cellular responses to DNA strand breaks caused by a DNA topoisomerase I poison in conjunction with DNA replication and RNA transcription. *Genes Genet. Syst.* **2016**, *90*, 187–194. [[CrossRef](#)] [[PubMed](#)]
66. Lee, J.H.; Berger, J.M. Cell cycle-dependent control and roles of DNA topoisomerase II. *Genes* **2019**, *10*, 859. [[CrossRef](#)] [[PubMed](#)]
67. Li, M.; Liu, Y. Topoisomerase I in human disease pathogenesis and treatments. *Genom. Proteom. Bioinform.* **2016**, *14*, 166–171. [[CrossRef](#)] [[PubMed](#)]
68. Larsen, A.K.; Escargueil, A.E.; Skladanowski, A. Catalytic topoisomerase II inhibitors in cancer therapy. *Pharm. Ther.* **2003**, *99*, 167–181. [[CrossRef](#)]
69. Hu, W.; Huang, X.S.; Wu, J.F.; Yang, L.; Zheng, Y.T.; Shen, Y.M.; Li, Z.Y.; Li, X. Discovery of novel topoisomerase II inhibitors by medicinal chemistry approaches. *J. Med. Chem.* **2018**, *61*, 8947–8980. [[CrossRef](#)] [[PubMed](#)]
70. Castelli, S.; Goncalves, M.B.; Katkar, P.; Stuchi, G.C.; Couto, R.A.A.; Petrilli, H.M.; da Costa Ferreira, A.M. Comparative studies of oxindolimine-metal complexes as inhibitors of human DNA topoisomerase IB. *J. Inorg. Biochem.* **2018**, *186*, 85–94. [[CrossRef](#)]
71. Katkar, P.; Coletta, A.; Castelli, S.; Sabino, G.L.; Alves Couto, R.A.; da Costa Ferreira, A.M.; Desideri, A. Effect of oxindolimine copper(II) and zinc(II) complexes on human topoisomerase I activity. *Metalomics* **2014**, *6*, 117–125. [[CrossRef](#)]
72. Cerchiaro, G.; Aquilano, K.; Filomeni, G.; Rotilio, G.; Ciriolo, M.R.; Ferreira, A.M. Isatin-Schiff base copper(II) complexes and their influence on cellular viability. *J. Inorg. Biochem.* **2005**, *99*, 1433–1440. [[CrossRef](#)]
73. Filomeni, G.; Cerchiaro, G.; Da Costa Ferreira, A.M.; De Martino, A.; Pedersen, J.Z.; Rotilio, G.; Ciriolo, M.R. Pro-apoptotic activity of novel Isatin-Schiff base copper(II) complexes depends on oxidative stress induction and organelle-selective damage. *J. Biol. Chem.* **2007**, *282*, 12010–12021. [[CrossRef](#)]
74. Chew, S.T.; Lo, K.M.; Lee, S.K.; Heng, M.P.; Teoh, W.Y.; Sim, K.S.; Tan, K.W. Copper complexes with phosphonium containing hydrazone ligand: Topoisomerase inhibition and cytotoxicity study. *Eur. J. Med. Chem.* **2014**, *76*, 397–407. [[CrossRef](#)] [[PubMed](#)]
75. Chen, Z.F.; Tan, M.X.; Liu, L.M.; Liu, Y.C.; Wang, H.S.; Yang, B.; Peng, Y.; Liu, H.G.; Liang, H.; Orvig, C. Cytotoxicity of the traditional chinese medicine (tcm) plumbagin in its copper chemistry. *Dalton Trans.* **2009**, *48*, 10824–10833. [[CrossRef](#)] [[PubMed](#)]
76. Seng, H.L.; Wang, W.S.; Kong, S.M.; Alan Ong, H.K.; Win, Y.F.; Raja Abd Rahman, R.N.Z.; Chikira, M.; Leong, W.K.; Ahmad, M.; Khoo, A.S.B.; et al. Biological and cytoselective anticancer properties of copper(II)-polypyridyl complexes modulated by auxiliary methylated glycine ligand. *BioMetals* **2012**, *25*, 1061–1081. [[CrossRef](#)] [[PubMed](#)]
77. Ikotun, O.F.; Higbee, E.M.; Ouellette, W.; Doyle, R.P. Pyrophosphate-bridged complexes with picomolar toxicity. *J. Inorg. Biochem.* **2009**, *103*, 1254–1264. [[CrossRef](#)]
78. Tabassum, S.; Afzal, M.; Arjmand, F. Synthesis of heterobimetallic complexes: In vitro DNA binding, cleavage and antimicrobial studies. *J. Photochem. Photobiol. B Biol.* **2012**, *114*, 108–118. [[CrossRef](#)]
79. Chauhan, M.; Banerjee, K.; Arjmand, F. DNA binding studies of novel copper(II) complexes containing l-tryptophan as chiral auxiliary: in vitro antitumor activity of cu–sn2 complex in human neuroblastoma cells. *Inorg. Chem.* **2007**, *46*, 3072–3082. [[CrossRef](#)]
80. Afzal, M.; Al-Lohedan, H.A.; Usman, M.; Tabassum, S. Carbohydrate-based heteronuclear complexes as topoisomerase I α inhibitor: Approach toward anticancer chemotherapeutics. *J. Biomol. Struct. Dyn.* **2019**, *37*, 1494–1510. [[CrossRef](#)]

81. Tabassum, S.; Ahmad, A.; Khan, R.A.; Hussain, Z.; Srivastav, S.; Srikrishna, S.; Arjmand, F. Chiral heterobimetallic complexes targeting human DNA-topoisomerase I α . *Dalton Trans.* **2013**, *42*, 16749–16761. [[CrossRef](#)]
82. Tabassum, S.; Asim, A.; Khan, R.A.; Arjmand, F.; Rajakumar, D.; Balaji, P.; Akbarsha, A.M. A multifunctional molecular entity CuII–SnIV heterobimetallic complex as a potential cancer chemotherapeutic agent: DNA binding/cleavage, SOD mimetic, topoisomerase Ia inhibitory and in vitro cytotoxic activities. *RSC Adv.* **2015**, *5*, 47439–47450. [[CrossRef](#)]
83. Lee, S.K.; Tan, K.W.; Ng, S.W. Zinc, copper and nickel derivatives of 2-[2-bromoethyliminomethyl]phenol as topoisomerase inhibitors exhibiting anti-proliferative and antimetastatic properties. *RSC Adv.* **2014**, *4*, 60280–60292. [[CrossRef](#)]
84. Lee, S.K.; Tan, K.W.; Ng, S.W. Topoisomerase I inhibition and DNA cleavage by zinc, copper, and nickel derivatives of 2-[2-bromoethyliminomethyl]-4-[ethoxymethyl]phenol complexes exhibiting anti-proliferation and anti-metastasis activity. *J. Inorg. Biochem.* **2016**, *159*, 14–21. [[CrossRef](#)] [[PubMed](#)]
85. Vutey, V.; Castelli, S.; D’Annessa, I.; Sâmia, L.B.; Souza-Fagundes, E.M.; Beraldo, H.; Desideri, A. Human topoisomerase IB is a target of a thiosemicarbazone copper(II) complex. *Arch. Biochem. Biophys.* **2016**, *606*, 34–40. [[CrossRef](#)] [[PubMed](#)]
86. Haleel, A.K.; Mahendiran, D.; Rafi, U.M.; Veena, V.; Shobana, S.; Rahiman, A.K. Tetrazolo[1,5-a]pyrimidine-based metal(II) complexes as therapeutic agents: DNA interaction, targeting topoisomerase I and cyclin-dependent kinase studies. *Inorg. Nano Met. Chem.* **2019**, *48*, 569–582. [[CrossRef](#)]
87. Haleel, A.K.; Mahendiran, D.; Veena, V.; Sakthivel, N.; Rahiman, A.K. Antioxidant, DNA interaction, VEGFR2 kinase, topoisomerase I and in vitro cytotoxic activities of heteroleptic copper(II) complexes of tetrazolo[1,5-a]pyrimidines and diimines. *Mater. Sci. Eng. C Mater. Biol. Appl.* **2016**, *68*, 366–382. [[CrossRef](#)] [[PubMed](#)]
88. Tabassum, S.; Al-Asbahy, W.M.; Afzal, M.; Arjmand, F.; Bagchi, V. Molecular drug design, synthesis and structure elucidation of a new specific target peptide based metallo drug for cancer chemotherapy as topoisomerase I inhibitor. *Dalton Trans.* **2012**, *41*, 4955–4964. [[CrossRef](#)] [[PubMed](#)]
89. Hasinoff, B.B.; Wu, X.; Yadav, A.A.; Patel, D.; Zhang, H.; Wang, D.S.; Chen, Z.S.; Yalowich, J.C. Cellular mechanisms of the cytotoxicity of the anticancer drug elesclomol and its complex with Cu(II). *Biochem. Pharm.* **2015**, *93*, 266–276. [[CrossRef](#)] [[PubMed](#)]
90. Foo, J.B.; Ng, L.S.; Lim, J.H.; Tan, P.X.; Lor, Y.Z.; Loo, J.S.; Low, M.L.; Chan, L.C.; Beh, C.Y.; Leong, S.W.; et al. Induction of cell cycle arrest and apoptosis by copper complex Cu(SBCM)2 towards oestrogen-receptor positive MCF-7 breast cancer cells. *RSC Adv.* **2019**, *9*, 18359–18370. [[CrossRef](#)]
91. Foo, J.B.; Low, M.L.L.; Lim, J.H.; Lor, Y.Z.; Abidin, R.Z.; Dam, V.E.; Rahman, N.A.; Beh, C.Y.; Chan, L.C.; How, C.W.; et al. Copper complex derived from S-benzyldithiocarbazate and 3-acetylcoumarin induced apoptosis in breast cancer cell. *Biometals* **2018**, *4*, 505–515. [[CrossRef](#)]
92. Yeh, E.T.H.; Ewer, M.; Moslehi, J.; Dlugosz-Danecka, M.; Banchs, J.; Chang, H.M.; Minotti, G. Mechanisms and clinical course of cardiovascular toxicity of cancer treatment I. Oncology. *Semin. Oncol.* **2019**, *46*, 397–402. [[CrossRef](#)]
93. Pendleton, M.; Lindsey, R.H., Jr.; Felix, C.A.; Grimwade, D.; Osheroff, N. Topoisomerase II and leukemia. *Ann. N. Y. Acad. Sci.* **2014**, *1310*, 98–110. [[CrossRef](#)]
94. West, D.X.; Thientanavanich, I.; Liberta, A.E. Copper(II) complexes of 6-methyl-2-acetylpyridine N(4)-substituted thiosemicarbazones. *Trans. Met. Chem.* **1995**, *20*, 303–308. [[CrossRef](#)]
95. Miller, M.C.; Bastow, K.F.; Stineman, C.N.; Vance, J.R.; Song, S.C.; West, D.X.; Hall, I.H. The Cytotoxicity of 2-Formyl and 2-Acetyl-(6-picoly)-4 N-Substituted Thiosemicarbazones and Their Copper(II) Complexes. *Arch. Pharm. Pharm. Med. Chem.* **1998**, *331*, 121–127. [[CrossRef](#)]
96. Khan, T.; Rahmad, R.; Joshi, S.; Khan, A.R. Anticancer potential of metal thiosemicarbazone complexes: A review. *Chem. Sin.* **2015**, *6*, 1–11.
97. Huang, H.; Chen, Q.; Xin, K.; Meng, L.; Lin, L.; Wang, X.; Zhu, C.; Wang, Y.; Chen, Z.; Li, M.; et al. A series of α -heterocyclic carboxaldehyde thiosemicarbazones inhibit topoisomerase II α catalytic activity. *J. Med. Chem.* **2010**, *53*, 3048–3064. [[CrossRef](#)] [[PubMed](#)]
98. Conner, J.D.; Medawala, W.; Stephens, M.T.; Morris, W.H.; Deweese, J.E.; Kent, P.L.; Rice, J.J.; Jiang, X.; Lisic, E.C. Cu(II) benzoylpyridine thiosemicarbazone complexes: Inhibition of human topoisomerase II α and activity against breast cancer cells. *Open J. Inorg. Chem.* **2016**, *6*, 146–154. [[CrossRef](#)]

99. Wilson, J.T.; Jiang, X.; McGill, B.C.; Lisic, E.C.; Deweese, J.E. Examination of the impact of copper(II) α -(n)-heterocyclic thiosemicarbazone complexes on dna topoisomerase II α . *Chem. Res. Toxicol.* **2016**, *29*, 649–658. [[CrossRef](#)]
100. Keck, J.M.; Conner, J.D.; Wilson, J.T.; Jiang, X.; Lisic, E.C.; Deweese, J.E. Clarifying the mechanism of copper(II) α -(N)-heterocyclic thiosemicarbazone complexes on DNA topoisomerase II α and II β . *Chem. Res. Toxicol.* **2019**, *32*, 2135–2143. [[CrossRef](#)]
101. Miller, M.C.; Stineman, C.N.; Vance, J.R.; West, D.X. Multiple Mechanisms for Cytotoxicity Induced by Copper(II) Complexes of 2-Acetylpyrazine-N-substituted Thiosemicarbazones. *Appl. Organometal. Chem.* **1999**, *13*, 9–19. [[CrossRef](#)]
102. Zeglis, B.M.; Divilov, V.; Lewis, J.S. Role of metalation in the topoisomerase II α inhibition and antiproliferation activity of a series of α -heterocyclic-N4-substituted thiosemicarbazones and their Cu(II) complexes. *J. Med. Chem.* **2011**, *54*, 2391–2398. [[CrossRef](#)]
103. Lisic, E.C.; Rand, V.G.; Ngo, L.; Kent, P.; Rice, J.; Gerlach, D.; Papish, E.T.; Jiang, X. Cu(II) propionyl-thiazole thiosemicarbazone complexes: Crystal structure, inhibition of human topoisomerase II α , and activity against breast cancer cells. *Open J. Med. Chem.* **2018**, *8*, 30–46. [[CrossRef](#)]
104. Morris, W.H.; Ngo, L.; Wilson, J.T.; Medawala, W.; Brown, A.R.; Conner, J.D.; Fabunmi, F.; Cashman, D.J.; Lisic, E.; Yu, T.; et al. Structural and metal ion effects on human topoisomerase II α inhibition by α -(N)-heterocyclic thiosemicarbazones. *Chem. Res. Toxicol.* **2019**, *32*, 90–99. [[CrossRef](#)] [[PubMed](#)]
105. Sandhaus, S.; Taylor, R.; Edwards, T.; Huddleston, A.; Wooten, Y.; Venkatraman, R.; Weber, R.T.; González-Sarriás, A.; Martin, P.M.; Cagle, P.; et al. A novel copper(II) complex identified as a potent drug against colorectal and breast cancer cells and as a poison inhibitor for human topoisomerase II α . *Inorg. Chem. Commun.* **2016**, *64*, 45–49. [[CrossRef](#)] [[PubMed](#)]
106. Bacher, F.; Enyedy, É.; Nagy, N.V.; Rockenbauer, A.; Bognár, G.M.; Trondl, R.; Novak, M.S.; Klapproth, E.; Kiss, T.; Arion, V.B. Copper(II) complexes with highly water-soluble L- and D-proline-thiosemicarbazone conjugates as potential inhibitors of Topoisomerase II α . *Inorg. Chem.* **2013**, *52*, 8895–8908. [[CrossRef](#)] [[PubMed](#)]
107. Bisceglie, F.; Musiari, A.; Pinelli, S.; Alinovi, R.; Menozzi, I.; Polverini, E.; Tarasconi, P.; Tavone, M.; Pelosi, G. Quinoline-2-carboxaldehyde thiosemicarbazones and their Cu(II) and Ni(II) complexes as topoisomerase IIa inhibitors. *J. Inorg. Biochem.* **2015**, *152*, 10–19. [[CrossRef](#)]
108. Chen, J.; Huang, Y.W.; Liu, G.; Afrasiabi, Z.; Sinn, E.; Padhye, S.; Ma, Y. The cytotoxicity and mechanisms of 1,2-naphthoquinone thiosemicarbazone and its metal derivatives against MCF-7 human breast cancer cells. *Toxicol. Appl. Pharm.* **2004**, *197*, 40–48. [[CrossRef](#)]
109. Palanimuthu, D.; Shinde, S.V.; Somasundaram, K.; Samuelson, A.G. In vitro and in vivo anticancer activity of copper bis(thiosemicarbazone) complexes. *J. Med. Chem.* **2013**, *56*, 722–734. [[CrossRef](#)]
110. Nair, R.S.; Potti, M.E.; Thankappan, R.; Chandrika, S.K.; Kurup, M.R.; Srinivas, P. Molecular trail for the anticancer behavior of a novel copper carbohydrazone complex in BRCA1 mutated breast cancer. *Mol. Carcinog.* **2017**, *56*, 1501–1514. [[CrossRef](#)]
111. Arjmand, F.; Jamsheera, A.; Afzal, M.; Tabassum, S. Enantiomeric specificity of biologically significant Cu(II) and Zn(II) chromone complexes towards DNA. *Chirality* **2012**, *24*, 977–986. [[CrossRef](#)]
112. Duff, B.; Thangella, V.R.; Creaven, B.S.; Walsh, M.; Egan, D.A. Anti-cancer activity and mutagenic potential of novel copper(II) quinolinone Schiff base complexes in hepatocarcinoma cells. *Eur. J. Pharm.* **2012**, *689*, 45–55. [[CrossRef](#)]
113. Khan, R.A.; Usman, M.; Dhivya, R.; Balaji, P.; Alsalme, A.; AlLohedan, H.; Arjmand, F.; AlFarhan, K.; Akbarsha, M.A.; Marchetti, F.; et al. Heteroleptic copper(I) complexes of “scorpionate” bis-pyrazolyl carboxylate ligand with auxiliary phosphine as potential anticancer agents: An insight into cytotoxic mode. *Sci. Rep.* **2017**, *7*, 45229–45246. [[CrossRef](#)]
114. Ross, W.; Rowe, T.; Glisson, B.; Yalowich, J.; Liu, L. Role of topoisomerase II in mediating epipodophyllotoxin-induced DNA cleavage. *Cancer Res.* **1984**, *44*, 5857–5860. [[PubMed](#)]
115. Furuta, T.; Takemura, H.; Liao, Z.Y.; Aune, G.J.; Redon, C.; Sedelnikova, O.A.; Pilch, D.R.; Rogakou, E.P.; Celeste, A.; Chen, H.T.; et al. Phosphorylation of histone H2AX and activation of Mre11, Rad50, and Nbs1 in response to replication-dependent DNA double-strand breaks induced by mammalian DNA topoisomerase I cleavage complexes. *J. Biol. Chem.* **2003**, *278*, 20303–20312. [[CrossRef](#)] [[PubMed](#)]

116. Bonner, W.M.; Redon, C.E.; Dickey, J.S.; Nakamura, A.J.; Sedelnikova, O.A.; Solier, S.; Pommier, Y. Gamma H2AX and cancer. *Nat. Rev. Cancer* **2008**, *8*, 957–967. [CrossRef] [PubMed]
117. Sordet, O.; Redon, E.C.; Guirouilh-Barbat, J.; Smith, S.; Solier, S.; Douarre, C.; Conti, C.; Nakamura, J.A.; Das, B.B.; Nicolas, E.; et al. Ataxia telangiectasia mutated activation by transcription- and topoisomerase I-induced DNA double-strand breaks. *EMBO Rep.* **2009**, *10*, 887–893. [CrossRef]
118. Pilié, P.G.; Tang, C.; Mills, G.B.; Yap, T.A. State-of-the-art strategies for targeting the DNA damage response in cancer. *Nat. Rev. Clin. Oncol.* **2019**, *16*, 81–104. [CrossRef] [PubMed]
119. Hakem, R. DNA-damage repair; the good, the bad and the ugly. *EMBO J.* **2008**, *27*, 589–605. [CrossRef] [PubMed]
120. Donzelli, M.; Draetta, F.G. Regulating mammalian checkpoints through Cdc25 inactivation. *EMBO Rep.* **2003**, *4*, 671–677. [CrossRef]
121. Banin, S.; Moyal, L.; Shieh, S.; Taya, Y.; Anderson, C.W.; Chessa, L.; Smorodinsky, N.I.; Prives, C.; Reiss, Y.; Shiloh, Y.; et al. Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science* **1998**, *281*, 1674–1677. [CrossRef]
122. Canman, C.E.; Lim, D.S.; Cimprich, K.A.; Taya, Y.; Tamai, K.; Sakaguchi, K.; Appella, E.; Kastan, M.B.; Siliciano, J.D. Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science* **1998**, *281*, 1677–1679. [CrossRef]
123. Hirao, A.; Kong, Y.Y.; Matsuoka, S.; Wakeham, A.; Ruland, J.; Yoshida, H.; Liu, D.; Elledge, S.J.; Mak, T.W. DNA damage-induced activation of p53 by the checkpoint kinase Chk2. *Science* **2000**, *287*, 1824–1827. [CrossRef]
124. Shieh, S.Y.; Ahn, J.; Tamai, K.; Taya, Y.; Prives, C. The human homologs of checkpoint kinases Chk1 and Cds1 (Chk2) phosphorylate p53 at multiple DNA damage-inducible sites. *Genes Dev.* **2000**, *14*, 289–300. [PubMed]
125. Xiong, Y.; Hannon, G.J.; Zhang, H.; Casso, D.; Kobayashi, R.; Beach, D. p21 is a universal inhibitor of cyclin kinases. *Nature* **1993**, *366*, 701–704. [CrossRef] [PubMed]
126. Hengstschläger, M.; Braun, K.; Soucek, T.; Miloloza, A.; Hengstschläger-Ottnad, E. Cyclin-dependent kinases at the G1-S transition of the mammalian cell cycle. *Mutat. Res.* **1999**, *436*, 1–9. [CrossRef]
127. Xiao, D.; Herman-Antosiewicz, A.; Antosiewicz, J.; Xiao, H.; Brisson, M.; Lazo, J.S.; Singh, S.V. Diallyl trisulfide-induced G(2)-M phase cell cycle arrest in human prostate cancer cells is caused by reactive oxygen species-dependent destruction and hyperphosphorylation of Cdc 25 C. *Oncogene* **2005**, *24*, 6256–6268. [CrossRef] [PubMed]
128. Okoh, V.O.; Garba, N.A.; Penney, R.B.; Das, J.; Deoraj, A.; Singh, K.P.; Sarkar, S.; Felty, Q.; Yoo, C.; Jackson, R.M.; et al. Redox signalling to nuclear regulatory proteins by reactive oxygen species contributes to oestrogen-induced growth of breast cancer cells. *Br. J. Cancer* **2015**, *112*, 1687–1702. [CrossRef]
129. Macip, S.; Kosoy, A.; Lee, S.W.; O’Connell, M.J.; Aaronson, S.A. Oxidative stress induces a prolonged but reversible arrest in p53-null cancer cells, involving a Chk1-dependent G2 checkpoint. *Oncogene* **2006**, *25*, 6037–6047. [CrossRef]
130. He, L.; Nan, M.H.; Oh, H.C.; Kim, Y.H.; Jang, J.H.; Erikson, R.L.; Ahn, J.S.; Kim, B.Y. Asperlin induces G₂/M arrest through ROS generation and ATM pathway in human cervical carcinoma cells. *Biochem. Biophys. Res. Commun.* **2011**, *409*, 489–493. [CrossRef]
131. Tubbs, A.; Nussenzweig, A. Endogenous DNA damage as a source of genomic instability in cancer. *Cell* **2017**, *168*, 644–656. [CrossRef]
132. Deiss, K.; Lockwood, N.; Howell, M.; Segeren, H.A.; Saunders, R.E.; Chakravarty, P.; Soliman, T.N.; Martini, S.; Rocha, N.; Semple, R.; et al. A genome-wide RNAi screen identifies the SMC5/6 complex as a non-redundant regulator of a Topo2a-dependent G2 arrest. *Nucleic Acids Res.* **2019**, *47*, 2906–2921. [CrossRef]
133. Bower, J.J.; Zhou, Y.; Zhou, T.; Simpson, D.A.; Arlander, S.J.; Paules, R.S.; Cordeiro-Stone, M.; Kaufmann, W.K. Revised genetic requirements for the decatenation G2 checkpoint: The role of ATM. *Cell Cycle* **2010**, *9*, 1617–1628. [CrossRef]
134. Bower, J.J.; Karaca, G.F.; Zhou, Y.; Simpson, D.A.; Cordeiro-Stone, M.; Kaufmann, W.K. Topoisomerase IIalpha maintains genomic stability through decatenation G(2) checkpoint signaling. *Oncogene* **2010**, *29*, 4787–4799. [CrossRef] [PubMed]
135. Yoshida, C.; Hishiyama, K.; Miyazaki, K.; Watanabe, M.; Kanbe, M.; Yamada, Y.; Matsuzaki, K.; Miyashita, K.; Kitanaka, S.; Miyata, S. Analysis of inhibition of topoisomerase IIalpha and cancer cell proliferation by ingenolEZ. *Cancer Sci.* **2010**, *101*, 374–378. [CrossRef] [PubMed]

136. Dykhuizen, E.C.; Hargreaves, D.C.; Miller, E.L.; Cui, K.; Korshunov, A.; Kool, M.; Pfister, S.; Cho, Y.J.; Zhao, K.; Crabtree, G.R. BAF complexes facilitate decatenation of DNA by topoisomerase II α . *Nature* **2013**, *497*, 624–627. [CrossRef] [PubMed]
137. D Arcy, N.; Gabrielli, B. Topoisomerase II inhibitors and poisons, and the influence of cell cycle checkpoints. *Curr. Med. Chem.* **2017**, *24*, 1504–1519. [CrossRef] [PubMed]
138. Hjaltelin, J.X.; Izarzugaza, J.; Jensen, L.J.; Russo, F.; Westergaard, D.; Brunak, S. Identification of hyper-re wired genomic stress non-oncogene addiction genes across 15 cancer types. *NPJ Syst. Biol. Appl.* **2019**, *5*, 27–37. [CrossRef]
139. Al-Matouq, J.; Holmes, T.R.; Hansen, L.A. CDC25B and CDC25C overexpression in nonmelanoma skin cancer suppresses cell death. *Mol. Carcinog.* **2019**, *58*, 1691–1700. [CrossRef]
140. Butz, H.; Németh, K.; Czenke, D.; Likó, I.; Czirják, S.; Zivkovic, V.; Baghy, K.; Korbonits, M.; Kovácszky, I.; Igaz, P.; et al. Systematic investigation of expression of G2/M transition genes reveals CDC25 alteration in nonfunctioning pituitary adenomas. *Pathol. Oncol. Res.* **2017**, *23*, 633–641. [CrossRef]
141. Mantovani, F.; Collavin, L.; Del Sal, G. Mutant p53 as a guardian of the cancer cell. *Cell Death Differ.* **2019**, *26*, 199–212. [CrossRef]
142. Hayman, L.; Chaudhry, W.R.; Revin, V.V.; Zhelev, N.; Bourdon, J.C. What is the potential of p53 isoforms as a predictive biomarker in the treatment of cancer? *Expert Rev. Mol. Diagn.* **2019**, *19*, 149–159. [CrossRef]
143. Lin, Z.P.; Zhu, Y.L.; Ratner, E.S. Targeting cyclin-dependent kinases for treatment of gynecologic cancers. *Front. Oncol.* **2018**, *8*, 303–314. [CrossRef]
144. Roskoski, R., Jr. Cyclin-dependent protein serine/threonine kinase inhibitors as anticancer drugs. *Pharm. Res.* **2019**, *139*, 471–488. [CrossRef] [PubMed]
145. Kachalaki, S.; Ebrahimi, M.; Mohamed Khosroshahi, L.; Mohammadinejad, S.; Baradaran, B. Cancer chemoresistance: biochemical and molecular aspects: A brief overview. *Eur. J. Pharm. Sci.* **2016**, *89*, 20–30. [CrossRef] [PubMed]
146. Gongora, C.; Vezzio-Vie, N.; Tuduri, S.; Denis, V.; Causse, A.; Auzanneau, C.; Collod-Beroud, G.; Coquelle, A.; Pasero, P.; Pourquier, P.; et al. New Topoisomerase I mutations are associated with resistance to camptothecin. *Mol. Cancer* **2011**, *10*, 64–77. [CrossRef] [PubMed]
147. Tsurutani, J.; Nitta, T.; Hirashima, T.; Komiya, T.; Uejima, H.; Tada, H.; Syunichi, N.; Tohda, A.; Fukuoka, M.; Nakagawa, K. Point mutations in the topoisomerase I gene in patients with non-small cell lung cancer treated with irinotecan. *Lung Cancer* **2002**, *35*, 299–304. [CrossRef]
148. Bassermann, F.; Eichner, R.; Pagano, M. The ubiquitin proteasome system—implications for cell cycle control and the targeted treatment of cancer. *Biochim. Biophys. Acta* **2014**, *1843*, 150–162. [CrossRef] [PubMed]
149. Lee, K.C.; Swan, R.L.; Sondka, Z.; Padgett, K.; Cowell, I.G.; Austin, C.A. Effect of TDP2 on the Level of TOP2-DNA Complexes and SUMOylated TOP2-DNA Complexes. *Int. J. Mol. Sci.* **2018**, *19*, 2056. [CrossRef]
150. Gardner, L.; Malik, R.; Shimizu, Y.; Mullins, N.; ElShamy, W.M. Geminin overexpression prevents the completion of topoisomerase II α chromosome decatenation, leading to aneuploidy in human mammary epithelial cells. *Breast Cancer Res.* **2011**, *13*, R53. [CrossRef]
151. Rozav, A.G.; Chikamori, K.; Kozuki, T.; Grabowski, D.R.; Bukowski, R.M.; Willard, B.; Kinter, M.; Andersen, A.H.; Ganapathi, R.; Ganapathi, M.K. Casein kinase I delta phosphorylates topoisomerase II at serine-1106 and modulates DNA cleavage activity. *Nucleic Acids Res.* **2009**, *37*, 382–392.
152. Kurz, E.U.; Leader, K.B.; Kroll, D.J.; Clark, M.; Gieseler, F. Modulation of human DNA topoisomerase II function by interaction with 14–3–3”. *J. Biol. Chem.* **2000**, *275*, 13948–13954. [CrossRef]
153. Visconti, R.; Della Monica, R.; Grieco, D. Cell cycle checkpoint in cancer: A therapeutically targetable double-edged sword. *J. Exp. Clin. Cancer Res.* **2016**, *35*, 153–161. [CrossRef]
154. Allday, M.J.; Inman, G.J.; Crawford, D.H.; Farrell, P.J. DNA damage in human B cells can induce apoptosis, proceeding from G1/S when p53 is transactivation competent and G2/M when it is transactivation defective. *EMBO J.* **1995**, *14*, 4994–5005. [CrossRef] [PubMed]
155. Vitale, I.; Galluzzi, L.; Castedo, M.; Kroemer, G. Mitotic catastrophe: A mechanism for avoiding genomic instability. *Nat. Rev. Mol. Cell Biol.* **2011**, *12*, 385–392. [CrossRef] [PubMed]
156. Yazinski, S.A.; Comaills, V.; Buisson, R.; Genois, M.M.; Nguyen, H.D.; Ho, C.K.; Todorova Kwan, T.; Morris, R.; Lauffer, S.; Nussenzweig, A.; et al. ATR inhibition disrupts rewired homologous recombination and fork protection pathways in PARP inhibitor-resistant BRCA-deficient cancer cells. *Genes Dev.* **2017**, *31*, 318–332. [CrossRef] [PubMed]

157. Su, Z.; Yang, Z.; Xu, Y.; Chen, Y.; Yu, Q. Apoptosis, autophagy, necroptosis, and cancer metastasis. *Mol. Cancer* **2015**, *14*, 48–62. [[CrossRef](#)]
158. Singh, R.; Letai, A.; Sarosiek, K. Regulation of apoptosis in health and disease: The balancing act of BCL-2 family proteins. *Nat. Rev. Mol. Cell Biol.* **2019**, *20*, 175–193. [[CrossRef](#)]
159. Pfeffer, C.M.; Singh, A.T.K. Apoptosis: A target for anticancer therapy. *Int. J. Mol. Sci.* **2018**, *19*, 448. [[CrossRef](#)]
160. Lee, S.Y.; Ju, M.K.; Jeon, H.M.; Jeong, E.K.; Lee, Y.J.; Kim, C.H.; Park, H.G.; Han, S.I.; Kang, H.S. Regulation of tumor progression by programmed necrosis. *Oxid. Med. Cell. Longev.* **2018**, *2018*, 3537471–3537499. [[CrossRef](#)]
161. Eguchi, Y.; Shimizu, S.; Tsujimoto, Y. Intracellular ATP levels determine cell death fate by apoptosis or necrosis. *Cancer Res.* **1997**, *57*, 1835–1840.
162. Vakkila, J.; Lotze, M.T. Inflammation and necrosis promote tumour growth. *Nat. Rev. Immunol.* **2004**, *4*, 641–648. [[CrossRef](#)]
163. Kang, R.; Chen, R.; Zhang, Q.; Hou, W.; Wu, S.; Cao, L.; Huang, J.; Yu, Y.; Fan, X.G.; Yan, Z.; et al. MGB1 in health and disease. *Mol. Asp. Med.* **2014**, *40*, 1–116. [[CrossRef](#)]
164. Gomes, L.R.; Menck, C.F.M.; Leandro, G.S. Autophagy roles in the modulation of DNA repair pathways. *Int. J. Mol. Sci.* **2017**, *18*, 2351. [[CrossRef](#)] [[PubMed](#)]
165. Yang, Y.; Li, C.; Fu, Y.; Liu, Y.; Zhang, Y.; Zhang, Y.; Zhou, P.; Yuan, Y.; Zhou, S.; Li, S.; et al. Redox cycling of a copper complex with benzaldehyde nitrogen mustard-2-pyridine carboxylic acid hydrazone contributes to its enhanced antitumor activity, but no change in the mechanism of action occurs after chelation. *Oncol. Rep.* **2016**, *3*, 1636–1644. [[CrossRef](#)] [[PubMed](#)]
166. Chen, C.; Lu, L.; Yan, S.; Yi, H.; Yao, H.; Wu, D.; He, G.; Tao, X.; Deng, X. Autophagy and doxorubicin resistance in cancer. *Anticancer Drugs* **2018**, *29*, 1–9. [[CrossRef](#)] [[PubMed](#)]
167. Delgado, J.L.; Hsieh, C.M.; Chan, N.L.; Hiasa, H. Topoisomerases as anticancer targets. *Biochem. J.* **2018**, *475*, 373–398. [[CrossRef](#)]
168. Dehshahri, A.; Ashrafizadeh, M.; Ghasemipour Afshar, E.; Pardakhty, A.; Mandegary, A.; Mohammadinejad, R.; Sethi, G. Topoisomerase inhibitors: Pharmacology and emerging nanoscale delivery systems. *Pharm. Res.* **2020**, *151*, 104551–104563. [[CrossRef](#)]
169. Fujita, K.; Kubota, Y.; Ishida, H.; Sasaki, Y. Irinotecan, a key chemotherapeutic drug for metastatic colorectal cancer. *World J. Gastroenterol.* **2015**, *21*, 12234–12248. [[CrossRef](#)]
170. Woo, W.; Carey, E.T.; Choi, M. Spotlight on liposomal irinotecan for metastatic pancreatic cancer: Patient selection and perspectives. *Onco Targets Ther.* **2019**, *12*, 1455–1463. [[CrossRef](#)]
171. Kondo, R.; Watanabe, S.; Shoji, S.; Ichikawa, K.; Abe, T.; Baba, J.; Tanaka, J.; Tsukada, H.; Terada, M.; Sato, K.; et al. A Phase II Study of Irinotecan for Patients with Previously Treated Small-Cell Lung Cancer. *Oncology* **2018**, *94*, 223–232. [[CrossRef](#)]
172. Xu, F.; Ren, X.; Chen, Y.; Li, Q.; Li, R.; Chen, Y.; Xia, S. Irinotecan-platinum combination therapy for previously untreated extensive-stage small cell lung cancer patients: A meta-analysis. *BMC Cancer* **2018**, *18*, 808–820. [[CrossRef](#)]
173. Lihua, P.; Chen, X.Y.; Wu, T.X. Topotecan for ovarian cancer. *Cochrane Database Syst. Rev.* **2008**, *2008*, CD005589.
174. Pignata, S.; Pisano, C.; Di Napoli, M.; Cecere, S.C.; Tambaro, R.; Attademo, L. Treatment of recurrent epithelial ovarian cancer. *Cancer* **2019**, *24*, 4609–4615. [[CrossRef](#)] [[PubMed](#)]
175. Rosen, V.M.; Guerra, I.; McCormack, M.; Nogueira-Rodrigues, A.; Sasse, A.; Munk, V.C.; Shang, A. Systematic review and network meta-analysis of bevacizumab plus first-line topotecan-paclitaxel or cisplatin-paclitaxel versus non-bevacizumab-containing therapies in persistent, recurrent, or metastatic cervical cancer. *Int. J. Gyn. Cancer* **2017**, *27*, 1237–1246. [[CrossRef](#)] [[PubMed](#)]
176. Qin, A.; Kalemkerian, G.P. Treatment options for relapsed small-cell lung cancer: What progress have we made? *J. Oncol. Pract.* **2018**, *14*, 369–370. [[CrossRef](#)] [[PubMed](#)]
177. Armenian, S.; Bhatia, S. Predicting and preventing anthracycline-related cardiotoxicity. *Am. Soc. Clin. Oncol. Educ. Book* **2018**, *38*, 3–12. [[CrossRef](#)] [[PubMed](#)]
178. Vu, K.; Ai, W. Update on the treatment of anaplastic large cell lymphoma. *Curr. Hematol. Malig. Rep.* **2018**, *13*, 135–141. [[CrossRef](#)] [[PubMed](#)]
179. Liu, W.; Jiang, Q.; Zhou, Y. Advances of systemic treatment for adult soft-tissue sarcoma. *Chin. Clin. Oncol.* **2018**, *7*, 42–55. [[CrossRef](#)]

180. D'Ambrosio, L.; Touati, N.; Blay, J.Y.; Grignani, G.; Flippot, R.; Czarnecka, A.M.; Piperno-Neumann, S.; Martin-Broto, J.; Sanfilippo, R.; Katz, D.; et al. European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. Doxorubicin plus dacarbazine, doxorubicin plus ifosfamide, or doxorubicin alone as a first-line treatment for advanced leiomyosarcoma: A propensity score matching analysis from the European Organization for Research and Treatment of Cancer Soft Tissue and Bone Sarcoma Group. *Cancer* **2020**, *126*, 2637–2647. [[PubMed](#)]
181. Carvalho, C.; Santos, R.X.; Cardoso, S.; Correia, S.; Oliveira, P.J.; Santos, M.S.; Moreira, P.I. Doxorubicin: The good, the bad and the ugly effect. *Curr. Med. Chem.* **2009**, *16*, 3267–3285. [[CrossRef](#)]
182. Banke, A.; Fosbøl, E.L.; Møller, J.E.; Gislason, G.H.; Andersen, M.; Bernsdorf, M.; Jensen, M.B.; Schou, M.; Ejlerksen, B. Long-term effect of epirubicin on incidence of heart failure in women with breast cancer: Insight from a randomized clinical trial. *Eur. J. Heart Fail.* **2018**, *20*, 1447–1453. [[CrossRef](#)]
183. Werntz, R.P.; Adamic, B.; Steinberg, G.D. Emerging therapies in the management of high-risk non-muscle invasive bladder cancer (HRNMIBC). *World J. Urol.* **2019**, *37*, 2031–2040. [[CrossRef](#)]
184. Ravandi, F.; Assi, R.; Daver, N.; Benton, C.B.; Kadia, T.; Thompson, P.A.; Borthakur, G.; Alvarado, Y.; Jabbour, E.J.; Konopleva, M.; et al. Idarubicin, cytarabine, and nivolumab in patients with newly diagnosed acute myeloid leukaemia or high-risk myelodysplastic syndrome: A single-arm, phase 2 study. *Lancet Haematol.* **2019**, *6*, e480–e488. [[CrossRef](#)]
185. Evison, B.J.; Sleefs, B.E.; Watson, K.G.; Phillips, D.R.; Cutts, S.M. Mitoxantrone, More than Just another topoisomerase II poison. *Med. Res. Rev.* **2016**, *36*, 248–299. [[CrossRef](#)] [[PubMed](#)]
186. Barrenetxea Lekue, C.; Grasso Cicala, S.; Leppä, S.; Stauffer Larsen, T.; Herráez Rodríguez, S.; Alonso Caballero, C.; Jørgensen, J.M.; Toldbod, H.; Leal Martínez, I.; D'Amore, F. Pixantrone beyond monotherapy: A review. *Ann. Hematol.* **2019**, *98*, 2025–2033. [[CrossRef](#)] [[PubMed](#)]
187. Minotti, G.; Han, H.; Cattan, V.; Egorov, A.; Bertoni, F. Pixantrone: Novel mode of action and clinical readouts. *Expert Rev. Hematol.* **2018**, *11*, 587–596. [[CrossRef](#)] [[PubMed](#)]
188. Alsdorf, W.; Seidel, C.; Bokemeyer, C.; Oing, C. Current pharmacotherapy for testicular germ cell cancer. *Expert Opin. Pharm.* **2019**, *20*, 837–850. [[CrossRef](#)] [[PubMed](#)]
189. Bernhardt, E.B.; Jalal, S.I. Small Cell Lung Cancer. *Cancer Treat. Res.* **2016**, *170*, 301–322. [[PubMed](#)]
190. Reveiz, L.; Rueda, J.R.; Cardona, A.F. Chemotherapy for brain metastases from small cell lung cancer. *Cochrane Database Syst. Rev.* **2012**, *6*, CD007464. [[CrossRef](#)]
191. Li, J.; Chen, W.; Zhang, P.; Li, N. Topoisomerase II trapping agent teniposide induces apoptosis and G2/M or S phase arrest of oral squamous cell carcinoma. *World J. Surg. Oncol.* **2006**, *4*, 41–47. [[CrossRef](#)]
192. Joyce, M.J.; Pollock, B.H.; Devidas, M.; Buchanan, G.R.; Camitta, B. Chemotherapy for initial induction failures in childhood acute lymphoblastic leukemia: A Children's Oncology Group Study (POG 8764). *J. Pediatr. Hematol. Oncol.* **2013**, *35*, 32–35. [[CrossRef](#)]
193. Pommier, Y.; Cushman, M.; Doroshow, J.H. Novel clinical indenoisoquinoline topoisomerase I inhibitors: A twist around the camptothecins. *Oncotarget* **2018**, *9*, 37286–37288. [[CrossRef](#)]
194. Bailly, C. Contemporary challenges in the design of topoisomerase II inhibitors for cancer chemotherapy. *Chem. Rev.* **2012**, *112*, 3611–3640. [[CrossRef](#)] [[PubMed](#)]
195. Lovrics, A.; Pape, V.F.S.; Szisz, D.; Kalászi, A.; Heffeter, P.; Magyar, C.; Szakács, G. Identifying new topoisomerase II poison scaffolds by combining publicly available toxicity data and 2D/3D-based virtual screening. *J. Cheminform.* **2019**, *11*, 67–81. [[CrossRef](#)]
196. Ortega, J.A.; Riccardi, L.; Minniti, E.; Borgogno, M.; Arencibia, J.M.; Greco, M.L.; Minarini, A.; Sissi, C.; De Vivo, M. Pharmacophore hybridization to discover novel topoisomerase II poisons with promising antiproliferative activity. *J. Med. Chem.* **2018**, *61*, 1375–1379. [[CrossRef](#)] [[PubMed](#)]
197. Beck, D.E.; Abdelmalak, M.; Lv, W.; Reddy, P.V.; Tender, G.S.; O'Neill, E.; Agama, K.; Marchand, C.; Pommier, Y.; Cushman, M. Discovery of potent indenoisoquinoline topoisomerase I poisons lacking the 3-nitro toxicophore. *J. Med. Chem.* **2015**, *58*, 3997–4015. [[CrossRef](#)] [[PubMed](#)]
198. Nitiss, J.L. Targeting DNA topoisomerase II in cancer chemotherapy. *Nat. Rev. Cancer* **2009**, *9*, 338–350. [[CrossRef](#)]
199. Liu, T.; Karlsgen, M.; Karlberg, A.M.; Redalen, K.R. Hypoxia imaging and theranostic potential of [64Cu][Cu(ATSM)] and ionic Cu(II) salts: A review of current evidence and discussion of the retention mechanisms. *Ejnnmci Res.* **2020**, *10*, 33–47. [[CrossRef](#)]

200. Kantarjian, H. Acute myeloid leukemia—Major progress over four decades and glimpses into the future. *Am. J. Hematol.* **2016**, *91*, 131–145.
201. Bornhäuser, M. Vosaroxin in acute myeloid leukaemia. *Lancet Oncol.* **2015**, *16*, 1000–1001. [CrossRef]
202. Skok, Ž.; Zidar, N.; Kikelj, D.; Ilaš, J. Dual inhibitors of human DNA topoisomerase II and other cancer-related targets. *J. Med. Chem.* **2020**, *63*, 884–904. [CrossRef]
203. Kim, S.O.; Sakchaisri, K.; Thimmegowda, N.R.; Soung, N.K.; Jang, J.H.; Kim, Y.S.; Lee, K.S.; Kwon, Y.T.; Asami, Y.; Ahn, J.S.; et al. STK295900, a dual inhibitor of topoisomerase 1 and 2, induces G(2) arrest in the absence of DNA damage. *PLoS ONE* **2013**, *8*, e53908. [CrossRef]
204. Yi, J.M.; Zhang, X.F.; Huan, X.J.; Song, S.S.; Wang, W.; Tian, Q.T.; Sun, Y.M.; Chen, Y.; Ding, J.; Wang, Y.Q.; et al. Dual targeting of microtubule and topoisomerase II by α -carboline derivative YCH337 for tumor proliferation and growth inhibition. *Oncotarget* **2015**, *6*, 8960–8973. [CrossRef]
205. Seo, Y.H. Dual inhibitors against topoisomerases and histone deacetylases. *J. Cancer Prev.* **2015**, *20*, 85–91. [CrossRef] [PubMed]
206. Ferreira, R.; Schneekloth, J.S., Jr.; Panov, K.I.; Hannan, K.M.; Hannan, R.D. Targeting the RNA polymerase I transcription for cancer therapy comes of age. *Cells* **2020**, *9*, 266. [CrossRef] [PubMed]
207. Li, X.; Chen, Y.; Zhao, J.; Shi, J.; Wang, M.; Qiu, S.; Hu, Y.; Xu, Y.; Cui, Y.; Liu, C.; et al. The specific inhibition of SOD1 selectively promotes apoptosis of cancer cells via regulation of the ROS signaling network. *Oxid. Med. Cell Longev.* **2019**, *2019*, 9706792–9706814. [CrossRef] [PubMed]
208. Huang, P.; Feng, L.; Oldham, E.A.; Keating, M.J.; Plunkett, W. Superoxide dismutase as a target for the selective killing of cancer cells. *Nature* **2000**, *407*, 390–395. [CrossRef]
209. Ceci, C.; Atzori, M.G.; Lacal, P.M.; Graziani, G. Role of VEGFs/VEGFR-1 signaling and its inhibition in modulating tumor invasion: Experimental evidence in different metastatic cancer models. *Int. J. Mol. Sci.* **2020**, *21*, 1388.
210. Roskoski, R., Jr. Properties of FDA-approved small molecule protein kinase inhibitors: A 2020 update. *Pharm. Res.* **2020**, *152*, 104609–104628. [CrossRef]
211. Yang, S.W.; Burgin, A.B., Jr.; Huijzen, B.N.; Robertson, C.A.; Yao, K.C.; Nash, H.A. A eukaryotic enzyme that can disjoin dead-end covalent complexes between DNA and type I topoisomerases. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 11534–11539. [CrossRef]
212. Cortes Ledesma, F.; El Khamisy, S.F.; Zuma, M.C.; Osborn, K.; Caldecott, K.W. A human 5'-tyrosyl DNA phosphodiesterase that repairs topoisomerase-mediated DNA damage. *Nature* **2009**, *461*, 674–678. [CrossRef]
213. Zakharenko, A.; Dyrikheva, N.; Lavrik, O. Dual DNA topoisomerase 1 and tyrosyl-DNA phosphodiesterase 1 inhibition for improved anticancer activity. *Med. Res. Rev.* **2019**, *39*, 1427–1441. [CrossRef]
214. Wu, X.; Xue, X.; Wang, L.; Wang, W.; Han, J.; Sun, X.; Zhang, H.; Liu, Y.; Che, X.; Yang, J.; et al. Suppressing autophagy enhances disulfiram/copper-induced apoptosis in non-small cell lung cancer. *Eur. J. Pharm.* **2018**, *827*, 1–12. [CrossRef]
215. Marinello, J.; Delcuratolo, M.; Capranico, G. Anthracyclines as topoisomerase II poisons: From early studies to new perspectives. *Int. J. Mol. Sci.* **2018**, *19*, 3480. [CrossRef] [PubMed]
216. Brown, J.S.; Sundar, R.; Lopez, J. Combining DNA damaging therapeutics with immunotherapy: More haste, less speed. *Br. J. Cancer* **2018**, *118*, 312–324. [CrossRef] [PubMed]
217. Heinhuis, K.M.; Ros, W.; Kok, M.; Steeghs, N.; Beijnen, J.H.; Schellens, J.H.M. Enhancing antitumor response by combining immune checkpoint inhibitors with chemotherapy in solid tumors. *Ann. Oncol.* **2019**, *30*, 219–235. [CrossRef] [PubMed]
218. Kim, N.; Jinks-Robertson, S. The Top1 paradox: Friend and foe of the eukaryotic genome. *DNA Repair* **2017**, *56*, 33–41. [CrossRef] [PubMed]
219. Strickland, K.C.; Howitt, B.E.; Shukla, S.A.; Rodig, S.; Ritterhouse, L.; Liu, J.F.; Garber, J.E.; Chowdhury, D.; Catherine, J.; Andrea, A.D.D.; et al. Association and prognostic significance of BRCA1/2-mutation status with neoantigen load, number of tumor-infiltrating lymphocytes and expression of PD-1/PD-L1 in high grade serous ovarian cancer. *Oncotarget* **2016**, *7*, 1–12. [CrossRef]
220. Li, A.; Yi, M.; Qin, S.; Song, Y.; Chu, Q.; Wu, K. Activating cGAS-STING pathway for the optimal effect of cancer immunotherapy. *J. Hematol. Oncol.* **2019**, *12*, 35–47. [CrossRef]
221. Zhao, B.; Liu, P.; Fukumoto, T.; Fatkhutdinov, N.; Wu, S.; Lin, J.; Aird, K.M.; Tang, H.Y.; Liu, Q.; Speicher, D.W.; et al. Topoisomerase 1 cleavage complex enables pattern recognition and inflammation during senescence. *Nat. Commun.* **2020**, *11*, 908–919. [CrossRef]

222. Wang, Z.; Chen, J.; Hu, J.; Zhang, H.; Xu, F.; He, W.; Wang, X.; Li, M.; Lu, W.; Zeng, G.; et al. cGAS/STING axis mediates a topoisomerase II inhibitor-induced tumor immunogenicity. *J. Clin. Investig.* **2019**, *130*, 4850–4862. [[CrossRef](#)]
223. Wilkinson, R.D.A.; McCabe, N.; Parkes, E.E.; Barros, E.M.; Johnston, D.I.; Ali, R.M.M.; Lappin, K.; Greenberg, R.A.; Harkin, D.P.; McIntosh, S.A.; et al. Topoisomerase II inhibitors induce cGAS-STING dependent inflammation resulting in cytokine induction and immune checkpoint activation. *bioRxiv* **2019**. [[CrossRef](#)]
224. Srinivas, U.S.; Tan, B.; Vellayappan, B.A.; Jeyasekharan, A.D. ROS and the DNA damage response in cancer. *Redox Biol.* **2019**, *25*, 101084–101093. [[CrossRef](#)]
225. Corces, M.R.; Granja, J.M.; Shams, S.; Louie, B.H.; Seoane, J.A.; Zhou, W.; Silva, T.C.; Groeneveld, C.; Wong, C.K.; Cho, S.W.; et al. The chromatin accessibility landscape of primary human cancers. *Science* **2018**, *362*, eaav1898. [[CrossRef](#)] [[PubMed](#)]
226. Temko, D.; Tomlinson, I.P.M.; Severini, S.; Schuster-Böckler, B.; Graham, T.A. The effects of mutational processes and selection on driver mutations across cancer types. *Nat. Commun.* **2018**, *9*, 1857–1867. [[CrossRef](#)] [[PubMed](#)]
227. Alexandrov, L.B.; Alexandrov, L.B.; Nik-Zainal, S.; Wedge, D.C.; Aparicio, S.A.; Behjati, S.; Biankin, A.V.; Bignell, G.R.; Bolli, N.; Borg, A.; et al. Signatures of mutational processes in human cancer. *Nature* **2013**, *500*, 415–421. [[CrossRef](#)] [[PubMed](#)]
228. Helleday, T. Homologous recombination in cancer development, treatment and development of drug resistance. *Carcinogenesis* **2010**, *31*, 955–960. [[CrossRef](#)]
229. Palla, V.V.; Karaolanis, G.; Katafigiotis, I.; Anastasiou, I.; Patapis, P.; Dimitroulis, D.; Perrea, D. gamma-H2AX: Can it be established as a classical cancer prognostic factor? *Tumour Biol.* **2017**, *39*, 1010428317695931. [[CrossRef](#)]
230. Murai, J.; Thomas, A.; Miettinen, M.; Pommier, Y. Schlafen 11 (SLFN11), a restriction factor for replicative stress induced by DNA-targeting anti-cancer therapies. *Pharmacol. Ther.* **2019**, *201*, 94–102. [[CrossRef](#)]
231. Infante Lara, L.; Fenner, S.; Ratcliffe, S.; Isidro-Llobet, A.; Hann, M.; Bax, B.; Osheroff, N. Coupling the core of the anticancer drug etoposide to an oligonucleotide induces topoisomerase II-mediated cleavage at specific DNA sequences. *Nucleic Acids Res.* **2018**, *46*, 2218–2233. [[CrossRef](#)]
232. Ke, X.; Lin, W.; Li, X.; Wang, H.; Xiao, X.; Guo, Z. Synergistic dual-modified liposome improves targeting and therapeutic efficacy of bone metastasis from breast cancer. *Drug Deliv.* **2017**, *24*, 1680–1689. [[CrossRef](#)]
233. Asakura, T.; Yokoyama, M.; Shiraishi, K.; Aoki, K.; Ohkawa, K. Chemotherapeutic effect of CD147 antibody-labeled micelles encapsulating doxorubicin conjugate targeting cd147-expressing carcinoma cells. *Anticancer Res.* **2018**, *38*, 1311–1316.
234. Shi, J.; Su, Y.; Liu, W.; Chang, J.; Zhang, Z. A nanoliposome-based photoactivatable drug delivery system for enhanced cancer therapy and overcoming treatment resistance. *Int. J. Nanomed.* **2017**, *12*, 8257–8275. [[CrossRef](#)]
235. Goffart, S.; Hangas, A.; Pohjoismäki, J.L.O. Twist and Turn-Topoisomerase Functions in Mitochondrial DNA Maintenance. *Int. J. Mol. Sci.* **2019**, *20*, 2041. [[CrossRef](#)] [[PubMed](#)]
236. Baechler, S.A.; Factor, V.M.; Dalla Rosa, I.; Ravji, A.; Becker, D.; Khiati, S.; Miller Jenkins, L.M.; Lang, M.; Sourbier, C.; Michaels, S.A.; et al. The mitochondrial type IB topoisomerase drives mitochondrial translation and carcinogenesis. *Nat. Commun.* **2019**, *10*, 83–96. [[CrossRef](#)] [[PubMed](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).