



HAL
open science

Efficient but less active monitoring system in individuals with high aggressive predispositions.

Fanny Grisetto, Yvonne Delevoye, Clemence Roger

► To cite this version:

Fanny Grisetto, Yvonne Delevoye, Clemence Roger. Efficient but less active monitoring system in individuals with high aggressive predispositions.. International Journal of Psychophysiology, 2019, International journal of psychophysiology : official journal of the International Organization of Psychophysiology, 146, pp.125-132. 10.1016/j.ijpsycho.2019.10.006 . hal-03134085

HAL Id: hal-03134085

<https://hal.univ-lille.fr/hal-03134085>

Submitted on 15 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Grisetto, Delevoeye-Turrell & Roger.

1 **Title:**

2 Efficient but less active monitoring system in individuals with high aggressive predispositions

3

4 **Authors:** Fanny Grisetto, Yvonne N. Delevoeye-Turrell & Clémence Roger

5 **Affiliations:** Univ. Lille, CNRS, CHU Lille, UMR 9193 - SCALab - Sciences Cognitives et
6 Sciences Affectives, F-59000 Lille, France

7 **Corresponding author:** PhD. Clémence Roger, clemence.roger@univ-lille.fr

8

9 **Abstract:** Aggressive behaviors in pathological and healthy populations have been largely
10 related to poor cognitive control functioning. However, few studies have investigated the
11 influence of aggressive traits (i.e., aggressiveness) on cognitive control. In the current study,
12 we investigated the effects of aggressiveness on cognitive control abilities and particularly, on
13 performance monitoring. Thirty-two participants performed a Simon task while
14 electroencephalography (EEG) and electromyography (EMG) were recorded. Participants
15 were classified as having high and low levels of aggressiveness using the BPAQ
16 questionnaire (Buss & Perry, 1992). EMG recordings were used to reveal three response types
17 by uncovering small incorrect muscular activations in ~15% of correct trials (i.e., partial-
18 errors) that must be distinguished from full-error and pure-correct responses. For these three
19 response types, EEG recordings were used to extract fronto-central negativities indicative of
20 performance monitoring, the error and correct (-related) negativities (ERN/Ne and CRN/Nc).
21 Behavioral results indicated that the high aggressiveness group had a larger congruency effect
22 compared to the low aggressiveness group, but there were no differences in accuracy. EEG
23 results revealed a global reduction in performance-related negativity amplitudes in all the
24 response types in the high aggressiveness group compared to the low aggressiveness group.
25 Interestingly, the distinction between the ERN/Ne and the CRN/Nc components was
26 preserved both in high and low aggressiveness groups. In sum, high aggressive traits do not
27 affect the capacity to self-evaluate erroneous from correct actions but are associated with a

1 decrease in the importance given to one's own performance. The implication of these findings
2 are discussed in relation to pathological aggressiveness.

3

4 **Keywords:** aggressiveness; cognitive control; EEG; ERN/Ne; CRN/Nc; performance
5 monitoring

6

7 **Introduction**

8 Aggressiveness is defined as an individual's predispositions to respond aggressively,
9 to experience negative emotions and to hold hostile thoughts (Buss & Perry, 1992). Previous
10 studies in pathological and healthy individuals have demonstrated that aggressive tendencies
11 are linked to poor executive capacities. In maladapted populations, for example, violent
12 offenders, Hancock, Tapscott, & Hoaken (2010) showed that deficits in executive functioning
13 predicted the frequency and severity of intentional acts in physical aggression. Even in a
14 normal population, higher aggressive traits have been associated with low inhibitory control
15 capacities (Pawliczek et al., 2013; Zajenkowski & Zajenkowska, 2015). Conversely, higher
16 cognitive control abilities predicted less aggressive behaviors in response to provocation
17 (Wilkowski, Robinson, & Troop-Gordon, 2010). These evidences indicate that executive
18 functioning and in particular cognitive control, may be valuable research targets to gain a
19 better understanding of aggressiveness.

20 Cognitive control is a set of executive functions that are orchestrated to adjust
21 behaviors, according to internal goals and environmental constraints (Ridderinkhof,
22 Forstmann, Wylie, Burle, & van den Wildenberg, 2011). Among cognitive control
23 mechanisms, some of them are mobilized proactively and others reactively, according to
24 when a risk of making an error is detected (Braver, Gray, & Burgess, 2007; Braver, 2012).
25 Indeed, the risk of making an error must be monitored efficiently in order to involve the

1 appropriate proactive or reactive adjustment mechanism (Nieuwenhuis, Ridderinkhof, Blom,
2 Band, & Kok, 2001). Monitoring capacities are often investigated using
3 electroencephalography (EEG) in participants performing choice reaction time tasks in which
4 the stimulus-response congruency is manipulated, such as the Flanker or the Simon tasks.
5 Falkenstein, Hohnsbein, Hoormann, & Blanke (1991) and Gehring, Goss, Coles, Meyer, &
6 Donchin (1993) reported a negative fronto-central activity emerging at the time of the
7 response and peaking around 100 ms after the response, but only when participants were
8 making errors. This event-related potential (ERP) has been called the error (-related)
9 negativity (ERN/Ne) and is predominantly considered to reflect the involvement of the error
10 detection mechanisms (e.g., Hajcak, Moser, Yeung, & Simons, 2005; Maier, Scarpazza,
11 Starita, Filogamo, & Ladavas, 2016). Indeed, when the situation emphasizes accuracy over
12 speed (e.g., financial penalties for errors), it is important not to make errors and the ERN/Ne
13 is increased, whereas the ERN/Ne is reduced in situations for which the errors are not
14 meaningful because the instructions emphasize speed over accuracy (e.g., Gehring et al.,
15 1993; Hajcak et al., 2005). Another ERP component linked to the performance monitoring
16 system was also reported by Falkenstein, Hohnsbein, & Hoormann (1995) and Falkenstein et
17 al. (1991). This parameter referred as the error positivity (Pe) is a later centro-parietal
18 component that peaks between 200 and 400 ms after the response. It is only observable during
19 trials for which participants consciously detected their errors (Nieuwenhuis et al., 2001); its
20 magnitude varies according to the degree of error consciousness (Leuthold & Sommer, 1999).
21 Therefore, whereas the ERN/Ne indicates automatic internal performance feedback, the Pe is
22 considered to reflect the conscious detection and evaluation of errors (Falkenstein, Hoormann,
23 Christ, & Hohnsbein, 2000).

24 The error-specificity of the ERN/Ne has been challenged by the findings of Vidal,
25 Hasbroucq, Grapperon, & Bonnet (2000) who observed an ERN/Ne-like in non-error trials

1 thanks to the use of electromyography (EMG) and a methodology to increase the spatial
2 resolution of EEG recordings. More specifically, the EMG was recorded to reveal partial-
3 errors that are engaged erroneous actions that are successfully detected, inhibited and
4 corrected (Eriksen, Coles, Morris, & O'hara, 1985; Hasbroucq, Possamaï, Bonnet, & Vidal,
5 1999). Furthermore, in their work, the low spatial resolution of EEG was improved using the
6 Laplacian transform technique (Babiloni, Cincotti, Carducci, Rossini, & Babiloni, 2001;
7 Burle et al., 2015). Vidal et al. (2000) reported the classical ERN/Ne in errors, but also
8 described an ERN/Ne-like following both partial-errors and correct responses, which differed
9 by their magnitudes. Indeed, the ERN/Ne-like observed in correct responses, also called the
10 correct-related negativity (CRN/Nc), is largely smaller in amplitude than the ERN/Ne
11 observed after a partial-error, which is in turn smaller than the ERN/Ne in full-error
12 responses. Despite these differences in amplitude, the three ERPs have similar temporal
13 dynamics, topographies and are generated by the same cerebral regions: the supplementary
14 motor area and/or the dorsal anterior cingulate cortex (Bonini et al., 2014; Fu et al., 2019;
15 Roger, Bénar, Vidal, Hasbroucq, & Burle, 2010). Therefore, these components are thought to
16 reflect an identical process varying in degrees according to performance (Weinberg,
17 Dieterich, & Riesel, 2015). Even though the debates remain around the question of the
18 functional significance of these brain activities, the observed differences in magnitudes
19 between full-error ERN/Ne, partial-error ERN/Ne and CRN/Nc negativities confirm the
20 capacity of the brain to monitor behavioral motor performances. Additionally, the combined
21 analysis of the performance-related negativities (i.e., ERN/Ne and CRN/Nc) enables the
22 precise distinction between the ability to self-evaluate one's own performance from the
23 strength of the monitoring processes that is mobilized during the task. On the one hand, a
24 reduced ERN/Ne in error trials along with an increased CRN/Nc in correct trials indicates a
25 loss in the ability to self-evaluate the ongoing performance with less differentiation between

1 erroneous and correct actions. This is the case for example in schizophrenic patients, in de
2 novo patients with Parkinson disease and in patients with frontal lesions (Mathalon et al.,
3 2002; Turken & Swick, 2008; Willemsen, Müller, Schwarz, Falkenstein, & Beste, 2009). On
4 the other hand, a reduced ERN/Ne combined to a reduced CRN/Nc indicates a global
5 reduction of the importance given to the evaluation of the motor performance. Such a pattern
6 suggests that the monitoring system is less activated throughout the task, but that the ability to
7 self-evaluate an ongoing performance is preserved. Consequently, investigating the CRN/Nc
8 component appears to be relevant since it enables to decide between several interpretations
9 that could be drawn from the reduction in ERN/Ne. Using the same methodology as Vidal et
10 al. (2000), the current study combined the analysis of the negativities of all response types to
11 highlight the monitoring processes in high and low aggressive individuals.

12 Numerous factors differentially modulate the magnitude of the ERN/Ne and the Pe
13 (for a review, see Overbeek, Nieuwenhuis, & Ridderinkhof, 2005). In particular, a reduction
14 in ERN/Ne is often associated with psychiatric disorders such as borderline personality
15 disorders (de Bruijn et al., 2006) and schizophrenia (Charles et al., 2017; Mathalon et al.,
16 2002). These studies suggest that a reduced ERN/Ne is a marker of psychopathology (Olvet &
17 Hajcak, 2008). However, reduced ERN/Ne are also observed in externalizing populations in a
18 broader sense such as conduct disorders and substance dependences (Hall, Bernat, & Patrick,
19 2007; Pasion & Barbosa, 2019) and in juvenile offenders (Vilà-Balló, Hdez-Lafuente, Rostan,
20 Cunillera, & Rodriguez-Fornells, 2014). Aggressive behaviors, a common factor between
21 psychiatric populations, externalizing behaviors and offenders (e.g., Mancke, Herpertz, &
22 Bertsch, 2015; Zhou et al., 2016), thus seem to be associated with a reduction in ERN/Ne.
23 Fewer studies investigated the effect of aggressiveness on the Pe amplitudes. Moreover, their
24 results are less consistent. Comparing offenders and controls, Brazil et al., (2009) showed
25 reduced Pe amplitudes in the offenders whereas Vilà-Balló et al. (2014) did not find any

1 difference between the two groups. This inconsistency might be explained by the presence of
2 psychopathic traits in the population of the Brazil et al. (2009) study compared to Vilà-Balló
3 et al. (2014). Indeed, Steele, Maurer, Bernat, Calhoun, & Kiehl (2016) found larger Pe
4 amplitudes in offenders scoring high in psychopathic traits compared to those scoring low.
5 Psychopathic traits rather than aggressiveness itself may affect Pe amplitudes in these
6 aggressive populations.

7 In the current study, the main goal was to investigate the effect of aggressiveness on
8 performance monitoring. Considering the inconsistency of the findings relative to the Pe
9 component, we did not set a hypothesis of the effect of aggressiveness on its amplitude.
10 However, based upon previous studies (e.g. Charles et al., 2017; Hall et al., 2007; Vilà-Balló
11 et al., 2014), we expected that the reduction in ERN/Ne in full-errors would be revealed in
12 individuals showing high aggressive traits compared to those with low aggressive traits.
13 Moreover, because our participants were well-adapted non-clinical adults, we hypothesized
14 that the monitoring system in our sample would remain as efficient in distinguishing between
15 erroneous and correct actions unlike what it is found in pathological populations (Mathalon et
16 al., 2002; Turken & Swick, 2008; Willemsen et al., 2009). However, if the reduction in the
17 ERN/Ne in erroneous trials is confirmed, then we should observe a similar reduction in the
18 other ERPs to preserve the distinction between each ERN/Ne in the three response types.
19 Consequently, in addition of the reduction in the ERN/Ne in full-error trials, we should
20 observe a reduction both in the ERN/Ne in partial-error trials, and in the CRN/Nc in pure-
21 correct responses.

22

23 **Material and methods**

24 **1. Participants**

1 Thirty-two right-handed volunteers recruited at the University of Lille participated in
2 the study (18 women, mean age = 22.40 years, range from 19 to 28). Handedness was
3 assessed with the Edinburgh Handedness Inventory (Oldfield, 1971). Exclusion criteria
4 included motor and/or sensory disorders and a current medical treatment. They all gave
5 written informed consent for taking part in this study. The ethics committee of the University
6 of Lille (2015-9-S35) approved the experiment.

7

8 **2. Procedure and task**

9 *2.1 Experimental task*

10 The participant sat in a closed room facing a computer screen. She/he performed a
11 modified version of the Simon task (Simon, 1990). Visual stimuli were created in the shape of
12 a circle and of a square. Each participant was invited to respond as quickly and accurately as
13 possible as a function of the shape of the stimulus. For example, holding a response button in
14 each hand, the participant was required to press with the right hand if the stimulus was a circle
15 and with the left hand if it was a square. Shape-to-response mapping rules were
16 counterbalanced across participants. Importantly, the shapes were displayed on the right or on
17 the left part of the screen. Although this dimension of the stimuli was salient, it was irrelevant
18 for the task. Hence, 50% of the trials were labeled as “congruent” since the expected response
19 was ipsilateral to the position of the stimulus (*e.g.* when a circle requiring a left response was
20 presented on the left side of the display). Inversely, 50% of the trials were labeled as
21 “incongruent” since the expected response was contralateral to the position of the stimulus
22 (*e.g.* when a circle requiring a left response was presented on the right side of the display).

23 Each trial begun with the presentation of a fixation cross at the center of the screen
24 during 300 ms. The stimulus appeared and remained displayed until a response was given or

1 after a 1000 ms time lapse. Then, a black screen was presented during 1000 ms before the
2 start of the next trial.

3 The experiment begun with a training block of 20 trials. During this training, visual
4 feedback appeared for 500 ms after each response providing information about the accuracy
5 of the current trial ("*Bonne réponse*" for a correct response, "*Mauvaise réponse*" for an error,
6 "*Essayez d'aller plus vite*" for responses longer than 1000 ms). Then, the participant
7 performed 10 blocks of 129 trials. A pause of 15 seconds was implemented between each
8 block. The experiment lasted about 30 min.

9

10 2.2 Aggressiveness indices

11 Participants responded to the BPAQ Aggression Questionnaire (Buss & Perry, 1992;
12 french version: Pham, Ducro, & Saloppé, 2011) after the end of the Simon task to avoid
13 potential influences of the questionnaire on behaviors.

14 The BPAQ reveals traits in aggression and contains four subscales. The subscales
15 *Physical Aggression* and *Verbal Aggression*) evaluate the external forms of aggression (i.e.,
16 the tendency to act with the focus to hurt someone). The subscale *Anger* evaluates the
17 affective aspect of aggression and is defined as the physiological arousal associated with the
18 preparation for aggression. This subscale assesses the individual differences in the frequency
19 of experiencing the urge to act and the behavioral reactivity towards angry feelings (Poland,
20 Monks, & Tsermentseli, 2016). Finally, the subscale *Hostility* relates to a more cognitive
21 aspect of aggression and is defined as the tendency to evaluate negatively other people, which
22 is often accompanied by a desire to harm others (Poland et al., 2016). Internal consistency
23 was adequate in our sample for the total BPAQ scores, $\alpha = .86$, 95% CI [.79, .93] as well as
24 for the four subscores: *Anger*, $\alpha = .68$, 95% CI [.51, .85], *Hostility*, $\alpha = .76$, 95% CI [.63, .89],
25 *Physical Aggression*, $\alpha = .86$, 95% CI [.80, .93] and *Verbal Aggression*, $\alpha = .55$, 95% CI [.30,

1 .79]. These internal consistency values correspond to those found by Pham et al. (2011) with
2 Cronbach's α below .70 both for the *Anger* and for the *Verbal Aggression* subscales.

3

4 3. **Data Acquisition and Preprocessing**

5 All electrophysiological data were recorded simultaneously using Ag/AgCl electrodes
6 with the BioSemi© system (BioSemi ActiveTwo electrodes, Amsterdam). EEG signals were
7 collected with 64 electrodes (10-20 system positions) mounted on an elastic cap. The vertical
8 electro-oculogram (EOG) was recorded by means of two external electrodes placed below and
9 above the left eye. The horizontal EOG was recorded by means of two external electrodes
10 placed on the temples. The EOGs measurements were recorded to control for eye movement
11 artefacts. The left and the right electromyographic activities (EMG) were recorded by means
12 of two pairs of electrodes placed on the surface of the skin above the thumb-*flexor pollicis*
13 *brevis* of each hand. These EMG measurements were recorded to detect partial-errors and the
14 onset of all the muscular activities. The sampling rate was set to 1024 Hz.

15 Electrophysiological data were collected during the experimental blocks of the Simon task
16 only.

17

18 4. **Electrophysiological data pre-processing**

19 All the electrophysiological data pre-processing steps were done using BrainVision
20 Analyzer 2.1© software (Brain Products, Munich, Germany).

21 The EMG data were filtered with a 10 Hz high-pass filter. Onsets of EMG activities were
22 manually marked after visual inspection as it remained more precise than automatic
23 algorithms (Staupe, Flachenecker, Daumer, & Wolf, 2001). Experimenters were not aware of
24 the nature of the trial being inspected. Based on the manual markers, all trials were classified
25 as (1) pure-correct trials (i.e., trials with only one muscular burst on the correct side), (2) full-

1 error trials (i.e., trials with only one muscular burst on the incorrect side), and (3) partial-error
2 trials (i.e., trials containing two EMG activations, one on the incorrect side preceding the
3 correct response).

4 The raw EEG data were filtered with a 0.16 Hz high-pass filter only and were
5 referenced offline to the left mastoid. The EOGs were used to perform the ocular corrections
6 on the EEG signals following the statistical method by Gratton, Coles, & Donchin (1983). All
7 other artifacts were manually rejected after visual inspection of individual traces. To improve
8 the spatial and the temporal resolutions of the EEG signals, the Laplacian transform was
9 applied to the monopolar data (Babiloni et al., 2001; Burle et al., 2015). To perform this
10 operation, signals were interpolated with the spherical spline interpolation procedure using 3
11 as the degree of spline and 15 degrees, for the Legendre polynomial (Perrin, Bertrand, &
12 Pernier, 1987; Perrin, Pernier, Bertrand, & Echallier, 1989). Then, the second derivatives in
13 two dimensions of space were computed. Thus, electrical brain activities are expressed in
14 $\mu\text{V}/\text{cm}^2$.

15 **5. EEG data processing**

16 Information related to EMG onsets was superimposed upon the EEG signals. EEG
17 data were segmented with respect to the EMG onsets of pure-correct, full-error, and partial-
18 error EMG bursts. EEG segments ranging from 500 ms before and 500 ms after EMG onsets
19 were baseline corrected (100 ms pre-EMG window). Time courses were averaged as a
20 function of the response type.

21 Previous studies showed that the ERN/Ne is maximal at the FCz electrode (e.g. Bates,
22 Kiehl, Laurens, & Liddle, 2002; Ladouceur et al., 2018; Taylor, Visser, Fueggle, Bellgrove, &
23 Fox, 2018; Weinberg et al., 2016). Therefore, the magnitudes of the central negativities were
24 measured at the FCz electrode. A peak-to-peak method (i.e., baseline-free method,
25 Falkenstein, Hoormann, Christ, & Hohnsbein, 2000; Meckler, Carbonnell, Ramdani,

1 Hasbroucq, & Vidal, 2017; Olvet, Hatchwell, & Hajcak, 2010) was applied in the time
2 window between 50 ms and 250 ms after EMG onsets. However, as a peak-to-peak method to
3 measure Pe amplitudes would have been contaminated by the variability of the ERN/Ne
4 amplitudes, the mean positivity in a window frame between 200 ms and 400 ms after EMG
5 onsets was used as an index of Pe amplitudes.

6

7 **6. Experimental groups and statistical analyses**

8 The aim of this study was to evaluate the relation between cognitive control capacities
9 and aggressiveness. In the present study, ERN/Ne amplitudes in all response types were used
10 as an indicator of the performance monitoring capacities. Partial-error rates were used as an
11 indicator of the efficiency in reactive control. Finally, the post-error slowing and the Gratton
12 effects (Gratton, Coles, & Donchin, 1992; Rabbitt, 1966) were used as indicators of the
13 efficiency in proactive control.

14 The BPAQ median score was used to categorize participants as possessing high/low
15 aggressiveness trait personalities. In our sample, the BPAQ scores ranged from 44 to 101,
16 with 67 as the median score. The median-split method categorized 15 participants as low
17 aggressive (i.e. they scored strictly less than 67 in the BPAQ) and 14 participants as highly
18 aggressive (i.e. they scored strictly more than 67 in the BPAQ). Three participants were
19 excluded from ANOVA analyses because their BPAQ scores were equal to the median score.

20 ANOVAs were performed using the R *aov* function available in the *stats* package (R
21 Core Team, 2018). The behavioral performances (i.e., reaction times – RTs, accuracy,
22 reactive and proactive control indices) were submitted to a one-level ANOVA with
23 Aggressiveness as between-group factor. Performance monitoring indices were submitted to a
24 two-level ANOVA with Aggressiveness (2) as between-group factor and Response-Type (3)
25 as within-group factor. Post hoc Scheffé were applied when required and the effect sizes were

Grisetto, Delevoye-Turrell & Roger.

1 calculated as eta-squared and partial eta-squared (η^2 and η_p^2 , respectively) using the R
2 *etaSquared* function available in the *lsr* package (Navarro, 2015). The alpha level was set to
3 .05 for all analyses.

4

5

1 **Results**

2 The following results present the findings obtained during the Simon task in the total
3 sample of 32 participants. We then report the results obtained in the sub-groups after
4 categorizing participants with high and low aggressiveness traits using the total BPAQ score.

6 **Global analyses**

7 **1. Accuracy and reaction times**

8 Among the exploitable EMG recordings, a total of 72.6 %, 23.2 % and 4.1 % of trials
9 were classified as pure-correct, partial-error and full-error, respectively. As classically found,
10 incongruent RTs (478 ms) were longer than congruent RTs (453 ms), $t(31) = -12.41, p < .001$,
11 Cohen's $d = 2.19$. The RTs in full-error trials were shorter (416 ms) than the RTs measured
12 when correct responses were observed (i.e., combined pure-correct and partial-error trials, 465
13 ms), $t(31) = 10.65, p < .001$, Cohen's $d = 1.88$. The post-error slowing was significant: RTs in
14 correct trials following an error were longer (511 ms) than the RTs in correct trials following
15 a correct response (463 ms), $t(31) = -6.96, p < .001$, Cohen's $d = 1.23$. The Gratton effect was
16 significant: the congruency effect was smaller after incompatible trials (-17 ms) compared to
17 the congruency effect observed after compatible trials (69 ms), $F(1,124) = 26.88, p < .001, \eta_p^2$
18 $= .18$. Overall, the classical effects of the Simon task were replicated.

19

20 **2. Performance monitoring**

21 The ANOVA revealed the classical main effect of Response Type on ERN/Ne
22 amplitudes, $F(2,93) = 23.52, p < .001, \eta^2 = .34$. The post hoc Scheffé test confirmed that
23 ERN/Ne was larger in full-error trials than both in partial-error trials ($p < .001$) and in pure-
24 correct trials ($p < .001$). ERN/Ne amplitudes were smaller in pure-correct trials than in partial-
25 error trials ($p < .001$).

1 **Task performance, cognitive control as a function of aggressiveness**

2 ANOVA revealed no main effects of Aggressiveness on error rates, on RTs in full-
 3 error trials and on RTs in correct trials $F(1,27) = 0.12, p = .737, \eta^2 < .01, F(1,27) = 0.55, p =$
 4 $.466, \eta^2 = .02$ and $F(1,27) = 0.34, p = .565, \eta^2 = .01$, respectively. These results are
 5 presented in Table 1. There was a main effect of Aggressiveness on the Simon effect, $F(1,27)$
 6 $= 7.35, p = .012, \eta^2 = .21$ (cf. Figure 1). The high aggressiveness group had a larger Simon
 7 effect (31.13 ms) than the low aggressiveness group (21.74 ms). When considering the
 8 partial-error rates reflecting reactive control, results revealed no main effects of
 9 Aggressiveness, $F(1,27) = 0.06, p = .816, \eta^2 < .01$. When considering proactive control, the
 10 ANOVA revealed no main effects of Aggressiveness on neither the post-error slowing nor the
 11 Gratton effect, $F(1,27) = 1.45, p = .239, \eta^2 = .05$ and $F(1,27) = 0.38, p = .545, \eta^2 = .01$,
 12 respectively.

Table 1.


Means and Standard Deviations for the Task Performance Indices in the High and the Low Aggressiveness Groups.

13

14 *Note.* Correct trials refer to both pure-correct and partial-error trials. Low and high aggressiveness groups were
 15 created based upon the median BPAQ score. *M* = mean, *SD* = standard deviation, RT = reaction time, η^2 = eta-
 16 squared.

Behavioral indices	Low aggressiveness		High aggressiveness		<i>F</i>	<i>p</i>	η^2
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>			
Full-error rates (%)	3.55	1.52	3.77	1.97	0.12	.737	< .01
Partial-error rates (%)	21.88	10.14	20.95	11.12	0.06	.816	< .01
RTs in full-error trials (ms)	382.51	44.32	369.69	49.11	0.55	.466	.02
RTs in correct trials (ms)	417.87	40.10	408.34	47.98	0.34	.565	.01

17



10 *Figure 1.* Mean Simon effect (ms) in the high and the low aggressiveness groups. Error bars
11 represent standard deviations. *: $p < .05$.

12

13 **Performance monitoring and aggressiveness**

14 The EEG traces are represented in Figure 2 as a function of aggressiveness groups and
15 responses types. The analyses revealed a main effect of Aggressiveness on ERN/Ne
16 amplitudes, $F(1,81) = 10.53$, $p = .002$, $\eta_p^2 = .12$. The ERN/Ne amplitudes were smaller in the
17 high aggressiveness group ($-0.56 \mu\text{V}/\text{cm}^2$) than in the low aggressiveness group (-0.77
18 $\mu\text{V}/\text{cm}^2$). Reductions in ERN/Ne amplitudes in the high aggressiveness group were not
19 modulated by Response Type, $F(1,81) = 0.18$, $p = .833$, $\eta_p^2 < .01$. Concerning the Pe
20 observed in full-error trials, there were no differences in amplitudes between the high and the
21 low aggressiveness groups, $F(1, 27) = 1.00$, $p = .326$, $\eta_p^2 = .04$. Table 2 presents the ANOVA
22 results for the performance-related negativities (i.e., both ERN/Ne and CRN/Nc) and the Pe
23 amplitudes.

24

25

Table 2.

ANOVA Results for Mean Performance-Related Negativities and Pe Amplitudes.

Factors	Performance-related negativities			Pe		
	<i>F</i>	<i>p</i>	η_p^2	<i>F</i>	<i>p</i>	η_p^2
Aggressiveness	10.53	.002**	.12	1.00	.326	.04
Response Type	23.21	<.001***	.36			
Aggressiveness x Response Type	0.18	.832	< .01			

1

2

3

4

5

6

Note. Performance-related negativities refer to ERN/Ne in full-errors and in partial-errors responses and CRN/Nc in pure-correct responses. Bold text refers to significant effects. **: $p < .01$, ***: $p < .001$. η_p^2 : partial eta-squared.

7

8

9

10

11

12

13

14

15


16

17

18

19

20



1 *Figure 2.* The Laplacian transformed performance-related negativities (both ERN/Ne and
2 CRN/Nc) and Pe (FCz electrode) and the corresponding Laplacian topographies measured in
3 the full-error (top panel), in the partial-error (middle panel) and in the pure-correct responses
4 (bottom panel). Continuous lines and dotted lines represent the results obtained in the high
5 and the low aggressiveness groups, respectively. Signals were locked to the EMG onset of the
6 motor responses. The topographical maps (right column) show a top/horizontal view of the
7 scalp (nose up) with the actual distribution of the Laplacian-transformed EEG data, separately
8 for each response type, at the time of the negativity peak maximum.

11 **Discussion**

12 Aggressiveness is often associated with poor executive functioning and reduced
13 performance monitoring in pathological and maladapted populations (e.g., Hancock et al.,
14 2010; Vilà-Balló et al., 2014; Zajenkowski & Zajenkowska, 2015). The aim of the current
15 study was to explore cognitive control capacities, and in particular performance monitoring,
16 in a non-clinical population characterized with low and high aggressiveness personality traits.
17 At the behavioral level, the findings showed that the high aggressiveness group globally
18 performed as well as the low aggressiveness group in terms of accuracy and RTs. However,
19 individuals with high aggressive traits were characterized by a greater congruency effect
20 compared to those with low aggressive traits. At the brain level, the current study confirmed
21 the expected reduction in ERN/Ne amplitudes in full-error trials in the high aggressiveness
22 group compared to the low aggressiveness group. Interestingly, the results extended this
23 finding by also revealing a reduction in performance-related negativities after both partial-
24 errors and pure-correct responses. Additionally, the global reduction that was observed in all
25 performance-related negativities amplitudes in the high aggressiveness group did not interact
26 with the response types. Overall, the present results showed a reduction in the global
27 activation of the performance monitoring system, and not a decrease in its efficiency.

1 **Aggressiveness effects on cognitive control at behavioral level**

2 Aggressive behaviors are known to be related to disturbed cognitive control abilities.
3 Indeed, several studies investigating aggressiveness in tasks manipulating the stimulus-
4 response congruency have reported worse performances in pathological and populations with
5 problematic behaviors than in healthy controls (e.g., Gastaldo, Umiltà, Bianchin, & Prior,
6 2002; Hancock et al., 2010). In the current study, the performances of the participants in a
7 choice RT task requiring cognitive control mechanisms were considered through the prism of
8 their predispositions to act aggressively. We selected the BPAQ to assess the levels of
9 aggressiveness through a 35-item questionnaire (Buss & Perry, 1992; Pham et al., 2011). A
10 modified version of the Simon task manipulating the stimulus-response congruency was used
11 to reveal cognitive control functioning. Results showed no differences in the global
12 performance in the Simon task between groups of individuals categorized as having high and
13 low traits in aggressiveness (i.e., RTs and error rates). Additionally, there were no differences
14 neither in partial-error rates nor in behavioral adjustments indicating no effects of aggressive
15 traits, both on reactive and proactive control mechanisms, respectively. However, the high
16 aggressiveness group showed a larger congruency effect than the low aggressiveness group,
17 revealing higher difficulty in inhibiting irrelevant information with higher aggressive
18 tendencies. This result is consistent with a previous study showing longer reaction times in
19 incongruent trials in schizophrenic patients compared to control participants (Gastaldo et al.,
20 2002). While the few differences at the behavioral level between the two groups in our study
21 may seem disconcerting at first sight, they might not be so surprising. Firstly, our participants
22 were young individuals, recruited at the University, without maladapted behaviors. Secondly,
23 the experimental setup was made to not induce aggressive behaviors, like it has been done in
24 some previous studies (e.g., Krämer, Kopyciok, Richter, Rodriguez-Fornells, & Münte, 2011;
25 Pawliczek et al., 2013). Therefore, the traits in aggressiveness were not amplified, and

1 consequently the impact of such a trait may be too weak to be observable within the global
2 indices of behavioral performances (i.e., RTs and error rates). Even if the global performances
3 were not sensitive enough to the trait of aggressiveness in our study, the brain responses
4 revealed a different pattern.

5

6

7

8 **Aggressiveness effects on cognitive control at the brain level**

9 Electroencephalographic recordings were used to assess performance monitoring
10 abilities through the analysis of a specific response-locked ERP that is the error(-related)
11 negativity (ERN/Ne) and its equivalent in pure-correct trials (CRN/Nc). These EEG
12 components are fronto-central negativities peaking rapidly after the response. The amplitude
13 of the ERN/Ne is known to vary as a function of the need to self-evaluate behavioral
14 performances. The more the error is meaningful, the larger is the ERN/Ne amplitude (Gehring
15 et al., 1993; Hajcak et al., 2005). In the current study, we choose to use the Laplacian
16 transform to improve the temporal and spatial resolutions of the EEG (Babiloni et al., 2001;
17 Burle et al., 2015) and especially to uncover the CRN/Nc, usually masked by a large parietal
18 positivity (Roger et al., 2010; Vidal, Burle, Bonnet, Grapperon, & Hasbroucq, 2003; Vidal et
19 al., 2000). We conducted a combined analysis of the negativities in full-errors, in partial-
20 errors and in pure-correct trials. This joint analysis provides the means to specify how a
21 variable can affect patterns of performance monitoring abilities (e.g. task instructions,
22 disorders). In our opinion, only investigating the effect of a variable on the amplitude of the
23 ERN/Ne can mislead interpretations. A reduced ERN/Ne cannot be a marker of error
24 monitoring deficit if the CRN/Nc is also reduced: the monitoring system still distinguishes
25 erroneous and correct actions. This pattern of results should be interpreted as a global

1 decrease in monitoring engagement (i.e., reduction in the value placed on the performance).
2 On the contrary, if a reduced ERN/Ne goes along with a large CRN/Nc, this pattern should be
3 interpreted as a specific difficulty in the evaluation of one's own performance: the monitoring
4 system becomes less discriminant.

5 In the current study, the performance-related negativities peaks were observed within a
6 time window of 100 to 200 ms after the onset of the muscular responses recorded using EMG.
7 Its amplitude was the highest in full-error, intermediate in partial-error and the smallest in
8 pure-correct responses. This pattern of results replicated previously reported findings (e.g.,
9 Meckler et al., 2011; Roger et al., 2010; Roger, Castellar, Pourtois, & Fias, 2014; Vidal et al.,
10 2003, 2000). Considering the aggressiveness effect, the ERN/Ne in full-errors was reported to
11 be decreased in individuals showing high traits in aggressiveness compared to those with low
12 aggressive traits. The current study replicated the findings by Vilà-Balló et al. (2014) in an
13 adapted population and without manipulating the aggressive states of the participants.
14 Interestingly, thanks to the Laplacian transform and the use of EMG, the current study
15 extended these findings by revealing a reduction in the ERN/Ne in partial-errors and in
16 CRN/Nc in pure-correct trials in the high aggressiveness group compared to the low
17 aggressiveness group. Since all the negativities were affected in the same way by
18 aggressiveness, the current findings suggest that aggressiveness does not affect the ability to
19 self-evaluate actions, but instead reduces the importance attached to one's own performance.
20 This global reduction in negativities amplitudes is consistent with several previous studies.
21 Indeed, aggressiveness has already been linked to reduced prefrontal activities in an emotion-
22 manipulated context (Pawliczek et al., 2013) and after exposure to violent video games
23 (Hummer, Kronenberger, Wang, & Mathews, 2019). Nevertheless, the study showed that high
24 aggressive predispositions are characterized by a less active, but still efficient monitoring
25 system even in a neutral situation. It seems that high aggressive individuals care less about

1 their performance than low aggressive individuals do, but are still able to clearly evaluate it.
2 Further studies should consider the functioning of the monitoring system in high aggressive
3 individuals in motivational situations to understand whether the global reduction observed
4 here, in a neutral environment, is due to a genuine inability to mobilize the monitoring
5 system.

6 Aggressive behaviors are characteristic of psychiatric populations (e.g., Mancke et al.,
7 2015; Zhou et al., 2016) and these disorders are also associated with a reduction in the
8 ERN/Ne amplitude (e.g., Charles et al., 2017; de Bruijn et al., 2006). Olvet & Hajcak (2008)
9 even proposed that the ERN/Ne should be considered as an endophenotype of
10 psychopathology. In the current study, we did not carry out precise psychiatric screenings for
11 ethical reasons and, thus, it might be possible that in our sample, especially in the high
12 aggressiveness group, some may meet criteria of psychiatric disorders. Moreover, aggressive
13 predispositions are risk factors for psychiatric disorders (e.g., Mula et al., 2015). Considering
14 the high aggressiveness group as at risk to develop psychiatric disorders, the current results
15 suggest that the global reduction in performance monitoring activities reflects a
16 neurophysiological marker for psychiatric vulnerability. However, this decrease in
17 involvement of the monitoring resources itself is not sufficient to suggest the existence of a
18 psychopathological state. In contrast, a difficulty in distinguishing between erroneous and
19 correct actions may be a more accurate marker of maladapted behaviors. Accordingly to this
20 hypothesis, Hall, Bernat, & Patrick (2007) and Mathalon et al. (2002) both showed weaker
21 differences between ERN/Ne and CRN/Nc amplitudes in externalizing disorders and in
22 schizophrenia, respectively, than in healthy individuals. The interpretation of the reduction in
23 ERN/Ne as a marker of psychopathology (Olvet & Hajcak, 2008) should also take into
24 account the modulation of the CRN/Nc to confirm the presence of disturbed self-evaluation

1 capacities and to specify the true nature of abnormal cognitive control functioning of the
2 monitoring system.

3

4 **Conclusions**

5 This study used the BPAQ questionnaire in order to evaluate aggressive tendencies in
6 a non-clinical and adapted population to compare cognitive control capacities between
7 individuals with low and high aggressive personality traits. At the behavioral level, the high
8 aggressiveness group was associated with a larger congruency effect compared to the low
9 aggressiveness group. However, this difficulty in inhibiting irrelevant information was not
10 reflected in performance: the high and low aggressiveness groups differed neither on error
11 rates nor on reaction times. More particularly, the aim of the current research was to study the
12 influence of these personality traits on the performance monitoring system, which plays a
13 crucial role in cognitive control. Individuals with high aggressive traits showed a reduction in
14 performance-related negativities amplitudes independent of the response type compared to
15 those in the low aggressiveness group. This reduction reveals a decrease in the value placed
16 on performance, but an intact capacity to self-evaluate one's own performance in high
17 aggressiveness. Further studies should be conducted in order to disentangle the influence of
18 aggressive personality traits from the influence of a psychiatric diagnosis on performance
19 monitoring capacities.

20

21 **Acknowledgements**

22 This research was funded in part by the Maison Européenne des Sciences Humaines et
23 Sociales (MESHS; project IMPULS-ACT) and was supported by a grant from the Ministry of
24 Research (program "Investments for the Future") and managed by the National Research
25 Agency under the reference ANR-11-EQPX-0023, as well as by European funds through the

1 program FEDER SCV-IrDIVE. The authors thank the engineers working on the IrDIVE
2 platform for their assistance and Pierre Le Denmat for the proofreading of this paper. This
3 research was supported by a doctoral grant from the University of Lille and the region Hauts-
4 de-France to F.G.

5 The authors declare that there is no conflict of interest regarding the publication of this
6 article.

7

8

9 **References**

- 10 Babiloni, F., Cincotti, F., Carducci, F., Rossini, P. M., & Babiloni, C. (2001). Spatial enhancement of EEG
11 data by surface Laplacian estimation: the use of magnetic resonance imaging-based head
12 models. *Clinical Neurophysiology*, *112*(5), 724-727. [https://doi.org/10.1016/S1388-](https://doi.org/10.1016/S1388-2457(01)00494-1)
13 [2457\(01\)00494-1](https://doi.org/10.1016/S1388-2457(01)00494-1)
- 14 Bates, A. T., Kiehl, K. A., Laurens, K. R., & Liddle, P. F. (2002). Error-related negativity and correct
15 response negativity in schizophrenia. *Clinical Neurophysiology*, *113*(9), 1454-1463.
16 [https://doi.org/10.1016/S1388-2457\(02\)00154-2](https://doi.org/10.1016/S1388-2457(02)00154-2)
- 17 Bonini, F., Burle, B., Liégeois-Chauvel, C., Régis, J., Chauvel, P., & Vidal, F. (2014). Action Monitoring
18 and Medial Frontal Cortex: Leading Role of Supplementary Motor Area. *Science*, *343*(6173),
19 888-891. <https://doi.org/10.1126/science.1247412>
- 20 Braver, T. S., Gray, J. R., & Burgess, G. C. (2007). Explaining the many varieties of working memory
21 variation: dual mechanisms of cognitive control. In A. R. A. Conway, C. Jarrold, & M. J. Kane,
22 *Variation in working memory* (Oxford University Press, p. 76-106). New York.
- 23 Braver, Todd S. (2012). The variable nature of cognitive control: a dual mechanisms framework.
24 *Trends in Cognitive Sciences*, *16*(2), 106-113. <https://doi.org/10.1016/j.tics.2011.12.010>
- 25 Brazil, I. A., de Bruijn, E. R. A., Bulten, B. H., von Borries, A. K. L., van Lankveld, J. J. D. M., Buitelaar, J.
26 K., & Verkes, R. J. (2009). Early and Late Components of Error Monitoring in Violent

- 1 Offenders with Psychopathy. *Biological Psychiatry*, 65(2), 137-143.
2 <https://doi.org/10.1016/j.biopsych.2008.08.011>
- 3 Burle, B., Spieser, L., Roger, C., Casini, L., Hasbroucq, T., & Vidal, F. (2015). Spatial and temporal
4 resolutions of EEG: Is it really black and white? A scalp current density view. *International*
5 *Journal of Psychophysiology*, 97(3), 210-220. <https://doi.org/10.1016/j.ijpsycho.2015.05.004>
- 6 Buss, A. H., & Perry, M. (1992). The Aggression Questionnaire. *Journal of Personality and Social*
7 *Psychology*, 63(3), 452.
- 8 Charles, L., Gaillard, R., Amado, I., Krebs, M.-O., Bendjema, N., & Dehaene, S. (2017). Conscious and
9 unconscious performance monitoring: Evidence from patients with schizophrenia.
10 *NeuroImage*, 144, 153-163. <https://doi.org/10.1016/j.neuroimage.2016.09.056>
- 11 de Bruijn, E. R. A., Grootens, K. P., Verkes, R. J., Buchholz, V., Hummelen, J. W., & Hulstijn, W. (2006).
12 Neural correlates of impulsive responding in borderline personality disorder: ERP evidence
13 for reduced action monitoring. *Journal of Psychiatric Research*, 40(5), 428-437.
14 <https://doi.org/10.1016/j.jpsychires.2005.09.004>
- 15 Eriksen, C. W., Coles, M. G. H., Morris, L. R., & O'hara, W. P. (1985). An electromyographic
16 examination of response competition. *Bulletin of the Psychonomic Society*, 23(3), 165-168.
17 <https://doi.org/10.3758/BF03329816>
- 18 Falkenstein, M., Hohnsbein, J., & Hoormann, J. (1995). Event-related potential correlates of errors in
19 reaction tasks. *Electroencephalography and Clinical Neurophysiology. Supplement*, 44,
20 287-296. [https://doi.org/Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7649035](https://doi.org/Retrieved%20from%20http://www.ncbi.nlm.nih.gov/pubmed/7649035)
- 21 Falkenstein, M., Hohnsbein, J., Hoormann, J., & Blanke, L. (1991). Effects of crossmodal divided
22 attention on late ERP components. II. Error processing in choice reaction tasks.
23 *Electroencephalography and Clinical Neurophysiology*, 78(6), 447-455.
24 [https://doi.org/10.1016/0013-4694\(91\)90062-9](https://doi.org/10.1016/0013-4694(91)90062-9)

- 1 Falkenstein, Michael, Hoormann, J., Christ, S., & Hohnsbein, J. (2000). ERP components on reaction
2 errors and their functional significance: a tutorial. *Biological Psychology*, *51*(2), 87-107.
3 [https://doi.org/10.1016/S0301-0511\(99\)00031-9](https://doi.org/10.1016/S0301-0511(99)00031-9)
- 4 Fu, Z., Wu, D.-A. J., Ross, I., Chung, J. M., Mamelak, A. N., Adolphs, R., & Rutishauser, U. (2019).
5 Single-Neuron Correlates of Error Monitoring and Post-Error Adjustments in Human Medial
6 Frontal Cortex. *Neuron*, *101*(1), 165-177.e5. <https://doi.org/10.1016/j.neuron.2018.11.016>
- 7 Gastaldo, S., Umiltà, C., Bianchin, G., & Prior, M. (2002). The Simon Effect in Schizophrenic Patients
8 with Negative Symptoms. *Cortex*, *38*(2), 149-159. [https://doi.org/10.1016/S0010-](https://doi.org/10.1016/S0010-9452(08)70647-6)
9 [9452\(08\)70647-6](https://doi.org/10.1016/S0010-9452(08)70647-6)
- 10 Gehring, W. J., Goss, B., Coles, M. G. H., Meyer, D. E., & Donchin, E. (1993). A Neural System for Error
11 Detection and Compensation. *Psychological Science*, *4*(6), 385-390.
12 <https://doi.org/10.1111/j.1467-9280.1993.tb00586.x>
- 13 Gratton, G., Coles, M. G. H., & Donchin, E. (1983). A new method for off-line removal of ocular
14 artifact. *Electroencephalography and Clinical Neurophysiology*, *55*(4), 468-484.
15 [https://doi.org/10.1016/0013-4694\(83\)90135-9](https://doi.org/10.1016/0013-4694(83)90135-9)
- 16 Gratton, G., Coles, M. G. H., & Donchin, E. (1992). Optimizing the use of information: Strategic
17 control of activation of responses. *Journal of Experimental Psychology*, *121*(4), 480-506.
- 18 Hajcak, G., Moser, J. S., Yeung, N., & Simons, R. F. (2005). On the ERN and the significance of errors.
19 *Psychophysiology*, *42*(2), 151-160. <https://doi.org/10.1111/j.1469-8986.2005.00270.x>
- 20 Hall, J. R., Bernat, E. M., & Patrick, C. J. (2007). Externalizing Psychopathology and the Error-Related
21 Negativity. *Psychological Science*, *18*(4), 326-333. [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-9280.2007.01899.x)
22 [9280.2007.01899.x](https://doi.org/10.1111/j.1467-9280.2007.01899.x)
- 23 Hancock, M., Tapscott, J. L., & Hoaken, P. N. S. (2010). Role of executive dysfunction in predicting
24 frequency and severity of violence. *Aggressive Behavior*, *36*(5), 338-349.
25 <https://doi.org/10.1002/ab.20353>

- 1 Hasbroucq, T., Possamaï, C.-A., Bonnet, M., & Vidal, F. (1999). Effect of the irrelevant location of the
2 response signal on choice reaction time: an electromyographic study in humans.
3 *Psychophysiology*, 36, 522-526.
- 4 Hummer, T. A., Kronenberger, W. G., Wang, Y., & Mathews, V. P. (2019). Decreased prefrontal
5 activity during a cognitive inhibition task following violent video game play: A multi-week
6 randomized trial. *Psychology of Popular Media Culture*, 8(1), 63-75.
7 <https://doi.org/10.1037/ppm0000141>
- 8 Krämer, U. M., Kopyciok, R. P. J., Richter, S., Rodriguez-Fornells, A., & Münte, T. F. (2011). The Role of
9 Executive Functions in the Control of Aggressive Behavior. *Frontiers in Psychology*, 2.
10 <https://doi.org/10.3389/fpsyg.2011.00152>
- 11 Ladouceur, C. D., Tan, P. Z., Sharma, V., Bylisma, L. M., Silk, J. S., Siegle, G. J., ... Ryan, N. D. (2018).
12 Error-related brain activity in pediatric anxiety disorders remains elevated following
13 individual therapy: a randomized clinical trial. *Journal of Child Psychology and Psychiatry*,
14 59(11), 1152-1161. <https://doi.org/10.1111/jcpp.12900>
- 15 Leuthold, H., & Sommer, W. (1999). ERP correlates of error processing in spatial S-R compatibility
16 tasks. *Clinical Neurophysiology*, 16.
- 17 Maier, M. E., Scarpazza, C., Starita, F., Filogamo, R., & Làdavas, E. (2016). Error monitoring is related
18 to processing internal affective states. *Cognitive, Affective, & Behavioral Neuroscience*, 16(6),
19 1050-1062. <https://doi.org/10.3758/s13415-016-0452-1>
- 20 Mancke, F., Herpertz, S. C., & Bertsch, K. (2015). Aggression in borderline personality disorder: A
21 multidimensional model. *Personality Disorders: Theory, Research, and Treatment*, 6(3),
22 278-291. <https://doi.org/10.1037/per0000098>
- 23 Mathalon, D. H., Fedor, M., Faustmann, W. O., Gray, M., Askari, N., & Ford, J. M. (2002). Response-
24 monitoring dysfunction in schizophrenia: an event-related brain potential study. *Journal of*
25 *Abnormal Psychology*, 111(1), 22-41. <http://dx.doi.org/10.1037/0021-843X.111.1.22>

Grisetto, Delevoeye-Turrell & Roger.

- 1 Meckler, C., Allain, S., Carbonnell, L., Hasbroucq, T., Burle, B., & Vidal, F. (2011). Executive control and
2 response expectancy: A Laplacian ERP study: Executive control and response expectancy.
3 *Psychophysiology*, *48*(3), 303-311. <https://doi.org/10.1111/j.1469-8986.2010.01077.x>
- 4 Meckler, C., Carbonnell, L., Ramdani, C., Hasbroucq, T., & Vidal, F. (2017). On-line action monitoring
5 of response execution: An electrophysiological study. *Biological Psychology*, *129*, 178-185.
6 <https://doi.org/10.1016/j.biopsycho.2017.09.002>
- 7 Mula, M., Agrawal, N., Mustafa, Z., Mohanalingham, K., Cock, H. R., Lozsadi, D. A., & von Oertzen, T.
8 J. (2015). Self-reported aggressiveness during treatment with levetiracetam correlates with
9 depression. *Epilepsy & Behavior*, *45*, 64-67. <https://doi.org/10.1016/j.yebeh.2015.03.018>
- 10 Navarro, D. J. (2015). *Learning statistics with R: a tutorial for psychology students and other*
11 *beginners*. University of Adelaide, Australia.
- 12 Nieuwenhuis, S., Ridderinkhof, K. R., Blom, J., Band, G. P. H., & Kok, A. (2001). Error-related brain
13 potentials are differentially related to awareness of response errors: Evidence from an
14 antisaccade task. *Psychophysiology*, *38*(5), 752-760.
- 15 Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
16 *Neuropsychologia*, *9*(1), 97-113. [https://doi.org/10.1016/0028-3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4)
- 17 Olvet, D. M., & Hajcak, G. (2008). The error-related negativity (ERN) and psychopathology: Toward an
18 endophenotype. *Clinical Psychology Review*, *28*(8), 1343-1354.
19 <https://doi.org/10.1016/j.cpr.2008.07.003>
- 20 Olvet, D. M., Hatchwell, E., & Hajcak, G. (2010). Lack of association between the 5-HTTLPR and the
21 error-related negativity (ERN). *Biological Psychology*, *85*(3), 504-508.
22 <https://doi.org/10.1016/j.biopsycho.2010.09.012>
- 23 Overbeek, T. J. M., Nieuwenhuis, S., & Ridderinkhof, K. R. (2005). Dissociable Components of Error
24 Processing. *Journal of Psychophysiology*, *19*(4), 319-329. [https://doi.org/10.1027/0269-](https://doi.org/10.1027/0269-8803.19.4.319)
25 [8803.19.4.319](https://doi.org/10.1027/0269-8803.19.4.319)

- 1 Pasion, R., & Barbosa, F. (2019). ERN as a transdiagnostic marker of the internalizing-externalizing
2 spectrum: A dissociable meta-analytic effect. *Neuroscience & Biobehavioral Reviews*, *103*,
3 133-149. <https://doi.org/10.1016/j.neubiorev.2019.06.013>
- 4 Pawliczek, C. M., Derntl, B., Kellermann, T., Kohn, N., Gur, R. C., & Habel, U. (2013). Inhibitory control
5 and trait aggression: Neural and behavioral insights using the emotional stop signal task.
6 *NeuroImage*, *79*, 264-274. <https://doi.org/10.1016/j.neuroimage.2013.04.104>
- 7 Perrin, F., Bertrand, O., & Pernier, J. (1987). Scalp Current Density Mapping: Value and Estimation
8 from Potential Data. *IEEE Transactions on Biomedical Engineering*, *BME-34*(4), 283-288.
9 <https://doi.org/10.1109/TBME.1987.326089>
- 10 Perrin, F., Pernier, J., Bertrand, O., & Echallier, J. F. (1989). Spherical splines for scalp potential and
11 current density mapping. *Electroencephalography and Clinical Neurophysiology*, *72*(2),
12 184-187. [https://doi.org/10.1016/0013-4694\(89\)90180-6](https://doi.org/10.1016/0013-4694(89)90180-6)
- 13 Pham, T. H., Ducro, C., & Saloppé, X. (2011). Psychometric assessment of the Buss and Perry
14 Aggression Questionnaire (1992) in a prison population. *Acta Psychiatrica Belgica*, *111*(2), 9.
- 15 Poland, S. E., Monks, C. P., & Tsermentseli, S. (2016). Cool and hot executive function as predictors of
16 aggression in early childhood: Differentiating between the function and form of aggression.
17 *British Journal of Developmental Psychology*, *34*(2), 181-197.
18 <https://doi.org/10.1111/bjdp.12122>
- 19 R Core Team. (2018). *R: A language and environment for statistical computing*. Consulté à l'adresse
20 <https://www.R-project.org/>
- 21 Rabbitt, P. . (1966). Errors and error correction in choice-response tasks. *Journal of Experimental*
22 *Psychology*, *71*(2), 264-272.
- 23 Ridderinkhof, K. R., Forstmann, B., Wylie, S., Burle, B., & van den Wildenberg, W. P. M. (2011).
24 Neurocognitive mechanisms of action control: resisting the call of the Sirens. *Wiley*
25 *Interdisciplinary Reviews: Cognitive Science*, *2*(2), 174-192. <https://doi.org/10.1002/wcs.99>

- 1 Roger, C., Bénar, C. G., Vidal, F., Hasbroucq, T., & Burle, B. (2010). Rostral Cingulate Zone and correct
2 response monitoring: ICA and source localization evidences for the unicity of correct- and
3 error-negativities. *NeuroImage*, *51*(1), 391-403.
4 <https://doi.org/10.1016/j.neuroimage.2010.02.005>
- 5 Roger, C., Castellar, E. ., Pourtois, G., & Fias, W. (2014). Changing your mind before it is too late: The
6 electrophysiological correlates of online error correction during response selection.
7 *Psychophysiology*, *51*(8), 746-760.
- 8 Simon, J. R. (1990). The effects of an irrelevant directional cue on human information processing. In
9 R. W. Proctor & T. G. Reeve (Éd.), *Advances in Psychology* (p. 31-86).
10 [https://doi.org/10.1016/S0166-4115\(08\)61218-2](https://doi.org/10.1016/S0166-4115(08)61218-2)
- 11 Staude, G., Flachenecker, C., Daumer, M., & Wolf, W. (2001). Onset Detection in Surface
12 Electromyographic Signals: A Systematic Comparison of Methods. *EURASIP Journal on*
13 *Advances in Signal Processing*, *2001*(2), 867853.
14 <https://doi.org/10.1155/S1110865701000191>
- 15 Steele, V. R., Maurer, J. M., Bernat, E. M., Calhoun, V. D., & Kiehl, K. A. (2016). Error-related
16 processing in adult males with elevated psychopathic traits. *Personality Disorders: Theory,*
17 *Research, and Treatment*, *7*(1), 80-90. <https://doi.org/10.1037/per0000143>
- 18 Taylor, J. B., Visser, T. A. W., Fueggle, S. N., Bellgrove, M. A., & Fox, A. M. (2018). The error-related
19 negativity (ERN) is an electrophysiological marker of motor impulsiveness on the Barratt
20 Impulsiveness Scale (BIS-11) during adolescence. *Developmental Cognitive Neuroscience*, *30*,
21 77-86.
- 22 Turken, A. U., & Swick, D. (2008). The effect of orbitofrontal lesions on the error-related negativity.
23 *Neuroscience Letters*, *441*(1), 7-10. <https://doi.org/10.1016/j.neulet.2008.05.115>
- 24 Vidal, F., Burle, B., Bonnet, M., Grapperon, J., & Hasbroucq, T. (2003). Error negativity on correct
25 trials: a reexamination of available data. *Biological Psychology*, *64*(3), 265-282.
26 [https://doi.org/10.1016/S0301-0511\(03\)00097-8](https://doi.org/10.1016/S0301-0511(03)00097-8)

- 1 Vidal, F., Hasbroucq, T., Grapperon, J., & Bonnet, M. (2000). Is the 'error negativity' specific to errors?
2 *Biological Psychology*, 51(2), 109-128. [https://doi.org/10.1016/S0301-0511\(99\)00032-0](https://doi.org/10.1016/S0301-0511(99)00032-0)
- 3 Vilà-Balló, A., Hdez-Lafuente, P., Rostan, C., Cunillera, T., & Rodriguez-Fornells, A. (2014).
4 Neurophysiological correlates of error monitoring and inhibitory processing in juvenile
5 violent offenders. *Biological Psychology*, 102, 141-152.
6 <https://doi.org/10.1016/j.biopsycho.2014.07.021>
- 7 Weinberg, A., Dieterich, R., & Riesel, A. (2015). Error-related brain activity in the age of RDoC: A
8 review of the literature. *International Journal of Psychophysiology*, 98(2, Part 2), 276-299.
9 <https://doi.org/10.1016/j.ijpsycho.2015.02.029>
- 10 Weinberg, A., Meyer, A., Hale-Rude, E., Perlman, G., Kotov, R., Klein, D. N., & Hajcak, G. (2016). Error-
11 related negativity (ERN) and sustained threat: Conceptual framework and empirical
12 evaluation in an adolescent sample: ERN and sustained threat. *Psychophysiology*, 53(3),
13 372-385. <https://doi.org/10.1111/psyp.12538>
- 14 Wilkowski, B. M., Robinson, M. D., & Troop-Gordon, W. (2010). How does cognitive control reduce
15 anger and aggression? The role of conflict monitoring and forgiveness processes. *Journal of*
16 *Personality and Social Psychology*, 98(5), 830-840. <https://doi.org/10.1037/a0018962>
- 17 Willemsen, R., Müller, T., Schwarz, M., Falkenstein, M., & Beste, C. (2009). Response Monitoring in
18 De Novo Patients with Parkinson's Disease. *PLoS ONE*, 4(3), e4898.
19 <https://doi.org/10.1371/journal.pone.0004898>
- 20 Zajenkowski, M., & Zajenkowska, A. (2015). Intelligence and aggression: The role of cognitive control
21 and test related stress. *Personality and Individual Differences*, 81, 23-28.
22 <https://doi.org/10.1016/j.paid.2014.12.062>
- 23 Zhou, J.-S., Zhong, B.-L., Xiang, Y.-T., Chen, Q., Cao, X.-L., Correll, C. U., ... Wang, X.-P. (2016).
24 Prevalence of aggression in hospitalized patients with schizophrenia in China: A meta-
25 analysis: Aggression and schizophrenia. *Asia-Pacific Psychiatry*, 8(1), 60-69.
26 <https://doi.org/10.1111/appy.12209>