



**HAL**  
open science

## O-GlcNAcylation: A New Cancer Hallmark?

Yann Fardini, Vanessa Dehennaut, Tony Lefebvre, Tarik Issad

► **To cite this version:**

Yann Fardini, Vanessa Dehennaut, Tony Lefebvre, Tarik Issad. O-GlcNAcylation: A New Cancer Hallmark?. *Frontiers in Endocrinology*, 2013, 4, pp.99. 10.3389/fendo.2013.00099 . hal-03173298

**HAL Id: hal-03173298**

**<https://hal.univ-lille.fr/hal-03173298>**

Submitted on 18 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# O-GlcNAcylation: a new cancer hallmark?

Yann Fardini<sup>1,2</sup>, Vanessa Dehennaut<sup>3</sup>, Tony Lefebvre<sup>3</sup> and Tarik Issad<sup>1,2\*</sup>

<sup>1</sup> Institut Cochin, Université Paris Descartes, CNRS (UMR8104), Paris, France

<sup>2</sup> INSERM, U1016, Paris, France

<sup>3</sup> CNRS/UMR 8576, Unit of Structural and Functional Glycobiology, Institut Fédératif de Recherche IFR 147, Lille 1 University, Villeneuve d'Ascq, France

## Edited by:

Kenneth Siddle, University of Cambridge, UK

## Reviewed by:

Ramasamy Paulmurugan, Stanford University, USA

Zhixiang Wang, University of Alberta, Canada

## \*Correspondence:

Tarik Issad, Department of Endocrinology, Metabolism and Diabetes, Institut Cochin, 22 rue Méchain, 75014 Paris, France  
e-mail: tarik.issad@inserm.fr

O-linked *N*-acetylglucosaminylation (O-GlcNAcylation) is a reversible post-translational modification consisting in the addition of a sugar moiety to serine/threonine residues of cytosolic or nuclear proteins. Catalyzed by O-GlcNAc-transferase (OGT) and removed by O-GlcNAcase, this dynamic modification is dependent on environmental glucose concentration. O-GlcNAcylation regulates the activities of a wide panel of proteins involved in almost all aspects of cell biology. As a nutrient sensor, O-GlcNAcylation can relay the effects of excessive nutritional intake, an important cancer risk factor, on protein activities and cellular functions. Indeed, O-GlcNAcylation has been shown to play a significant role in cancer development through different mechanisms. O-GlcNAcylation and OGT levels are increased in different cancers (breast, prostate, colon...) and vary during cell cycle progression. Modulating their expression or activity can alter cancer cell proliferation and/or invasion. Interestingly, major oncogenic factors have been shown to be directly O-GlcNAcylated (p53, MYC, NFκB, β-catenin...). Furthermore, chromatin dynamics is modulated by O-GlcNAc. DNA methylation enzymes of the Tet family, involved epigenetic alterations associated with cancer, were recently found to interact with and target OGT to multi-molecular chromatin-remodeling complexes. Consistently, histones are subjected to O-GlcNAc modifications which regulate their function. Increasing number of evidences point out the central involvement of O-GlcNAcylation in tumorigenesis, justifying the attention received as a potential new approach for cancer treatment. However, comprehension of the underlying mechanism remains at its beginnings. Future challenge will be to address directly the role of O-GlcNAc-modified residues in oncogenic-related proteins to eventually propose novel strategies to alter cancer development and/or progression.

**Keywords:** O-glycosylation, O-GlcNAc, post-translational modification, cancer, metastasis, cell cycle, epigenetics, transcription factors

## INTRODUCTION

Excess food intake associated with modern lifestyle constitutes an important cancer risk factor (1). Numerous epidemiological studies indicate that obesity or diabetic conditions increase the risk of cancer, including colon, esophageal, liver, pancreas kidney, endometrial, and breast cancers (2, 3). Increased body mass index (BMI) above 25 kg/m<sup>2</sup> is associated with a significantly increased relative risk of several cancers, and BMI higher than 30 kg/m<sup>2</sup> are associated with a two- to four-fold increase in colorectal, endometrial, esophageal, liver, gallbladder, and gastric cancers (4). Hyperglycemia also appears to be an important cancer risk factor. Indeed, in a 10-year prospective study involving 1,298 million Koreans, individuals with fasting serum glucose higher than 140 mg/dl had significant higher death rates from all cancers combined than those with fasting glucose lower than 90 mg/dl. Sustained weight loss, improvement of insulin resistance, and attenuation of metabolic syndrome observed after bariatric surgery are associated with reduction in cancer incidence (5). In mouse and rat models of diet-induced obesity, overfeeding is associated with accelerated development of tumors (6). In contrast, food restriction has inhibitory effects on tumor

growth in rodents (7) and reduces cancer incidence in non-human primates (8). Interestingly, low calorie intake habits in Okinawan population (9) is associated with reduced cancer risk compared to mainland Japan (10), suggesting that the anti-cancer effects of calorie restriction in rodent models may extend to primates, including humans.


Nutritional conditions, excess body weight, and insulin resistance can modulate tumor development by modifying circulating factors that affect signaling pathways involved in cell growth, proliferation, and apoptosis. For instance, chronic hyperinsulinemia is associated with increased risk of colon, pancreas, breast, and endometrium cancers (4). These effects could be directly mediated by insulin receptors on (pre)neoplastic target cells, or might be secondary to hyperinsulinemia. Thus, insulin promotes the synthesis of IGF1, and inhibits the expression of insulin-like growth factor binding proteins 1 and 2 (IGFBP1 and IGFBP2), resulting in increased bioavailability of this potent growth factor. In addition, insulin and IGF1 inhibit the expression of sex-hormone binding globulin (SHBG), resulting in increase in estrogen bioavailability, a breast cancer risk factor in post-menopausal women. Increased leptin circulating

levels associated with excess adiposity may also constitute a risk factor for breast cancer. Indeed, ObRb, the long form of the leptin receptor stimulates proliferative and anti-apoptotic pathways, and both leptin and leptin receptors are overexpressed in human primary and metastatic breast cancer cells (11, 12).

However, besides its effects on circulating hormones and adipokines, nutritional conditions also modulate the availability of nutrients important for growth and proliferation of cancer cells. Among them, glucose and glutamine are considered crucial, and cancer cells have been described as “addicted” to these two nutrients, from which they obtain biosynthetic precursors to build cell membranes, DNA, and proteins. Glucose and glutamine are the two most abundant extracellular nutrients (13). They contribute carbons to the synthesis of the three major classes of macromolecules (nucleic acids, lipids, and proteins) in proliferating tumor cells. In addition to its role as a carbon source, glutamine also donates nitrogen to nucleotide and amino acid synthesis. Thus, biosynthesis of nucleotides utilizes ribose 5-phosphate produced from the diversion of glycolytic intermediate into the pentose phosphate pathway, and glutamine. Fatty acid synthesis, used to produce lipids, requires acetylCoA generated from glucose. Protein synthesis requires amino acids, tRNAs, and ribosomes. Both glucose and glutamine are used to generate these molecules (13). However, in addition to their role as “molecular bricks” in building of cancer cell components, glucose and glutamine metabolism intervene in protein *O*-GlcNAcylation, a post-translational modification that regulates most aspects of cell life (14–18). *O*-GlcNAcylation of cytosolic and nuclear proteins is a reversible post-translational modification, analogous to phosphorylation, which controls protein subcellular localization, stability, or activity according to the nutritional environment. It corresponds to the addition of *N*-acetylglucosamine (GlcNAc) on serine or threonine residues. The reaction is catalyzed by *O*-GlcNAc-transferase (OGT), which uses UDP-GlcNAc as a substrate (Figure 1). UDP-GlcNAc, produced through the hexosamine biosynthetic pathway (HBP), can be considered as a sensor for the nutritional state of the cell, as it integrates glucose, glutamine, fatty acids (acetyl), uridine, and ATP metabolism (14–18). *O*-GlcNAc is removed from proteins by *O*-GlcNAcase (OGA), permitting dynamic regulation of *O*-GlcNAcylation levels. *O*-GlcNAcylation can be in competition directly or indirectly with phosphorylation on the same protein, providing a complex cross-talk between these two PTM to control the function of various proteins (19).

Excess of nutrients intake, hyperglycemia, and other metabolic perturbations associated with diabetes and obesity are believed to feed the HBP and promote abnormally elevated *O*-GlcNAcylation of key signaling molecules and transcription factors. These modifications have been proposed to play key roles in complications associated with the metabolic syndrome, diabetic conditions, neurodegenerative disease, and cancer (17, 20, 21).

On the other hand, *O*-GlcNAcylation regulates cell cycle, signaling intermediates, and transcription factors involved in the control of cell proliferation or cell death (22–27). A growing amount of studies indicates that *O*-GlcNAcylation may constitute an important regulator of cancer growth and invasion, providing a potential link between obesity, diabetes, and cancer (28, 29).



**FIGURE 1 | The hexosamine biosynthetic pathway and protein *O*-GlcNAcylation.** The hexosamine biosynthetic pathway (HBP) controls *O*-GlcNAc glycosylation (*O*-GlcNAcylation) of nuclear and cytosolic proteins. This dynamic and reversible post-translational modification controls activity, localization, or stability of substrate proteins, according to the rate of glucose availability. A fraction (2–3%) of glucose entering the cell is directed to the HBP. In this pathway, fructose-6-phosphate is converted to glucosamine-6-phosphate by the glutamine:fructose-6-phosphate amidotransferase (GFAT), the rate-limiting enzyme of the pathway. After a subset of reactions, UDP-*N*-acetylglucosamine (UDP-GlcNAc) is generated and used by the *O*-GlcNAc-transferase (OGT) as a substrate to add GlcNAc to serine or threonine residues of target proteins. *O*-GlcNAc moiety is removed from *O*-GlcNAc-modified proteins by the *O*-GlcNAcase (OGA). Experimentally, the level of *O*-GlcNAc-modified proteins in cells can be manipulated by exposing cells to high glucose concentrations or to glucosamine which enters the HBP downstream the rate-limiting GFAT-mediated reaction, as glucosamine-6-phosphate. In addition, OGA can be inhibited by pharmacological agents such as *O*-[2-acetamid-*O*-2-deoxy- $\beta$ -glucopyranosylidene] amino-*N*-phenylcarbamate (PUGNAc) or 1,2-dideoxy-2'-propyl- $\alpha$ - $\beta$ -glucopyranoso-[2,1-D]- $\Delta$ 2'-thiazoline (NButGT), resulting in an accumulation of *O*-GlcNAc-modified proteins in the cell.

## *O*-GLcNAcylation and *O*-GLcNAc-CYCLING ENZYMES IN CANCER

Increased protein *O*-GlcNAcylation and changes in OGT and/or OGA expression have now been described in different cancer types including breast (30–32), lung (33), colon (33), liver (34), bladder (35), endometrial (36) prostate (37), and chronic lymphocytic leukemia (CCL) cells (38).

### BREAST CANCER

Breast cancer is the most common cancer in women. The link between nutritional conditions, obesity, and breast cancer is well established (39). Initial studies by Slawson et al. suggested increased OGA activity in primary breast tumors compared to matched adjacent breast tissues (30). However, because the data set was relatively small, no clear correlation could be established with tumor grade or type. Moreover, these results contrasted with those

obtained by other groups in more recent investigations. Dahl et al. defined a set of 40 candidate genes that are predominantly localized in the most frequently altered chromosomal regions known to be important in the pathogenesis of breast and ovarian cancers (40). Systematic characterization of these candidate genes by both cDNA dot plot using cancer profiling array and real-time RT-PCR analysis revealed differential expression in breast cancer for nine genes, including *MGEA5*, the gene coding for OGA, which expression was reduced by about 56% in breast tumors (40). In agreement with decreased OGA in breast cancer, Gu et al. using immunohistochemistry analysis, observed that the global O-GlcNAcylation level in breast tumor tissues was significantly elevated compared to the corresponding adjacent normal tissue (31). Moreover, O-GlcNAcylation was also significantly enhanced in metastatic lymph nodes compared with their corresponding primary breast tissues, suggesting a potential influence of O-GlcNAcylation on malignant properties of the breast cancer cells (31). More recently, Krzeslak et al. observed increased OGT and decreased OGA mRNA expression in breast tumors (32), with poorly differentiated tumors (grade II and III) having significantly higher OGT and lower OGA mRNA expression than grade I tumors, respectively. Lymph node metastasis status was also significantly associated with decreased OGA mRNA expression, but not with OGT mRNA expression. OGT and OGA expression profiles showed no significant differences in association with different estrogen and progesterone status (32). These results have been confirmed very recently by Champattanachai and colleagues who analyzed breast tumors from grade I to III and observed increased total O-GlcNAc protein levels along with OGT levels compared to healthy tissue. This increase was also proportional to tumor grades (41).

Finally, a role for O-GlcNAcylation in tamoxifen sensitivity was recently described (42). In breast cancer, tamoxifen remains largely used as endocrine therapy, but resistance to this drug often occurs (43). Kanwal et al. observed that O-GlcNAcylation-inducing treatment protected MCF-7 cells from tamoxifen-induced cell death, whereas siRNA-mediated OGT inhibition had opposite effects. These data suggest that targeting OGT might be an interesting approach to overcome tamoxifen resistance in breast cancer (42).

### COLORECTAL CANCER

Colorectal cancers (CRCs) are one of the leading causes of mortality and morbidity by cancer, the second for women and the third for men, in Europe and the United States of America. Unhealthy lifestyle and nutrient excess favor emergence of CRCs and, more generally, these cancers are intimately linked to metabolic diseases that themselves reach epidemic proportions in the Western societies. As an example, type-2 diabetes doubles CRC emergence (44). Evaluation of O-GlcNAc in CRC using immunohistochemical analyses revealed that O-GlcNAcylation level and OGT expression are increased in colon tissues in comparison with the corresponding adjacent tissues, whereas no significant difference was observed for OGA (33).

### LIVER CANCER

Although liver cancers (LCs) only represent the fifth and the seventh most common cause of cancer in men and women

respectively, their frequency has been increasing steadily for 20 years. Indeed, as for CRCs, unhealthy lifestyle favors emergence of LCs, and diabetes and obesity increase the risk of LCs through the development of a non-alcoholic steatosis (45).

Zhu et al. (34) analyzed O-GlcNAcylation level and expression of OGT and OGA by real-time PCR in tissues of patients suffering hepatocellular carcinoma, the most common primary liver malignancy (600,000 new cases diagnosed each year). Immunohistochemistry analyses of patients having undergone a liver transplantation, that remains at the present time the best treatment for LC and cirrhosis despite the high incidence of tumor recurrence, indicate that O-GlcNAcylation was higher in cancer tissues compared to healthy tissues. Moreover, O-GlcNAcylation was higher again in patients diagnosed a recurrence of hepatocarcinoma. The authors tried to correlate OGT and OGA expression with the prognosis of patients who underwent liver transplantation. The 60 tumors used were segregated into two groups according to low/high amounts of OGT/OGA transcripts. More than three quarters of the recurrent tumor tissues show low OGA expression while the non-recurrent ones have high OGA levels. Intriguingly, no significant correlation for OGT was found. The recurrence-free survival was significantly better for patients showing high OGA than patients with low OGA, while OGT expression did not allow predicting the prognosis. The authors proposed that low expression of OGA may constitute an additional independent prognostic factor for the recurrence-free survival following liver transplantation.

### BLADDER AND ENDOMETRIAL CANCERS

Krzeslak et al. measured the level of OGA and OGT mRNA in bladder (35) and endometrial (36) cancers. Rozanski et al. observed that OGA mRNA was present in the urine (cells pelleted from centrifuged urines) of healthy and bladder cancer patients with almost the same proportion (47.1% of healthy and 52.3% of bladder cancer patients). OGT mRNA was found in 50% of patients suffering bladder cancer while it was not detected in healthy individuals. OGA mRNA level was higher in grade I tumors compared to grade III, whereas OGT mRNA was lower in grade I than in grades II and III. The authors concluded that measurement of OGT and OGA mRNA in urine might be an interesting parameter for the diagnosis bladder cancers (35).

Regarding the endometrial carcinomas, it appeared that OGT and OGA mRNA were significantly more elevated in grades II and III tumors than in grade I. In addition, OGT and OGA expression was higher in case of cancers with deep myometrial invasion. Although both OGT and OGA mRNA were increased, the relative expression of OGT was much higher than OGA. However, no significant differences in enzymes messengers were found for the different lymph nodes metastasis (36).

### OTHER CANCERS

Changes in O-GlcNAc levels and/or expression of O-GlcNAc-cycling enzymes have also been described in leukemia, lung, and prostate cancers (33, 37, 38).

Increased protein O-GlcNAcylation and OGT expression were observed in CCL cells compared to normal circulating and tonsillar B cells (38). However, higher O-GlcNAc levels in CCL cells

were associated with a favorable outcome, while lower levels were associated with more aggressive disease.

Mi et al. constructed a tissue microarray comprising 31 paraffin-embedded lung cancer tissues and their corresponding adjacent normal lung tissues (33). Using immunohistochemistry, the authors observed increased O-GlcNAcylation level and OGT expression in cancer tissues, whereas OGA expression level was not modified. In agreement with their results, examining Oncomine™ database, Mi et al. found increased OGT mRNA expression in lung cancers (33).

Lynch et al. (37) analyzed Oncomine™ database to determine whether OGT was overexpressed in tumors, and found four microarray gene expression studies showing elevated OGT mRNA levels in human prostate carcinoma as compared to adjacent tissue samples. In addition, a survey of the National Center for Biotechnology Information Gene Expression Omnibus indicated a positive correlation between high OGT expression and metastatic progression between normal, primary tumor, and metastatic tumor tissues. Moreover, an additional study of 94 patient tumor samples, which when stratified by level of OGT expression, indicated that disease-free survival 5 years post-treatment was higher in patients with low OGT expression profile compared to patients with increased OGT expression. In agreement with these data, using a panel of normal and prostate carcinoma cell lines, Lynch et al. observed an increased O-GlcNAc level and OGT expression associated with malignant properties in prostate cancer cells (37).

In summary, studies on different types of cancer generally indicate increased expression in O-GlcNAc level and OGT expression. However, contradictory results were obtained concerning OGA, with either decreased (31, 32, 34, 35, 40), no change (33), and even increased in OGA levels in some studies (30, 36, 46). These discrepancies in OGA results may originate from the fact that OGA expression itself is upregulated upon increase O-GlcNAcylation level in the cell (47), complicating the interpretation of the data.

From the literature reviewed in the previous sections, it appears that cancer cells display both increased uptake of nutrients involved in O-GlcNAc biosynthesis and increased capacity to O-GlcNAcylate proteins. As we shall see in the following sections, alteration in O-GlcNAcylation may directly affect important steps in tumorigenesis.

## O-GLcNACYLATION AND MITOGENIC SIGNALING PATHWAYS

As mentioned previously, the nutritional status impacts tumor development. Thus, several studies in rodent showed that dietary restriction can inhibit tumor growth. Mitogenic signals elicited by various receptors often involve common signaling cascades such as PI3K/Akt or MAPK pathways. Interestingly, when grown as tumor xenografts in mice, cancer cells bearing mutations that induce constitutive PI3K activation are resistant to dietary restriction, whereas cancer cells with mutations that constitutively activate the Ras/Raf/MAPK pathway remain sensitive to calorie restriction (48, 49). These observations suggest that the activity of the PI3K/Akt pathway is central to sensitivity of cancer cells to nutritional conditions. Interestingly, O-GlcNAcylation has been implicated in modulation of PI3K/Akt activity. For instance, in thyroid anaplastic cancer 8305C cells, increased O-GlcNAcylation

induced by OGA inhibition resulted in an increase in basal and IGF1 stimulated Akt activity (32). In MCF-7 breast cancer cells, Olivier-Van Stichelen et al. showed that serum-induced Akt activation was markedly impaired by siRNA-mediated OGT inhibition (27). Moreover, Kanwal et al. showed that treating MCF-7 cells with O-GlcNAcylation-inducing agents stimulated Akt phosphorylation (42). Interestingly, using a BRET-based assay which monitors PIP3 production at the plasma membrane in living cells (50), Kanwal et al. demonstrated that O-GlcNAcylation-inducing treatments stimulated the production of phosphatidylinositol 3,4,5-trisphosphate (PIP3) by PI3K in MCF-7 cells, suggesting that stimulation of Akt by these treatments resulted from activation of early steps in the signaling cascade (42). Several anti-cancer drugs targeting PI3K are under clinical development (51). Noteworthy, in a cancer cell line collection (60 diverse human cancer cell lines representing multiple tumor types), resistance to treatment with the PI3K inhibitor GDC-0941 correlated with OGT expression level (52). These authors also showed in different cell lines that silencing OGT increased sensitivity to GDC-0941, whereas increasing O-GlcNAc levels using PUGNAc promoted resistance to this drug (52).

On the other hand, growth factor signaling may also influence the O-GlcNAcylation pathway. For instance, it has been shown that upon insulin stimulation, OGT is recruited to the plasma membrane through a PIP3-binding domain, leading to O-GlcNAcylation and inhibition of insulin signaling intermediates (53). This is consistent with results from another study showing that insulin stimulation increased OGT shuttling from nucleus to cytoplasm. This was associated with OGT interaction with IR, allowing the latter to phosphorylate OGT on tyrosine residues and stimulate its catalytic activity (54). Very recently, OGT recruitment in response to insulin was shown to occur in lipid rafts through a PI3K/Akt dependent pathway in HepG2 cells. Notably, expression of OGT is required for proper IR expression and insulin signaling in HepG2 cells, illustrating the complex interconnections between these two pathways (55).

Growth factors can also affect O-GlcNAc signaling by regulating the expression of important enzymes of pathway. For instance, one of the first links between mitogenic signals and hexosamine pathway was provided by the observation that in MDA-468 breast cancer cells, epidermal growth factor activated the expression of the rate-limiting enzyme GFAT (56). More recently, serum stimulation was shown to increase OGT protein levels in MCF-7 cells through a post-transcriptional dependent pathway, and this effect appears to be necessary for cell growth (27). However, besides these specific examples, no systematic approaches on growth factors or hormones in relation to their ability to control O-GlcNAcylation and/or O-GlcNAc-cycling enzyme expression have been conducted. Clearly, the relevance of these mechanisms in breast cancer pathophysiology still deserves further investigations.

## O-GLcNACYLATION AND CELL CYCLE REGULATION THE FIRST OBSERVATIONS

Independent reports conducted during the 1990s supported that OGT and O-GlcNAc dynamics interfere with cell cycle progression of germinal and somatic cells.


It was first highlighted dynamic O-GlcNAcylation changes in response to mitogens (57). Activation of murine T lymphocytes with the lectin concanavalin A resulted in a rapid decrease in cytosolic proteins O-GlcNAc level and, conversely, an enhancement of O-GlcNAcylation of nuclear proteins. These observations suggested that O-GlcNAc cycling was necessary for the activation of T cells. In the human colon adenocarcinoma cell line HT29, colcemid or nocodazole-induced G2/M cell cycle arrest induced a concomitant enhancement in phosphorylation and O-GlcNAcylation of keratins K8 and K18 (two intermediate filament proteins overexpressed in cancer cells) (58). Removal of nocodazole slowly returned keratins O-GlcNAc to baseline levels, suggesting that O-GlcNAcylation of these proteins is cell cycle dependent. At this time, because of the lack of techniques allowing specific visualization of O-GlcNAcylation of proteins, the impact of O-GlcNAcylation on cell cycle progression was largely minimized. Since then, it appeared that O-GlcNAcylation actively contributes in cell cycle proceeding at different steps. For instance, inhibition of OGA using the non-selective hexosaminidase inhibitor PUGNAC

indicated that treated-cells progressed more slowly through the cell cycle than untreated ones (47).

### G0/G1

Most of the cells of the organism are in the G0 phase, a quiescent stage of non-division. Entry into G1 phase (the first phase of the cell cycle) requires the presence of exogenous mitogenic signals that lead to activation of the MAP kinase and PI3K pathways, and to transcription of the cyclin D1 gene, a key regulator of the G1 phase (Figure 2).

Recently, two independent groups reported that OGT is significantly increased following serum stimulation (G0/G1 transition) (59, 60). Blockade of OGT activity or interfering with its expression delays serum-stimulated cyclin D1 synthesis and cell proliferation (27). Kwei et al. observed also in the ovarian cancer cell line OVCAR-4 that cyclin D1 is down-regulated in siOGT-transfected cells (52). A decrease in the HBP flux also reduces the cell proliferating rate, while inhibiting OGA accelerates this process. OGT silencing diminishes PI3K and MAPK



**FIGURE 2 | O-GlcNAcylation and cell cycle.** A quiescent cell enters the cell cycle upon mitogenic signals. Cell cycle is divided into four phases: the G1 (Gap1) phase, during which cell grows, followed by the S phase of DNA replication, then the G2 (Gap2) phase which prepares the cell for the proper division phase called M phase. Progression of the cell through the different phases is highly controlled: at the G1/S and G2/M transitions a checkpoint exists to ensure that DNA is not damaged respectively before and after its replication. The G2/M checkpoint also controls that replication is ended before division of the cell into two daughter cells genetically identical. O-GlcNAcylation levels have been found to vary all along the cell cycle suggesting that it could regulate this process. OGT and O-GlcNAcylation levels increase when quiescent cells are stimulated by mitogenic signals to enter into the cell cycle (G0/G1 transition). On the contrary, OGA activity is increased at the G1/S transition leading to a decrease in global O-GlcNAcylation levels. At the G2/M checkpoint, a burst

in O-GlcNAcylation occurs. In agreement with these observations, O-GlcNAcylation has been demonstrated to be crucial for cell cycle entry and progress. Inhibition of OGT delays serum-stimulated MAPK and PI3K pathways activation and cell cycle entry whereas OGA inhibition accelerates the process. Moreover, inhibiting OGT in G2-like *Xenopus laevis* oocytes prevents entry into M phase. In addition, the levels of cyclin D1 and cyclin B1, two key regulators of the G1 and the M phase respectively, decrease when OGT is inhibited. Finally, O-GlcNAcylation could take part in the control of DNA replication since it modifies several histones and three proteins of the MCM (Mini Chromosome Maintenance Complex) 3, 6, and 7 that belong to the DNA pre-replication complex. As a consequence, deregulation of O-GlcNAcylation processes could contribute to perturbation in cell cycle control leading to anarchic proliferation, but also in accumulation of DNA mutations, two well established characteristics of cancer cells.

activation (27), demonstrating that OGT is indispensable for G0/G1 transition.

Therefore, studies that focused on G0/G1 transition indicate that OGT synthesis and a subsequent O-GlcNAcylation increase are required for cell cycle entry. It might be interesting to establish whether cyclin D is itself O-GlcNAcylated, and to determine its potential consequences on its expression and activity.

### G1/S

The S phase is the second phase of the cell cycle during which DNA replicates. At the G1/S boundary a checkpoint is set up to make sure that the DNA is not damaged. OGT, OGA, and O-GlcNAcylation levels were measured during the G1/S transition and throughout S-phase progression (59). A global decrease in O-GlcNAcylated proteins was observed during the G1/S transition and this change correlates with an increase in OGA expression and activity. In agreement with this observation, inhibition of HBP accelerated the S phase (47). Drougat et al. identified more than 50 proteins which O-GlcNAc status varies among which Minichromosome Maintenance (MCM) 3, 6, and 7, which are localized in the multiprotein pre-replicative complex needed for helicase activity and are involved in the replication of DNA (59). Zhang et al. showed that O-GlcNAcylation of histones is lower in S phase in comparison with G1, late S, and G2 phases (61) (see below for details). Accordingly, reduction of histones O-GlcNAcylation during S phase may help the pre-replicative complex to reach the chromatin.

Therefore, the few studies relating to the impact of O-GlcNAcylation on G1/S transition indicate that contrary to G0/G1, a global decrease in O-GlcNAcylation is observed. Intriguingly, not only OGA but also OGT levels increase during G1/S (59). It is tempting to hypothesize that OGT plays a function independent of its GlcNAc transferase activity as exemplified by its interaction with p38MAPK (62).

### G2/M

The S phase is followed by G2, a phase during which the cell grows and prepares for the proper division M phase (mitosis in the case of somatic cells and meiosis of germ cells). A second checkpoint exists at the G2/M transition to ensure that the DNA replication is ended and that the DNA is not damaged.

The *Xenopus laevis* oocyte has been widely used as a model for the characterization and the identification of many key-cell cycle components, such as the M-phase promoting factor (MPF) and the cytostatic factor (CSF) (63, 64) and for studying the regulation of the cell cycle, especially events occurring at G2/M. At the end of oogenesis, the oocyte is physiologically blocked in prophase of the first meiotic division in a G2-like stage; it is called immature oocyte. Upon progesterone stimulation, the oocyte resumes meiosis in a G2/M analog transition phase.

The micro-injection into *Xenopus* oocytes of bovine galactosyltransferase, an enzyme enabling the capping of terminal GlcNAc residues inhibited M-phase entry and blocked M to S-phase transition (65). Slawson and co-workers showed that the perturbation of *Xenopus* oocyte O-GlcNAcylation levels either by glucosamine or PUGNAc treatment modified the maturation kinetics (66). A few

years later, a set of studies conducted by the same group showed that hormonal stimulation of physiologically G2-blocked *Xenopus laevis* oocytes triggers a quick increase in O-GlcNAcylation levels and that inhibition of OGT impairs G2/M transition (67–70). OGT and O-GlcNAc localized on the meiotic spindle and chromosomes in metaphase-II *Xenopus laevis* oocytes. It was also observed that OGT expression and O-GlcNAcylation peaked at the M phase of the cell cycle of MEF and HEK293 cells (60). Slawson et al. observed that OGT localized to the mitotic spindle and midbody during mitosis and that its overexpression resulted in supernumerary chromosomes (47). In a second report, these authors showed that OGT and OGA interact with Aurora B and protein phosphatase 1 (PP1) to regulate the stability of the midbody and the phosphorylation and/or O-GlcNAcylation of vimentin at M phase (71). Regarding G2/M transition, both OGA and OGT knockdown decrease cyclin B1 expression, indicating that a correct expression of the two enzymes are necessary for cell cycle progression (27, 60).

As for cell cycle entry, O-GlcNAcylation increase is crucial for the cell to start mitosis. The cell cycle is a very complex process that requires a plethora of enzymes and regulatory components. These proteins were widely described to be regulated by PTM like phosphorylation, acetylation, methylation, ubiquitination, and many others. Recently, O-GlcNAcylation has joined this list of PTM, and manipulating OGT and OGA level or activity affects the progression of the cell cycle.

Altogether, the available data indicate that O-GlcNAcylation tends to increase all along the cycle with a drop at G1/S. The pattern of O-GlcNAcylation levels after the M phase and during cell cycle exit remains unknown. One could expect a dramatic decrease, a hypothesis that needs to be experimentally addressed.

## O-GLCNAcylation AND EPIGENETIC REGULATIONS

Recent whole-exon sequencing of human cancers has shown a high proportion of mutations in genes involved in regulation of DNA methylation, histone modification, and/or nucleosome remodeling (72). These discoveries firmly establish that interferences in epigenetic processes can lead to cancer and add credence to the idea that epigenetics is a major player in this disease. As described below, it is now clear that O-GlcNAcylation plays a part in the regulation of the epigenome.

Over the last 4 years, several groups focused on O-GlcNAc-mediated regulation of chromatin dynamics, a process crucial for DNA replication, DNA repair, gene expression, and mitosis (DNA compaction) (33, 73–75).

Chromatin condensation and relaxation is managed by histones that form octamers by assembling two copies of each core nucleosomal histone, H2A, H2B, H3, and H4. These oligomers interact with DNA in a nucleosome structure compacted by the linker histone H1. Histones, assisted by numerous post-translational modifications are therefore the main proteins responsible for chromatin remodeling and gene expression. Acetylation, methylation, and phosphorylation are the best-characterized histones PTM, the two formers being associated with activation of chromatin and the latter with both activation and repression [for review, see (76)]. Histones are also covalently modified by ADP-Ribosylation, SUMOylation, ubiquitination, and as recently described, O-GlcNAcylation [for review, see (77)], while O-GlcNAcylation of *Xenopus* oocytes

histones remained undetectable (69). The histone code is therefore very complex and far from being deciphered.

Sakabe and Hart pointed out the impact of OGT on histone H3 modification (74). These authors mapped three O-GlcNAcylation sites on H2A, H2B, and H4 at Thr101, Ser36, and Ser47 respectively, three phosphorylation sites necessary for the assembly of nucleosomes (74). H2B tail is also O-GlcNAcylated at Ser112 (78). This modification promotes H2B monoubiquitination at Lys120 allowing local transcription as suggested by visualization of O-GlcNAcylated H2B at Ser112 in transcribed gene loci (78). H3 tail is O-GlcNAcylated at the phosphorylation site Ser10 (61). Histone O-GlcNAcylation is coupled with other modifications associated with both active and inactive chromatin states, and covalent linkage of histones by O-GlcNAc fluctuates all along the cell cycle with a lower rate in S phase (61). Fong et al. worked on the same histone, and identified Thr32 as a major H3 O-GlcNAcylated site (75). O-GlcNAcylated H3 isoforms are higher in interphase cells than in mitosis. H3 phosphorylation at Ser10, Ser28, and Thr32 is associated to mitosis: increased H3 O-GlcNAcylation reduces phosphorylation and delays mitosis entry (75). It is noteworthy that H3 Ser10 and Ser28 are phosphorylated by Aurora B (79, 80) and dephosphorylated by PP1 (81), two enzymes physically interacting with OGT and OGA (71). It is tempting to hypothesize that regulation of histone assembly/disassembly is managed in part by a heterotetrameric complex made of Aurora B, PP1, OGT, and OGA.

O-GlcNAc-transferase and O-GlcNAc may also affect chromatin structure by modulating chromatin-remodeling enzyme activities. Fujiki et al. (82) highlighted the role O-GlcNAcylation of a histone lysine methyltransferase (MLL5) in the control of chromatin state. MLL5 O-activates RAR- $\alpha$  and interacts with OGT in an active multimeric complex. O-GlcNAcylation of MLL5 increases granulopoiesis of HL60 promyelocytes in response to retinoic acid. This facilitation occurs through an increased-MLL5 methylation induced by O-GlcNAcylation. One O-GlcNAcylation site was identified on MLL5 at Thr440 and modification of this residue potentiates its H3K4 methyltransferase activity. Another study by Sakabe and Hart also observed that moderate OGT overexpression prevented the phosphorylation of coactivator-associated arginine methyltransferase 1 (CARM1), resulting in decreased H3 methylation on Arg17 by CARM1 (74).

TET (ten-eleven translocation) proteins interact with and target OGT to chromatin (83). TET are DNA hydroxylases involved in DNA demethylation. These enzymes convert 5-methyl-Cytosine to 5-formyl-Cytosine and 5-carboxyl-Cytosine successively (84). TET are necessary for gene transcription, pre-mRNA splicing, zygotic epigenetic reprogramming, and TET mutations are responsible for myeloid cancers. In an attempt to identify TET proteins partners, Chen et al. (85) showed that OGT interacts with TET2 and TET3. While OGT does not influence TET function, physical interaction between TET2 and OGT is a precondition to address the glycosyltransferase to chromatin where it O-GlcNAcylates histones. Knockdown of TET2 decreases OGT interaction with chromatin and impairs H2A/B, H3, and H4 O-GlcNAcylation, and deregulation of TET2 more precisely reduces O-GlcNAcylation of H2B at Ser112. Genome-wide chromatin immunoprecipitation and sequencing analysis indicate that OGT and H2B Ser112 O-GlcNAc overlap a large amount of target genes with TET2 and


that the density of distribution is enriched at transcription start sites. As O-GlcNAcylation of H2B at Ser112 promotes monoubiquitination and transcriptional activation (78), TET2 may exert chromatin activation and gene expression by promoting demethylation of DNA and nucleosome opening by acting indirectly on histones. Therefore, TET2 and OGT form a complex that regulates gene transcription. Another study reports the interaction of TET1, TET2, Sin3a, and Hcfc1 with nuclear OGT (86). A genome-wide range analysis led by chromatin immunoprecipitation coupled to high-throughput DNA sequencing indicates that 62% of the 11552 OGT binding sites locate within promoter regions. Among the list of genes regulated by OGT, it was found genes involved in glycerolipid, glycerophospholipid, N- and O-glycosylation metabolism. OGT binding sites overlay with TET1 at H3K4me3 positive promoters, and both proteins interact together in the vicinity of transcription start sites enriched in unmethylated CpG-rich promoters. As for TET2, TET1 is necessary for recruiting OGT to the chromatin but it seems that, contrary to TET2 and TET3 that are neither O-GlcNAcylated nor regulated by OGT (85, 87), OGT regulates 5-hydroxymethyl-Cytosine levels by stabilizing TET1 at the promoters.

### O-GLCNAcylation REGULATES TRANSCRIPTION FACTOR ACTIVITIES

Transcription factors bind DNA to control the expression of an array of target genes, consisting for some in genuine “hubs” for cellular process regulation. Mutation in transcription factor-encoding gene, resulting in dysregulation of their expression or activity, is a common mechanism involved in many cancers (88). Nonetheless, post-translational modifications including O-GlcNAcylation also represent an important control mechanism of the expression and/or activity of transcription factors (16, 89). Rightfully, several transcription factors involved in cancer have been shown to be O-GlcNAcylated (Figure 3).

### MYC

The protein MYC which is normally expressed at low levels is largely overexpressed in proliferative cells and cancer cells. MYC is a preponderant oncogenic transcriptional factor driving forward cell cycle and replication (90). Upon growth signals, phosphorylation on Ser62 results in stimulation of MYC activity. This is a prerequisite for GSK3 $\beta$  to phosphorylate MYC at Thr58 near the transactivation domain, resulting in its degradation (91). Almost 20 years ago, Thr58 was shown to be also O-GlcNAcylated (22, 92). GSK3 $\beta$  inhibition or mutation of Ser62 to Ala increase O-GlcNAcylation of Thr58. Interestingly, in lymphomas, Thr58 is frequently mutated resulting in a more stable MYC (92). Although it has not been clearly demonstrated that O-GlcNAcylation of Thr58 *per se* is able to stabilize MYC, it is reasonable to assume that in a context of hyper-O-GlcNAcylation, phosphorylation of MYC on Thr58 by GSK3 $\beta$  will be reduced, preventing its degradation. This would result in a more stable and more active MYC, promoting tumorigenesis. Although experimental evidences are still required to support such a mechanism, a very recent report presented data in line with this hypothesis (93). In prostate cancer biopsies, OGT is upregulated in association with MYC levels. In three prostate cancer cell lines (LNCaP, VCaP, and PC3),



**FIGURE 3 | O-GlcNAcylation and oncogenic transcription factors.** Several transcription factors have been described to be O-GlcNAcylated by the O-GlcNAc-transferase (OGT) among which some that are involved in tumorigenesis. The proto-oncogene MYC has been shown to be O-GlcNAcylated on Thr58. This site is normally phosphorylated by GSK3 $\beta$  in response to MYC phosphorylation on Ser62 resulting in MYC degradation. Proper experimental evidences demonstrating that O-GlcNAcylation promotes MYC stability and pro-oncogenic activities remain to be described. NF- $\kappa$ B signaling is known to be promoted by O-GlcNAcylation. First, the IKK $\beta$  kinase responsible for NF- $\kappa$ B activation has been shown to be O-GlcNAcylated on Ser733. In a p53-null context, it was associated with pro-oncogenic activity in a model of colitis-associated cancer. Second, NF- $\kappa$ B itself is also O-GlcNAcylated at Thr322 and Thr352 in presence of high glucose concentration. In pancreatic cancer cells, Thr322 and Thr352 O-GlcNAcylation were shown to be involved in cell anchorage *in vitro*. In breast and prostatic cancer cells, OGT controls the expression of FOXO1, a pro-oncogenic transcriptional factor. However, FOXO1, itself, is not O-GlcNAcylated. Through

an unknown substrate and mechanism, OGT prevents FOXO1 degradation and thus promotes tumor development in breast cancer and metastasis in prostate cancer.  $\beta$ -Catenin, involved in cell adherent junctions, is also central to the Wnt/ $\beta$ -catenin pathway as a transcription factor in complex notably with LEF/TCF co-factors. Once activated,  $\beta$ -catenin promotes tumorigenesis, especially colorectal and liver cancers. Increased O-GlcNAcylation levels of  $\beta$ -catenin have been observed in colon carcinoma cells. Furthermore, interaction of OGT with  $\beta$ -catenin is stimulated during serum-induced proliferation in HeLa cells. These observations support a positive role of O-GlcNAc-modified  $\beta$ -catenin in cell proliferation. The well known tumor suppressor p53 is O-GlcNAcylated at Ser149 which prevents the phosphorylation at Thr155 by the COP9 signalosome. This results in an inhibition of p53 degradation. O-GlcNAcylation of p53 promotes p53 stability and thus its tumor suppressor activity in physiological context. In case of pro-oncogenic p53 mutants, O-GlcNAc-mediated stability of p53 may favor pro-oncogenic processes. This hypothesis has not been addressed experimentally yet.

pharmacologically inhibition of OGT decreased MYC protein stability without affecting its mRNA levels. The authors confirmed O-GlcNAcylation of MYC in LNCaP cell line suggesting that this PTM could be responsible for MYC stabilization (93). However, to our knowledge, the O-GlcNAc status of MYC in patient tumors still remains to be evaluated.

### p53

p53 is a central tumor suppressor whose gene is mutated in more than 50% of mutations in cancers (94). p53 level is tightly controlled by PTM (95). Upon various stress signals (DNA damage, oncogenic events, hypoxia...), p53 levels are stabilized through inhibition of proteasomal degradation. This is achieved by phosphorylation of p53, which blocks its interaction with the E3 ubiquitin ligase MDM2 (95). However, phosphorylation-induced degradation of p53 can also occur, as illustrated by phosphorylation performed by COP9 signalosome on Thr155 in the DNA-binding region of p53 (96). More recently, Yang et al. showed that O-GlcNAcylation of p53 at residue Ser149 inhibits its phosphorylation on Thr155 by COP9 signalosome, hence promoting p53 stabilization and activity (97). In the case of a wild-type p53,

O-GlcNAcylation should promote its tumor suppressor activity, as suggested by higher apoptosis in H1299 cells (97). However, examples of gain-of-function mutant of p53 favoring tumorigenesis have been described in the literature (98–100). In this context, stabilization-induced O-GlcNAcylation of gain-of-function mutant form of p53 may amplify its pro-oncogenic activity. Clearly, this hypothesis requires to be addressed experimentally.

### NF- $\kappa$ B

NF- $\kappa$ B transcription factors are involved in a wide variety of physiological and pathological processes including, immunity and inflammation, metabolism, and cancer (101, 102). In the basal state, NF- $\kappa$ B is sequestered by I $\kappa$ B in the cytosol. Canonical activation takes place by activation of I $\kappa$ B kinase complex which in turn phosphorylates and triggers I $\kappa$ B degradation. NF- $\kappa$ B is then released and enters the nucleus to activate its target genes (101). Originally, Yang et al. identified residues Thr322 and Thr352 as the sites of O-GlcNAc modification. However, only O-GlcNAcylation of Thr352 was central for transcriptional activation (103). O-GlcNAcylation of NF- $\kappa$ B in relation to cancer has not been directly addressed until recently. It was shown that mutation

of Thr322 and Thr352 to Ala, reduced the anchorage of pancreatic ductal adenocarcinoma cells *in vitro* (104). Therefore, it will be of great interest to determine *in vivo* the involvement of NF- $\kappa$ B O-GlcNAcylation in the context of tumorigenesis.

Links between NF- $\kappa$ B pathway and O-GlcNAcylation have also been characterized upstream NF- $\kappa$ B factors, in relation to the activating kinase IKK $\beta$ . In a context of p53-null mutation, aerobic glycolysis is upregulated through an IKK-NF- $\kappa$ B pathway (105). It was shown that IKK $\beta$  is activated by phosphorylation concomitantly with an increase in its Ser733 O-GlcNAcylation (106). Ser733 is usually the target of an inhibitory auto-phosphorylation by the catalytic domain of activated IKK $\beta$ . O-GlcNAcylation of Ser733 prevents inhibition of IKK $\beta$  which results in a sustained activity. Consequently, NF- $\kappa$ B pathway remains active, and this further promotes glycolysis (106). The role of NF- $\kappa$ B pathway in tumorigenesis is complex, either pro- or anti-oncogenic, depending on cancer type, environment, inflammation status, or p53 functionality (105, 107–110). Whereas it now clearly appears that O-GlcNAcylation play a significant role in NF- $\kappa$ B pathway, its involvement in tumorigenesis still remains to be precisely defined.

### FOXM1

FOXM1 is a transcriptional regulator expressed at high level in various cancers, pointing out its importance in tumorigenesis (111). Links between FOXM1 and O-GlcNAcylation pathway have been made by Reginato's laboratory (112). These authors showed that high levels of OGT are responsible for the increased activity of FOXM1 in breast cancer MCF-10A cells overexpressing the activated form of ErbB2. Although no O-GlcNAcylation form of FOXM1 could be detected, OGT expression was shown to be central to the control of FOXM1 levels and to its effects on cell proliferation (112). FOXM1 levels were dramatically decreased by OGT silencing. Consequently, the level of the FOXM1's target S-phase kinase-associated protein 2 (Skp2), involved in the SCF<sup>SKP2</sup> E3 ubiquitin ligase complex responsible for p27<sup>Kip1</sup> degradation, was markedly decreased. This resulted in an increase in cell cycle inhibitor p27<sup>Kip1</sup> protein levels and subsequent cell cycle arrest (112). Similarly, the same group showed that inhibition of OGT in the prostate cancer cell line PC3-ML was associated with an increase in FOXM1 degradation through a proteasome-mediated process (37). It was also observed that loss of FOXM1 upon modulation of OGT levels in PC3-ML cells affects their ability to form bone metastasis in mice (see below). Overexpression of a degradation-resistant mutant of FOXM1 rescued the effects of OGT inhibition (37). However, the direct target of OGT involved in the control of FOXM1 expression remains to be identified.

### $\beta$ -CATENIN

$\beta$ -Catenin is a versatile protein playing fundamental roles in cells from control of intercellular junction integrity through its interaction with cadherins and cytoskeleton, to regulation of transcriptional processes as a co-transcription factor. As a cell cycle and proliferation regulator,  $\beta$ -catenin is a key partner in the Wnt/ $\beta$ -catenin pathway. Upon Wnt ligand interaction with its receptor,  $\beta$ -catenin sequestered by the Adenomatous Polyposis Coli (APC)/axin degradation complex is released and re-localized to the nucleus, where it interacts with TCF/LEF or other co-factors

to form a transcriptional complex able to modulate the expression of a number of target genes (113). Mutations of  $\beta$ -catenin itself (accounting for 10% of mutations) or APC (one of the main component of the destruction complex, accounting for 80% of mutations), are found in 85–90% of CRC. These mutations increase  $\beta$ -catenin stability which therefore acquires oncogenic properties.

Recently, in an attempt to understand the link between nutrition and CRC, Olivier-Van Stichelen and colleagues showed that  $\beta$ -catenin expression correlated with the HBP flux and the O-GlcNAcylation levels in colon carcinoma cells (114) as previously pointed out in macrophages (115). Later, Olivier-Van Stichelen et al. showed that serum-induced proliferation increases the interaction between OGT and  $\beta$ -catenin in HeLa cells, supporting a positive role for O-GlcNAc-modified  $\beta$ -catenin in cell cycle progression (27). It is therefore proposed that hyper-O-GlcNAcylation in the colic and rectal mucosa may constitute a pro-oncogenic signal, explaining why metabolic disorders and over-nutrition increase the risk of CRC.

### OTHER TRANSCRIPTION FACTORS

Obviously, other transcription factors involved in cancer have been shown to be O-GlcNAc-modified as hypermethylated in cancer 1 (HIC1) (24), Jun (116, 117), Estrogen Receptor  $\beta$  (118, 119), the chimeric transcription factor EWS-FLI1 expressed in Ewing's sarcoma family tumors (120). However, significant work remains to be performed to better understand the exact role of the O-GlcNAc-modified residues of oncogenic-related transcription factors and their function in tumorigenesis.

### O-GLCNAcylation REGULATES CELL ADHESION AND MIGRATION

Numerous studies indicate that O-GlcNAc promotes cancer cell invasiveness and metastasis. Thus, manipulation of O-GlcNAc levels using chemical or shRNA-mediated inhibition of OGA or OGT affects *in vitro* and *in vivo* migration/invasion of breast (31, 112, 121), lung (31, 33), liver (34), prostate (37), and colon (33) cancer cells. Several lines of evidence suggest that O-GlcNAc favors cancer malignancy by impacting the E-cadherin/ $\beta$ -catenin system and by promoting expression of metalloproteinases.

### E-CADHERIN AND CANCER METASTASIS

E-cadherin, the prototypic member of the cadherin family, is a major component of adherent junctions. E-cadherin regulates cell–cell adhesion via homophilic interactions between its extracellular domains on opposing plasma membranes and by binding of its intracellular domain to the cytoskeleton via  $\beta$ -catenin. Down-regulation of E-cadherin is a critical step in the epithelial-mesenchymal transition (EMT), characteristic of carcinoma invasion. In breast cancer 4T1 cells, E-cadherin protein expression was increased by OGT shRNA, while it was decreased by treatment with the pharmacological OGA inhibitors PUGNAC and NButGT (31). Moreover, co-immunoprecipitation experiments showed that E-cadherin interaction with  $\beta$ -catenin and p120 was markedly increased in OGT depleted cells and decreased by OGA inhibitors (31). In addition, analysis of Triton-X100 soluble and insoluble fraction showed that association of E-cadherin,  $\beta$ -catenin, and p120 with cytoskeleton was increased

by shOGT and decreased in OGA inhibited cells. Immunofluorescence experiments indicated that shOGT increased colocalization of E-cadherin,  $\beta$ -catenin, and p120 at the cell surface, whereas OGA inhibition reduced it. *In vivo* 4T1 cells metastasis in the lung was markedly reduced by OGT shRNA, and inhibition of E-cadherin expression in these cells (shOGT + shCadherin) restored invasiveness to control levels, suggesting an important role for regulation of E-cadherin expression by O-GlcNAc in cancer cell invasiveness (31). Whereas no change in E-cadherin O-GlcNAcylation was reported in this study, O-GlcNAc could be detected on p120 and  $\beta$ -catenin, suggesting a potential mechanism for O-GlcNAc-induced inhibition of E-cadherin cell surface localization. In addition, works from other laboratories suggested that O-GlcNAcylation regulates E-cadherin expression at the transcriptional level (34, 121). Indeed, increasing O-GlcNAcylation through siRNA-mediated inhibition of OGA down-regulated E-cadherin mRNA expression in LC cells, whereas decreasing OGT expression had the opposite effect (34). Interestingly, Snail1, a key regulator of EMT program, and a major transcriptional repressor of E-cadherin, was O-GlcNAcylated in hyperglycemic conditions in an OGT-dependent manner (121). GSK3-mediated phosphorylation and degradation of Snail1 is suppressed by its O-GlcNAcylation. Stabilization of Snail1 increases its repressor function, resulting in inhibition of E-cadherin mRNA expression (121). Therefore, O-GlcNAc regulation decreases E-cadherin localization at the membrane through inhibition of its interaction with its partners p120 and  $\beta$ -catenin, but also through stabilization of its transcriptional repressor Snail1. Consistently, a recent study showed a positive correlation between O-GlcNAc protein levels, cell migration, and E-cadherin protein levels in ovarian cancer cells. In HO-8910PM cells, down-regulation of OGT resulted in an increase in E-cadherin protein content. On the opposite, increased O-GlcNAcylation level through PUGNAc or Thiamet-G treatment decreased E-cadherin protein content in OVCAR3 cells (122). The authors showed that increased O-GlcNAcylation inhibits E-cadherin-catenin complex formation and that both E-cadherin, p120 and  $\beta$ -catenin can be O-GlcNAcylated in these cells (122). Identification of the O-GlcNAc-modified sites in the different partners involved will be required to precisely understand the mechanism involving O-GlcNAcylation in E-cadherin-dependent cancer cell migration.

### INCREASED EXPRESSION OF MATRIX METALLOPROTEINASES

Matrix metalloproteinases (MMPs) are zinc-finger dependent extracellular matrix (ECM) remodeling enzymes. Overexpression of MMPs has been associated with epithelial to mesenchymal transition. MMPs play crucial roles in invasion and metastasis, through proteolytic degradation of ECM, alteration of cell–cell and cell–ECM interactions, migration, and angiogenesis. The proteolytic activity of MMPs is required for a cancer cell to degrade physical barriers during local expansion, intravasation at nearby blood vessels, as well as extravasation and invasion at a distant location (123). Reginato's group first demonstrated negative regulation of MMP2 expression upon OGT inhibition using OGT shRNA in breast cancer cells (an effect probably mediated by down-regulation of FOXM1 protein), associated with decreased cell invasiveness (112). In LCs cells, down-regulation of OGT expression

using siRNA resulted in decreased expression of MMP1, MMP2, and MMP3, correlated with the decreased migrating and invasive capabilities of these cells, whereas down-regulation of OGA had the opposite effects (34). More recently, the Reginato's group showed that reducing O-GlcNAc in prostate cancer cells also reduced the expression of MMP2 and MMP9, associated with decreased FOXM1 expression (37). Inhibition of OGT expression in these cells was also associated with decreased *in vitro* invasiveness and *in vivo* metastasis. Interestingly, VEGF expression and angiogenesis were also inhibited by OGT siRNA, further emphasizing the involvement of O-GlcNAcylation in malignant properties of the cells. O-GlcNAcylation then participates in an integrative strategy that promotes invasiveness and metastasis by increasing ECM degradation in order to clear the way for migration.

### CONCLUSION AND FUTURE DIRECTIONS

Our knowledge regarding the regulation of cell cycle by O-GlcNAcylation progresses at a relatively high rate. Conversely, apprehending the role of O-GlcNAcylation in cancer processes is only at its beginnings. As reviewed here, there are many proofs that O-GlcNAcylation processes are perturbed in different kinds of cancers. Mainly, it appeared that O-GlcNAcylation and its cycling enzymes are upregulated in cancers and that this elevation is positively correlated with the grade or the aggressiveness of the tumor. It also appeared that metastases harbor higher O-GlcNAcylation levels than the primary tumor. Nevertheless, as mentioned previously, a number of contradictions concerning the expression or activity of O-GlcNAc-cycling enzymes also appeared (30–36, 40, 46). The large variety of different types of cancers and cellular models used in these studies, and the fact that O-GlcNAc level may itself modulate the expression of cycling enzymes deeply complicates the interpretation of the data. Future development should certainly include detailed analysis of the regulation of the promoters of OGT and OGA, and the study of the transcription factors that control the activities of these promoters in different types of normal and cancer cells.

The lack of tools necessary to specifically study the impact of O-GlcNAcylation on carcinogenesis may also partly explain contradictions observed in literature and slow down the progression toward understanding whether and how O-GlcNAc and its cycling enzymes influence the development of cancers. A striking feature of O-GlcNAcylation in cancer is the variety of proteins targeted by this PTM. In addition to cell cycle proteins, signaling intermediates and transcription factors already discussed in this review, several other types proteins have been shown to be O-GlcNAcylated in cancer, including metabolic enzymes (124) and heat shock protein (125, 126). From this impressive diversity, the fact that only two enzymes, OGT and OGA, are the only catalytic proteins in charge, calls for further identification of their co-factors and targets and for a better understanding of their mutual interaction, as recently exemplified with TET2/3 and OGT in epigenetic control (85, 87).

Modulating O-GlcNAc pathway could constitute a promising approach for anti-cancer therapy, which could be used in synergy with conventional treatments. This will be possible with systematic identification of O-GlcNAc-modified sites on the proteins of interest, which remains the prerequisite to understand the role

of this PTM in their function. Our understanding in O-GlcNAc proteome is rising exponentially (20). However, strategies to alter O-GlcNAc-cycling enzymes and their targets in order to specifically block cancer-related processes still need to be developed. The important number of cancer types renders difficult to predict how conserved a mechanism would be from one tumor to the other. A way to tackle this issue may come from the determination of the O-GlcNAcylated protein signature in the different cancers. This clearly remains a wide field for investigation. Although pharmacological targeting of OGT is possible *in vitro* (127), important work is still required for *in vivo* application. Indeed, keeping in mind that O-GlcNAcylation in normal cells should remain untouched, targeting OGT specifically in cancer cell *in vivo* may represent one of the most important challenges in this field. O-GlcNAc-modulating strategies will indeed require caution, as increasing amount of evidences suggests that O-GlcNAcylation regulates highly important biological processes such as epigenetic (77) and affects molecular mechanisms involved in major human diseases such as Alzheimer's disease, cardiovascular disorders, and diabetes (17, 18).

As we have seen throughout this review, considerable amount of evidence indicates that O-GlcNAcylation plays an important role in cell in cancer, through a broad panel of mechanisms (cell cycle, chromatin dynamics, transcription factors, kinases, and phosphatases involved in cell signaling, cell adhesion. . .). However, it is not clear whether cancer development requires increase in global protein O-GlcNAcylation or only modification of specific key proteins. In addition, as O-GlcNAcylation level tightly depends on nutrient availability, an important challenge will be to establish, for each protein identified as modified by O-GlcNAc in cancer cells, whether this modification is a cause or a consequence of the cancerous phenotype of the cell. Indeed, metabolic reprogramming,

one of the so-called cancer hallmarks, include increased glucose and glutamine uptake, which might be sufficient to increase O-GlcNAc level in proteins. This may promote a general increase in O-GlcNAc level on proteins, including in proteins more specifically involved in the maintenance of the cancerous phenotype, leaving the question of causality difficult to answer.

Finally, because the analogy between O-GlcNAcylation and phosphorylation is striking, we are always tempted to apply to O-GlcNAc signaling reasoning schemes and concepts taken from our classical view of phosphorylation cascades and their alterations in cancer cells. However, important differences should be kept in mind. Unlike phosphorylation, O-GlcNAcylation, which is performed by a single enzyme, cannot organize in signaling cascades in which one O-GlcNAcylated protein will transmit the signal by O-GlcNAcylation another protein. Rather than acting as a switch that turns on or off signaling pathways, O-GlcNAcylation should be considered as a "rheostat" (17) that controls the intensity of signals traveling through various pathways according to nutrient and stress environment. Whether one of the ancestral functions selected by evolution in unicellular organisms was to use this "rheostat" to sense nutrient availability and promote proliferation when the nutritional environment was favorable remains an open question that might provide some clues to understand why O-GlcNAcylation appears to be central to cancer cell biology.

## ACKNOWLEDGMENTS

Our work was supported by the "Association pour la Recherche sur le Cancer" (Grant no. 1069 and Grant no. 75958), the "Ligue contre le Cancer" (Comité de Paris, Grant no. RS10/75-46, and Comité Scientifique du Septentrion), and by the ANR (ANR grant Genopath Diab-O-Glyc).

## REFERENCES

- Dossus L, Kaaks R. Nutrition, metabolic factors and cancer risk. *Best Pract Res Clin Endocrinol Metab* (2008) **22**:551–71. doi:10.1016/j.beem.2008.08.003
- Roberts DL, Dive C, Renehan AG. Biological mechanisms linking obesity and cancer risk: new perspectives. *Annu Rev Med* (2010) **61**:301–16. doi:10.1146/annurev.med.080708.082713
- Simon D, Balkau B. Diabetes mellitus, hyperglycaemia and cancer. *Diabetes Metab* (2010) **36**:182–91. doi:10.1016/j.diabet.2010.04.001
- Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* (2004) **4**:579–91. doi:10.1038/nrc1408
- Ashrafian H, Ahmed K, Rowland SP, Patel VM, Gooderham NJ, Holmes E, et al. Metabolic surgery and cancer: protective effects of bariatric procedures. *Cancer* (2010) **117**:1788–99. doi:10.1002/cncr.25738
- Dogan S, Hu X, Zhang Y, Maihle NJ, Grande JP, Cleary MP. Effects of high-fat diet and/or body weight on mammary tumor leptin and apoptosis signaling pathways in MMTV-TGF- $\alpha$  mice. *Breast Cancer Res* (2007) **9**:R91. doi:10.1186/bcr1840
- Gillette CA, Zhu Z, Westerlind KC, Melby CL, Wolfe P, Thompson HJ. Energy availability and mammary carcinogenesis: effects of calorie restriction and exercise. *Carcinogenesis* (1997) **18**:1183–8. doi:10.1093/carcin/18.6.1183
- Colman RJ, Anderson RM, Johnson SC, Kastman EK, Kosmatka KJ, Beasley TM, et al. Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science* (2009) **325**:201–4. doi:10.1126/science.1173635
- Willcox BJ, Willcox DC, Todoriki H, Fujiyoshi A, Yano K, He Q, et al. Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. *Ann N Y Acad Sci* (2007) **1114**:434–55. doi:10.1196/annals.1396.037
- Willcox DC, Willcox BJ, Todoriki H, Suzuki M. The Okinawan diet: health implications of a low-calorie, nutrient-dense, antioxidant-rich dietary pattern in glycemic load. *J Am Coll Nutr* (2009) **28**(Suppl):500S–16. doi:10.1080/07315724.2009.10718117
- Garofalo C, Koda M, Cascio S, Sulkowska M, Kanczuga-Koda L, Golaszewska J, et al. Increased expression of leptin and the leptin receptor as a marker of breast cancer progression: possible role of obesity-related stimuli. *Clin Cancer Res* (2006) **12**:1447–53. doi:10.1158/1078-0432.CCR-05-1913
- Cirillo D, Rachiglio AM, La Montagna R, Giordano A, Normanno N. Leptin signaling in breast cancer: an overview. *J Cell Biochem* (2008) **105**:956–64. doi:10.1002/jcb.21911
- Deberardinis RJ, Sayed N, Ditsworth D, Thompson CB. Brick by brick: metabolism and tumor cell growth. *Curr Opin Genet Dev* (2008) **18**:54–61. doi:10.1016/j.gde.2008.02.003
- Hanover JA. Glycan-dependent signaling: O-linked N-acetylglucosamine. *FASEB J* (2001) **15**:1865–76. doi:10.1096/fj.01-0094rev
- Hart GW, Housley MP, Slawson C. Cycling of O-linked beta-N-acetylglucosamine on nucleocytoplasmic proteins. *Nature* (2007) **446**:1017–22. doi:10.1038/nature05815
- Issad T, Kuo M. O-GlcNAc modification of transcription factors, glucose sensing and glucotoxicity. *Trends Endocrinol Metab* (2008) **19**:380–9. doi:10.1016/j.tem.2008.09.001
- Issad T, Masson E, Pagesy P. O-GlcNAc modification, insulin signaling and diabetic complications. *Diabetes Metab* (2010) **36**:423–35. doi:10.1016/j.diabet.2010.09.001
- Lefebvre T, Dehennaut V, Guinez C, Olivier S, Drougat L, Mir AM, et al. Dysregulation of the nutrient/stress sensor O-GlcNAcylation is involved in the etiology of cardiovascular disorders, type-2 diabetes and Alzheimer's disease. *Biochim Biophys Acta* (2010) **1800**:67–79. doi:10.1016/j.bbagen.2009.08.008

19. Hart GW, Slawson C, Ramirez-Correa G, Lagerlof O. Cross talk between O-GlcNAcylation and phosphorylation: roles in signaling, transcription, and chronic disease. *Annu Rev Biochem* (2011) **80**:825–58. doi:10.1146/annurev-biochem-060608-102511
20. Copeland RJ, Han G, Hart GW. O-GlcNAcomics – revealing roles of O-GlcNAcylation in disease mechanisms and development of potential diagnostics. *Proteomics Clin Appl* (2013). doi:10.1002/prca.201300001
21. Ruan HB, Singh JP, Li MD, Wu J, Yang X. Cracking the O-GlcNAc code in metabolism. *Trends Endocrinol Metab* (2013) **24**:301–9. doi:10.1016/j.tem.2013.02.002
22. Chou TY, Dang CV, Hart GW. Glycosylation of the c-Myc transactivation domain. *Proc Natl Acad Sci U S A* (1995) **92**:4417–21. doi:10.1073/pnas.92.10.4417
23. Chou TY, Hart GW. O-linked N-acetylglucosamine and cancer: messages from the glycosylation of c-Myc. *Adv Exp Med Biol* (2001) **491**:413–8. doi:10.1007/978-1-4615-1267-7\_26
24. Lefebvre T, Pinte S, Guerardel C, Deltour S, Martin-Soudant N, Sloimianny MC, et al. The tumor suppressor HIC1 (hypermethylated in cancer 1) is O-GlcNAc glycosylated. *Eur J Biochem* (2004) **271**:3843–54. doi:10.1111/j.1432-1033.2004.04316.x
25. Kuo M, Zilberfarb V, Gangneux N, Christeff N, Issad T. O-GlcNAc modification of FoxO1 increases its transcriptional activity: a role in the glucotoxicity phenomenon? *Biochimie* (2008) **90**:679–85. doi:10.1016/j.biochi.2008.03.005
26. Kuo M, Zilberfarb V, Gangneux N, Christeff N, Issad T. O-glycosylation of FoxO1 increases its transcriptional activity towards the glucose 6-phosphatase gene. *FEBS Lett* (2008) **582**:829–34. doi:10.1016/j.febslet.2008.02.010
27. Olivier-Van Stichelen S, Drougou L, Dehennaut V, El Yazidi-Belkoura I, Guinez C, Mir AM, et al. Serum-stimulated cell cycle entry promotes ncOGT synthesis required for cyclin D expression. *Oncogenesis* (2012) **1**:e36. doi:10.1038/oncsis.2012.36
28. Slawson C, Copeland RJ, Hart GW. O-GlcNAc signaling: a metabolic link between diabetes and cancer? *Trends Biochem Sci* (2010) **35**:547–55. doi:10.1016/j.tibs.2010.04.005
29. Slawson C, Hart GW. O-GlcNAc signalling: implications for cancer cell biology. *Nat Rev Cancer* (2011) **11**:678–84. doi:10.1038/nrc3114
30. Slawson C, Pidala J, Potter R. Increased N-acetyl-beta-glucosaminidase activity in primary breast carcinomas corresponds to a decrease in N-acetylglucosamine containing proteins. *Biochim Biophys Acta* (2001) **1537**:147–57. doi:10.1016/S0925-4439(01)00067-9
31. Gu Y, Mi W, Ge Y, Liu H, Fan Q, Han C, et al. GlcNAcylation plays an essential role in breast cancer metastasis. *Cancer Res* (2010) **70**:6344–51. doi:10.1158/0008-5472.CAN-09-1887
32. Krzeslak A, Forma E, Bernaciak M, Romanowicz H, Brys M. Gene expression of O-GlcNAc cycling enzymes in human breast cancers. *Clin Exp Med* (2011) **12**:61–5. doi:10.1007/s10238-011-0138-5
33. Mi W, Gu Y, Han C, Liu H, Fan Q, Zhang X, et al. O-GlcNAcylation is a novel regulator of lung and colon cancer malignancy. *Biochim Biophys Acta* (2011) **1812**:514–9. doi:10.1016/j.bbadis.2011.01.009
34. Zhu Q, Zhou L, Yang Z, Lai M, Xie H, Wu L, et al. O-GlcNAcylation plays a role in tumor recurrence of hepatocellular carcinoma following liver transplantation. *Med Oncol* (2012) **29**:985–93. doi:10.1007/s12032-011-9912-1
35. Rozanski W, Krzeslak A, Forma E, Brys M, Blewniewski M, Wozniak P, et al. Prediction of bladder cancer based on urinary content of MGEA5 and OGT mRNA level. *Clin Lab* (2012) **58**:579–83.
36. Krzeslak A, Wojcik-Krowiranda K, Forma E, Bienkiewicz A, Brys M. Expression of genes encoding for enzymes associated with O-GlcNAcylation in endometrial carcinomas: clinicopathologic correlations. *Ginekol Pol* (2012) **83**:22–6.
37. Lynch TP, Ferrer CM, Jackson SR, Shahriari KS, Vosseller K, Reginato MJ. Critical role of O-linked beta-N-acetylglucosamine transferase in prostate cancer invasion, angiogenesis, and metastasis. *J Biol Chem* (2012) **287**:11070–81. doi:10.1074/jbc.M111.302547
38. Shi Y, Tomic J, Wen F, Shaha S, Bahlo A, Harrison R, et al. Aberrant O-GlcNAcylation characterizes chronic lymphocytic leukemia. *Leukemia* (2010) **24**:1588–98. doi:10.1038/leu.2010.152
39. Pichard C, Plu-Bureau G, Neves ECM, Gompel A. Insulin resistance, obesity and breast cancer risk. *Maturitas* (2008) **60**:19–30. doi:10.1016/j.maturitas.2008.03.002
40. Dahl E, Sadr-Nabavi A, Klopocki E, Betz B, Grube S, Kreutzfeld R, et al. Systematic identification and molecular characterization of genes differentially expressed in breast and ovarian cancer. *J Pathol* (2005) **205**:21–8. doi:10.1002/path.1687
41. Champattanachai V, Netsiri-sawan P, Chaiyawat P, Phueaouan T, Charoenwattanasatien R, Chokchaichamnankit D, et al. Proteomic analysis and abrogated expression of O-GlcNAcylated proteins associated with primary breast cancer. *Proteomics* (2013) **13**:2088–99. doi:10.1002/pmic.201200126
42. Kanwal S, Fardini Y, Pagesy P, N'tumba-Byn T, Pierre-Eugène C, Masson E, et al. O-GlcNAcylation-inducing treatments inhibit estrogen receptor  $\alpha$  expression and confer resistance to 4-OH-Tamoxifen in human breast cancer-derived MCF-7 cells. *PLoS ONE* (2013) **8**(7):e69150. doi:10.1371/journal.pone.0069150
43. Hurvitz SA, Pietras RJ. Rational management of endocrine resistance in breast cancer: a comprehensive review of estrogen receptor biology, treatment options, and future directions. *Cancer* (2008) **113**:2385–97. doi:10.1002/cncr.23875
44. Pais R, Silaghi H, Silaghi AC, Rusu ML, Dumitrascu DL. Metabolic syndrome and risk of subsequent colorectal cancer. *World J Gastroenterol* (2009) **15**:5141–8. doi:10.3748/wjg.15.5141
45. Zhang ZJ, Zheng ZJ, Shi R, Su Q, Jiang Q, Kip KE. Metformin for liver cancer prevention in patients with type 2 diabetes: a systematic review and meta-analysis. *J Clin Endocrinol Metab* (2012) **97**:2347–53. doi:10.1210/jc.2012-1267
46. Krzeslak A, Pomorski L, Lipinska A. Elevation of nucleocytoplasmic beta-N-acetylglucosaminidase (O-GlcNAcase) activity in thyroid cancers. *Int J Mol Med* (2010) **25**:643–8. doi:10.3892/ijmm\_00000387
47. Slawson C, Zachara NE, Vosseller K, Cheung WD, Lane MD, Hart GW. Perturbations in O-linked beta-N-acetylglucosamine protein modification cause severe defects in mitotic progression and cytokinesis. *J Biol Chem* (2005) **280**:32944–56. doi:10.1074/jbc.M503396200
48. Hursting SD, Lavigne JA, Berrigan D, Perkins SN, Barrett JC. Calorie restriction, aging, and cancer prevention: mechanisms of action and applicability to humans. *Annu Rev Med* (2003) **54**:131–52. doi:10.1146/annurev.med.54.101601.152156
49. Kalaany NY, Sabatini DM. Tumours with PI3K activation are resistant to dietary restriction. *Nature* (2009) **458**:725–31. doi:10.1038/nature07782
50. Pierre-Eugène C, Pagesy P, Nguyen TT, Neullé M, Tschank G, Tenaegels N, et al. Effect of insulin analogues on insulin/IGF1 hybrid receptors: increased activation by glargine but not by its metabolites M1 and M2. *PLoS ONE* (2012) **7**(7):e41992. doi:10.1371/journal.pone.0041992
51. Engelman JA. Targeting PI3K signalling in cancer: opportunities, challenges and limitations. *Nat Rev Cancer* (2009) **9**:550–62. doi:10.1038/nrc2664
52. Kwei KA, Baker JB, Pelham RJ. Modulators of sensitivity and resistance to inhibition of PI3K identified in a pharmacogenomic screen of the NCI-60 human tumor cell line collection. *PLoS ONE* (2012) **7**:e46518. doi:10.1371/journal.pone.0046518
53. Yang X, Ongusaha PP, Miles PD, Havstad JC, Zhang F, So WV, et al. Phosphoinositide signalling links O-GlcNAc transferase to insulin resistance. *Nature* (2008) **451**:964–9. doi:10.1038/nature06668
54. Whelan SA, Lane MD, Hart GW. Regulation of the O-linked beta-N-acetylglucosamine transferase by insulin signaling. *J Biol Chem* (2008) **283**:21411–7. doi:10.1074/jbc.M800677200
55. Perez-Cervera Y, Dehennaut V, Aquino Gil M, Guedri K, Solorzano Mata CJ, Olivier-Van Stichelen S, et al. Insulin signaling controls the expression of O-GlcNAc transferase and its interaction with lipid microdomains. *FASEB J* (2013). doi:10.1096/fj.12-217984
56. Paterson AJ, Kudlow JE. Regulation of glutamine:fructose-6-phosphate amidotransferase gene transcription by epidermal growth factor and glucose. *Endocrinology* (1995) **136**:2809–16. doi:10.1210/en.136.7.2809
57. Kearsse KP, Hart GW. Lymphocyte activation induces rapid changes in nuclear and cytoplasmic glycoproteins. *Proc Natl Acad Sci U S A*

- A (1991) **88**:1701–5. doi:10.1073/pnas.88.5.1701
58. Haltiwanger RS, Philipsberg GA. Mitotic arrest with nocodazole induces selective changes in the level of O-linked N-acetylglucosamine and accumulation of incompletely processed N-glycans on proteins from HT29 cells. *J Biol Chem* (1997) **272**:8752–8. doi:10.1074/jbc.272.13.8752
59. Drougat L, Olivier-Van Stichelen S, Mortuaire M, Foulquier F, Lacoste AS, Michalski JC, et al. Characterization of O-GlcNAc cycling and proteomic identification of differentially O-GlcNAcylated proteins during G1/S transition. *Biochim Biophys Acta* (2012) **1820**:1839–48. doi:10.1016/j.bbagen.2012.08.024
60. Yang YR, Song M, Lee H, Jeon Y, Choi EJ, Jang HJ, et al. O-GlcNAcase is essential for embryonic development and maintenance of genomic stability. *Aging Cell* (2012) **11**:439–48. doi:10.1111/j.1474-9726.2012.00801.x
61. Zhang S, Roche K, Nasheuer HP, Lowndes NF. Modification of histones by sugar beta-N-acetylglucosamine (GlcNAc) occurs on multiple residues, including histone H3 serine 10, and is cell cycle-regulated. *J Biol Chem* (2011) **286**:37483–95. doi:10.1074/jbc.M111.284885
62. Cheung WD, Hart GW. AMP-activated protein kinase and p38 MAPK activate O-GlcNAcylation of neuronal proteins during glucose deprivation. *J Biol Chem* (2008) **283**:13009–20. doi:10.1074/jbc.M801222200
63. Masui Y, Markert CL. Cytoplasmic control of nuclear behavior during meiotic maturation of frog oocytes. *J Exp Zool* (1971) **177**:129–45. doi:10.1002/jez.1401770202
64. Masui Y. From oocyte maturation to the in vitro cell cycle: the history of discoveries of maturation-promoting factor (MPF) and cytosolic factor (CSF). *Differentiation* (2001) **69**:1–17. doi:10.1046/j.1432-0436.2001.690101.x
65. Fang B, Miller MW. Use of galactosyltransferase to assess the biological function of O-linked N-acetyl-d-glucosamine: a potential role for O-GlcNAc during cell division. *Exp Cell Res* (2001) **263**:243–53. doi:10.1006/excr.2000.5110
66. Slawson C, Shafii S, Amburgey J, Potter R. Characterization of the O-GlcNAc protein modification in *Xenopus laevis* oocyte during oogenesis and progesterone-stimulated maturation. *Biochim Biophys Acta* (2002) **1573**:121–9. doi:10.1016/S0304-4165(02)00369-0
67. Dehennaut V, Lefebvre T, Sellier C, Leroy Y, Gross B, Walker S, et al. O-linked N-acetylglucosaminyltransferase inhibition prevents G2/M transition in *Xenopus laevis* oocytes. *J Biol Chem* (2007) **282**:12527–36. doi:10.1074/jbc.M700444200
68. Dehennaut V, Hanouille X, Bodart JF, Vilain JP, Michalski JC, Landrieu I, et al. Microinjection of recombinant O-GlcNAc transferase potentiates *Xenopus* oocytes M-phase entry. *Biochem Biophys Res Commun* (2008) **369**:539–46. doi:10.1016/j.bbrc.2008.02.063
69. Dehennaut V, Slomianny MC, Page A, Vercoutter-Edouart AS, Jessus C, Michalski JC, et al. Identification of structural and functional O-linked N-acetylglucosamine-bearing proteins in *Xenopus laevis* oocyte. *Mol Cell Proteomics* (2008) **7**:2229–45. doi:10.1074/mcp.M700494-MCP200
70. Dehennaut V, Lefebvre T, Leroy Y, Vilain JP, Michalski JC, Bodart JF. Survey of O-GlcNAc level variations in *Xenopus laevis* from oogenesis to early development. *Glycoconj J* (2009) **26**:301–11. doi:10.1007/s10719-008-9166-0
71. Slawson C, Lakshmanan T, Knapp S, Hart GW. A mitotic GlcNAcylation/phosphorylation signaling complex alters the posttranslational state of the cytoskeletal protein vimentin. *Mol Biol Cell* (2008) **19**:4130–40. doi:10.1091/mbc.E07-11-1146
72. Baylin SB, Jones PA. A decade of exploring the cancer epigenome – biological and translational implications. *Nat Rev Cancer* (2011) **11**:726–34. doi:10.1038/nrc3130
73. Sakabe K, Hart GW. O-GlcNAc transferase regulates mitotic chromatin dynamics. *J Biol Chem* (2010) **285**:34460–8. doi:10.1074/jbc.M110.158170
74. Sakabe K, Wang Z, Hart GW. Beta-N-acetylglucosamine (O-GlcNAc) is part of the histone code. *Proc Natl Acad Sci U S A* (2010) **107**:19915–20. doi:10.1073/pnas.1009023107
75. Fong JJ, Nguyen BL, Bridger R, Medrano EE, Wells L, Pan S, et al. Beta-N-acetylglucosamine (O-GlcNAc) is a novel regulator of mitosis-specific phosphorylations on histone H3. *J Biol Chem* (2012) **287**:12195–203. doi:10.1074/jbc.M111.315804
76. Berger SL. The complex language of chromatin regulation during transcription. *Nature* (2007) **447**:407–12. doi:10.1038/nature05915
77. Hanover JA, Krause MW, Love DC. Bittersweet memories: linking metabolism to epigenetics through O-GlcNAcylation. *Nat Rev Mol Cell Biol* (2012) **13**:312–21. doi:10.1038/nrm3334
78. Fujiki R, Hashiba W, Sekine H, Yokoyama A, Chikanishi T, Ito S, et al. GlcNAcylation of histone H2B facilitates its monoubiquitination. *Nature* (2011) **480**:557–60.
79. Crosio C, Fimia GM, Loury R, Kimura M, Okano Y, Zhou H, et al. Mitotic phosphorylation of histone H3: spatio-temporal regulation by mammalian Aurora kinases. *Mol Cell Biol* (2002) **22**:874–85. doi:10.1128/MCB.22.3.874-885.2002
80. Goto H, Yasui Y, Nigg EA, Inagaki M. Aurora-B phosphorylates histone H3 at serine28 with regard to the mitotic chromosome condensation. *Genes Cells* (2002) **7**:11–7. doi:10.1046/j.1356-9597.2001.00498.x
81. Murnion ME, Adams RR, Callister DM, Allis CD, Earnshaw WC, Swedlow JR. Chromatin-associated protein phosphatase 1 regulates aurora-B and histone H3 phosphorylation. *J Biol Chem* (2001) **276**:26656–65. doi:10.1074/jbc.M102288200
82. Fujiki R, Chikanishi T, Hashiba W, Ito H, Takada I, Roeder RG, et al. GlcNAcylation of a histone methyltransferase in retinoic-acid-induced granulopoiesis. *Nature* (2009) **459**:455–9. doi:10.1038/nature07954
83. Mariappa D, Pathak S, van Aalten DM. A sweet TET-a-tet-synergy of TET proteins and O-GlcNAc transferase in transcription. *EMBO J* (2013) **32**:612–3. doi:10.1038/emboj.2013.26
84. Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, et al. Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* (2011) **333**:1300–3. doi:10.1126/science.1210597
85. Chen Q, Chen Y, Bian C, Fujiki R, Yu X. TET2 promotes histone O-GlcNAcylation during gene transcription. *Nature* (2013) **493**:561–4. doi:10.1038/nature11742
86. Vella P, Scelfo A, Jammula S, Chiacchiera E, Williams K, Cuomo A, et al. Tet proteins connect the O-linked N-acetylglucosamine transferase Ogt to chromatin in embryonic stem cells. *Mol Cell* (2013) **49**:645–56. doi:10.1016/j.molcel.2012.12.019
87. Deplus R, Delatte B, Schwinn MK, Defrance M, Mendez J, Murphy N, et al. TET2 and TET3 regulate GlcNAcylation and H3K4 methylation through OGT and SET1/COMPASS. *EMBO J* (2013) **32**:645–55. doi:10.1038/emboj.2012.357
88. Lee TI, Young RA. Transcriptional regulation and its misregulation in disease. *Cell* (2013) **152**:1237–51. doi:10.1016/j.cell.2013.02.014
89. Ozcan S, Andrali SS, Cantrell JE. Modulation of transcription factor function by O-GlcNAc modification. *Biochim Biophys Acta* (2010) **1799**:353–64. doi:10.1016/j.bbarm.2010.02.005
90. Meyer N, Penn LZ. Reflecting on 25 years with MYC. *Nat Rev Cancer* (2008) **8**:976–90. doi:10.1038/nrc2231
91. Gregory MA, Qi Y, Hann SR. Phosphorylation by glycogen synthase kinase-3 controls c-Myc proteolysis and subnuclear localization. *J Biol Chem* (2012) **278**:51606–12. doi:10.1074/jbc.M310722200
92. Chou TY, Hart GW, Dang CV. c-Myc is glycosylated at threonine 58, a known phosphorylation site and a mutational hot spot in lymphomas. *J Biol Chem* (1995) **270**:18961–5. doi:10.1074/jbc.270.32.18961
93. Ikonen HM, Minner S, Guldvik IJ, Sandmann MJ, Tsourlakis MC, Berge V, et al. O-GlcNAc transferase integrates metabolic pathways to regulate the stability of c-MYC in human prostate cancer. *Cancer Res* (2013). doi:10.1158/0008-5472.CAN-13-0549
94. Soussi T, Wiman KG. Shaping genetic alterations in human cancer: the p53 mutation paradigm. *Cancer Cell* (2007) **12**:303–12. doi:10.1016/j.ccr.2007.10.001
95. Dai C, Gu W. p53 Post-translational modification: deregulated in tumorigenesis. *Trends Mol Med* (2010) **16**:528–36. doi:10.1016/j.molmed.2010.09.002
96. Bech-Otschir D, Kraft R, Huang X, Henklein P, Kapelari B, Pollmann C, et al. COP9 signalosome-specific phosphorylation targets p53 to degradation by the ubiquitin system. *EMBO J* (2001) **20**:1630–9. doi:10.1093/emboj/20.7.1630
97. Yang WH, Kim JE, Nam HW, Ju JW, Kim HS, Kim YS, et al.

- Modification of p53 with O-linked N-acetylglucosamine regulates p53 activity and stability. *Nat Cell Biol* (2006) **8**:1074–83. doi:10.1038/ncb1470
98. Oren M, Rotter V. Mutant p53 gain-of-function in cancer. *Cold Spring Harb Perspect Biol* (2010) **2**:a001107. doi:10.1101/cshperspect.a001107
99. Kalo E, Kogan-Sakin I, Solomon H, Bar-Nathan E, Shay M, Shetzer Y, et al. Mutant p53R273H attenuates the expression of phase 2 detoxifying enzymes and promotes the survival of cells with high levels of reactive oxygen species. *J Cell Sci* (2012) **125**:5578–86. doi:10.1242/jcs.106815
100. Cooks T, Pateras IS, Tarcic O, Solomon H, Schetter AJ, Wilder S, et al. Mutant p53 prolongs NF-kappaB activation and promotes chronic inflammation and inflammation-associated colorectal cancer. *Cancer Cell* (2013) **23**:634–46. doi:10.1016/j.ccr.2013.03.022
101. Perkins ND. The diverse and complex roles of NF-kappaB subunits in cancer. *Nat Rev Cancer* (2012) **12**:121–32. doi:10.1038/nrc3204
102. Tornatore L, Thotakura AK, Bennett J, Moretti M, Franzoso G. The nuclear factor kappa B signaling pathway: integrating metabolism with inflammation. *Trends Cell Biol* (2012) **22**:557–66. doi:10.1016/j.tcb.2012.08.001
103. Yang WH, Park SY, Nam HW, Kim Do H, Kang JG, Kang ES, et al. NFkappaB activation is associated with its O-GlcNAcylation state under hyperglycemic conditions. *Proc Natl Acad Sci U S A* (2008) **105**:17345–50. doi:10.1073/pnas.0806198105
104. Ma Z, Vocadlo DJ, Vosseller K. Hyper-O-GlcNAcylation is anti-apoptotic and maintains constitutive NF-kappaB activity in pancreatic cancer cells. *J Biol Chem* (2012) **288**:15121–30. doi:10.1074/jbc.M113.470047
105. Kawachi K, Araki K, Tobiume K, Tanaka N. p53 Regulates glucose metabolism through an IKK-NF-kappaB pathway and inhibits cell transformation. *Nat Cell Biol* (2008) **10**:611–8. doi:10.1038/ncb1724
106. Kawachi K, Araki K, Tobiume K, Tanaka N. Loss of p53 enhances catalytic activity of IKKbeta through O-linked beta-N-acetyl glucosamine modification. *Proc Natl Acad Sci U S A* (2009) **106**:3431–6. doi:10.1073/pnas.0813210106
107. Dajee M, Lazarov M, Zhang JY, Cai T, Green CL, Russell AJ, et al. NF-kappaB blockade and oncogenic Ras trigger invasive human epidermal neoplasia. *Nature* (2003) **421**:639–43. doi:10.1038/nature01283
108. Greten FR, Eckmann L, Greten TF, Park JM, Li ZW, Egan LJ, et al. IKKbeta links inflammation and tumorigenesis in a mouse model of colitis-associated cancer. *Cell* (2004) **118**:285–96. doi:10.1016/j.cell.2004.07.013
109. Meylan E, Dooley AL, Feldser DM, Shen L, Turk E, Ouyang C, et al. Requirement for NF-kappaB signalling in a mouse model of lung adenocarcinoma. *Nature* (2009) **462**:104–7. doi:10.1038/nature08462
110. Mauro C, Leow SC, Anso E, Rocha S, Thotakura AK, Tornatore L, et al. NF-kappaB controls energy homeostasis and metabolic adaptation by upregulating mitochondrial respiration. *Nat Cell Biol* (2011) **13**:1272–9. doi:10.1038/ncb2324
111. Koo CY, Muir KW, Lam EW. FOXM1: from cancer initiation to progression and treatment. *Biochim Biophys Acta* (2012) **1819**:28–37. doi:10.1016/j.bbagr.2011.09.004
112. Caldwell SA, Jackson SR, Shahriari KS, Lynch TP, Sethi G, Walker S, et al. Nutrient sensor O-GlcNAc transferase regulates breast cancer tumorigenesis through targeting of the oncogenic transcription factor FoxM1. *Oncogene* (2010) **29**:2831–42. doi:10.1038/onc.2010.41
113. Valenta T, Hausmann G, Basler K. The many faces and functions of beta-catenin. *EMBO J* (2012) **31**:2714–36. doi:10.1038/emboj.2012.150
114. Olivier-Van Stichelen S, Guinez C, Mir AM, Perez-Cervera Y, Liu C, Michalski JC, et al. The hexosamine biosynthetic pathway and O-GlcNAcylation drive the expression of beta-catenin and cell proliferation. *Am J Physiol Endocrinol Metab* (2012) **302**:E417–24. doi:10.1152/ajpendo.00390.2011
115. Anagnostou SH, Shepherd PR. Glucose induces an autocrine activation of the Wnt/beta-catenin pathway in macrophage cell lines. *Biochem J* (2008) **416**:211–8. doi:10.1042/BJ20081426
116. Tai HC, Khidekel N, Ficarro SB, Peters EC, Hsieh-Wilson LC. Parallel identification of O-GlcNAc-modified proteins from cell lysates. *J Am Chem Soc* (2004) **126**:10500–1. doi:10.1021/ja047872b
117. Schlummer S, Vetter R, Kuder N, Henkel A, Chen YX, Li YM, et al. Influence of serine O-glycosylation or O-phosphorylation close to the vJun nuclear localisation sequence on nuclear import. *Chembiochem* (2006) **7**:88–97. doi:10.1002/cbic.200500212
118. Jiang MS, Hart GW. A subpopulation of estrogen receptors are modified by O-linked N-acetylglucosamine. *J Biol Chem* (1997) **272**:2421–8. doi:10.1074/jbc.272.4.2421
119. Chen YX, Du JT, Zhou LX, Liu XH, Zhao YF, Nakanishi H, et al. Alternative O-GlcNAcylation/O-phosphorylation of Ser16 induce different conformational disturbances to the N terminus of murine estrogen receptor beta. *Chem Biol* (2006) **13**:937–44. doi:10.1016/j.chembiol.2006.06.017
120. Bachmaier R, Aryee DN, Jug G, Kauer M, Kreppel M, Lee KA, et al. O-GlcNAcylation is involved in the transcriptional activity of EWS-FLI1 in Ewing's sarcoma. *Oncogene* (2009) **28**:1280–4. doi:10.1038/onc.2008.484
121. Park SY, Kim HS, Kim NH, Ji S, Cha SY, Kang JG, et al. Snail1 is stabilized by O-GlcNAc modification in hyperglycaemic condition. *EMBO J* (2010) **29**:3787–96. doi:10.1038/emboj.2010.254
122. Jin FZ, Yu C, Zhao DZ, Wu MJ, Yang Z. A correlation between altered O-GlcNAcylation, migration and with changes in E-cadherin levels in ovarian cancer cells. *Exp Cell Res* (2013) **319**:1482–90. doi:10.1016/j.yexcr.2013.03.013
123. Gialeli C, Theocharis AD, Karamanos NK. Roles of matrix metalloproteinases in cancer progression and their pharmacological targeting. *FEBS J* (2011) **278**:16–27. doi:10.1111/j.1742-4658.2010.07919.x
124. Yi W, Clark PM, Mason DE, Keenan MC, Hill C, Goddard WA III, et al. Phosphofructokinase 1 glycosylation regulates cell growth and metabolism. *Science* (2012) **337**:975–80. doi:10.1126/science.1222278
125. Guo K, Gan L, Zhang S, Cui FJ, Cun W, Li Y, et al. Translocation of HSP27 into liver cancer cell nucleus may be associated with phosphorylation and O-GlcNAc glycosylation. *Oncol Rep* (2012) **28**:494–500. doi:10.3892/or.2012.1844
126. Rambaruth ND, Greenwell P, Dwek MV. The lectin Helix pomatia agglutinin recognizes O-GlcNAc containing glycoproteins in human breast cancer. *Glycobiology* (2012) **22**:839–48. doi:10.1093/glycob/cws051
127. Dorfmüller HC, Borodkin VS, Blair DE, Pathak S, Navratilova I, van Aalten DM. Substrate and product analogues as human O-GlcNAc transferase inhibitors. *Amino Acids* (2011) **40**:781–92. doi:10.1007/s00726-010-0688-y

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 05 June 2013; paper pending published: 25 June 2013; accepted: 26 July 2013; published online: 12 August 2013.  
Citation: Fardini Y, Dehennaut V, Lefebvre T and Issad T (2013) O-GlcNAcylation: a new cancer hallmark? *Front. Endocrinol.* **4**:99. doi:10.3389/fendo.2013.00099

This article was submitted to *Frontiers in Molecular and Structural Endocrinology*, a specialty of *Frontiers in Endocrinology*. Copyright © 2013 Fardini, Dehennaut, Lefebvre and Issad. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.