



HAL
open science

Mediation role of cardiorespiratory fitness on the association between fatness and cardiometabolic risk in European adolescents: The HELENA study.

Carlos Cristi-Montero, Javier Courel-Ibáñez, Francisco B Ortega, Jose Castro-Piñero, Alba Santaliestra-Pasias, Angela Polito, Jeremy Vanhelst, Ascensión Marcos, Luis M Moreno, Jonatan R Ruiz

► To cite this version:

Carlos Cristi-Montero, Javier Courel-Ibáñez, Francisco B Ortega, Jose Castro-Piñero, Alba Santaliestra-Pasias, et al.. Mediation role of cardiorespiratory fitness on the association between fatness and cardiometabolic risk in European adolescents: The HELENA study.. *Journal of Sport and Health Science*, 2021, 10 (3), pp.360-367. 10.1016/j.jshs.2019.08.003 . hal-03231666

HAL Id: hal-03231666

<https://hal.univ-lille.fr/hal-03231666v1>

Submitted on 21 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Original article

Mediation role of cardiorespiratory fitness on the association between fatness and cardiometabolic risk in European adolescents: The HELENA study

Carlos Cristi-Montero ^{a,*}, Javier Courel-Ibáñez ^b, Francisco B. Ortega ^c, Jose Castro-Piñero ^d, Alba Santaliesra-Pasias ^{e,f}, Angela Polito ^g, Jérémy Vanhelst ^h, Ascensión Marcos ⁱ, Luis M. Moreno ^{e,f}, Jonatan R. Ruiz ^c, on behalf of the HELENA study group

^a IRyS Group, Physical Education School, Pontificia Universidad Católica de Valparaíso, Valparaíso 2530388, Chile

^b Department of Physical Activity and Sport, Faculty of Sport Sciences, University of Murcia, Murcia 30071, Spain

^c PROMoting FITness and Health through Physical Activity Research Group (PROFITH), Department of Physical Education and Sports, Faculty of Sport Sciences, University of Granada, Granada 18001, Spain

^d Department of Physical Education, Faculty of Education Sciences, University of Cádiz, Puerto Real 11003, Spain

^e Department of Health and Human Performance, School of Health Sciences, University of Zaragoza, Zaragoza 50001, Spain

^f Growth, Exercise, Nutrition and Development (GENUD) Research Group, Zaragoza 50001, Spain

^g National Institute for Food and Nutrition Research, Rome 80070, Italy

^h Lille Inflammation Research International Center, University of Lille, Lille 59000, France

ⁱ Immunonutrition Research Group, Department Metabolism and Nutrition, Institute of Food Science and Technology and Nutrition (ICTAN), Spanish National Research Council (CSIC), Madrid E-28040 Spain

Received 20 September 2018; revised 25 February 2019; accepted 9 June 2019

Available online 16 August 2019

2095-2546/© 2021 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Purpose: This study was aimed to analyze the mediation role of cardiorespiratory fitness (CRF) on the association between fatness and cardiometabolic risk scores (CMRs) in European adolescents.

Methods: A cross-sectional study was conducted in adolescents ($n = 525$; 46% boys; 14.1 ± 1.1 years old, mean \pm SD) from 10 European cities involved in the Healthy Lifestyle in Europe by Nutrition in Adolescence study. CRF was measured by means of the shuttle run test, while fatness measures included body mass index (BMI), waist to height ratio, and fat mass index estimated from skinfold thicknesses. A clustered CMRs was computed by summing the standardized values of homeostasis model assessment, systolic blood pressure, triglycerides, total cholesterol/high-density lipoprotein cholesterol ratio, and leptin.

Results: Linear regression models indicated that CRF acted as an important and partial mediator in the association between fatness and CMRs in 12–17-year-old adolescents (for BMI: coefficients of the indirect role $\beta = 0.058$ (95% confidence interval (95%CI): 0.023–0.101), Sobel test $z = 3.11$ (10.0% mediation); for waist to height ratio: $\beta = 4.279$ (95%CI: 2.242–7.059), $z = 3.86$ (11.5% mediation); and for fat mass index: $\beta = 0.060$ (95%CI: 0.020–0.106), $z = 2.85$ (9.4% mediation); all $p < 0.01$).

Conclusion: In adolescents, the association between fatness and CMRs could be partially decreased with improvements to fitness levels; therefore, CRF contribution both in the clinical field and public health could be important to consider and promote in adolescents independently of their fatness levels.

Keywords: Cardiovascular disease; Children; Fat mass; Fitness; Health; Physical activity

1. Introduction

Obesity in adolescents has increased dramatically in recent decades and has reached worldwide pandemic proportions.¹ Obese youths are likely to become obese adults and are at

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author.

E-mail address: carlos.cristi.montero@gmail.com (C. Cristi-Montero).

higher risk of developing cardiovascular diseases (CVD), dyslipidemia, and type 2 diabetes,^{2,3} which together constitute the main cause of death in adults.⁴ Both unhealthy dietary behaviors and physical inactivity⁵ seem to be the key determinants of obesity in youth; thus, early interventions for preventing the development of obesity-related CVD are needed not only for present health but also for future health.

To date, there has been no absolute clarity regarding the reverse or bidirectional relationship (cause–effect–response) between obesity and physical inactivity because the direction of causality cannot be inferred from cross-sectional associations.^{6,7} There is evidence from longitudinal studies that shows reciprocal associations and synergistic interactions among fatness with motor skill competence, physical activity, and cardiorespiratory fitness (CRF).^{8–10} In contrast, there are other studies that show that fatness leads to inactivity in children.^{6,7} These circumstances expose the ambiguity of the scientific literature in this matter and the need to investigate the interaction between fatness and fitness.

Of note is that physically inactive adolescents have lower CRF levels;^{11,12} and, in turn, this physiological marker is crucial for attenuating the association between fatness and increased cardiometabolic risk from childhood to adolescence.¹³ Hence, nowadays there is a special focus on improving CRF,¹⁴ especially in obese adolescents.¹⁵

An interesting approach to studying the role and importance of a variable over the association between others is mediation analysis, which can determine whether a predictor variable in an outcome is mediated through an intermediate variable (mediator).¹⁶ This innovative method has been applied in studies of children to examine the mediator role of obesity in the relationship between CRF and cardiometabolic risk¹⁷ and inflammation.¹⁸ The authors of these studies reported that, when including body mass index (BMI) as a mediator, the association between CRF and cardiometabolic risk disappears completely in girls and is considerably mitigated in boys.¹⁷ These results do not fully support the “fat-but-fit” paradigm, which suggests that a person with excess adiposity but with a high CRF (fat-fit phenotype) has a better cardiometabolic profile than a person with excess adiposity but with a low CRF.^{13,19} Accordingly, the role of fitness and fatness as mediators of cardiometabolic health still needs to be clarified, especially in children and adolescents.¹⁹

Based on the assumption that fatness leads to inactivity, the purpose of this study was to analyze the mediation role of CRF on the association between fatness markers and cardiometabolic risk scores (CMRs) clustering in European adolescents and to analyze the magnitude of CRF mediation in this relationship. We hypothesized that CRF would only partially mediate the relationship between fatness and CMRs. This investigation contributes to the field of public health by providing evidence about the bidirectional relationship between obesity and CRF.

2. Methods

The Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) cross-sectional study was designed to obtain reliable and comparable data on nutrition and health-related parameters from a sample of adolescents aged

12.5–17.5 years from 10 European cities in 9 countries. Data collection took place between 2006 and 2008. A sample of 3528 adolescents met the HELENA general inclusion criteria (age of 13.00–16.99 years, schooling in one of the 10 European cities, informed consent signed, had at least weight and height measured, and completed at least 75% of the other tests).²⁰ In the present study, 525 adolescents (54% girls) with valid data on the main variables and covariates were included in the analysis: cardiometabolic risk factors, including total cholesterol to high-density lipoprotein cholesterol ratio (TC/HDL-C), triglycerides (TG), homeostasis model assessment (HOMA), and systolic blood pressure (SBP); as well as leptin, CRF, age, sex, pubertal stage, and center (city involved in the study from each country). This subsample was representative to the total sample (assuming an error of 5%, a confidence interval (CI) of 95%, and 50% of heterogeneity). Moreover, there were no differences between the present subsample and the whole sample (boys and girls separately) in terms of age, body mass, height, and BMI (all $p > 0.05$).

The study was performed following the ethical guidelines of the 1964 Declaration of Helsinki (revision of Edinburgh 2000), Good Clinical Practice, and legislation regarding clinical research in humans in each of the participating countries. The protocol was approved by the Human Research Review Committees of the involved centers. Furthermore, all parents and guardians signed an informed consent form, and the adolescents agreed to participate in the study.²⁰

2.1. Physical examination

Adolescents participating in the study were barefoot and dressed in light clothing during anthropometric measurements. Body weight was measured to the nearest 0.1 kg with an electronic scale (Seca 861; Seca GmbH & Co., Hamburg, Germany), and body height was measured with a telescopic stadiometer (Seca 225; Seca GmbH & Co.) to the nearest 0.1 cm. BMI was calculated as body weight divided by the square of height (kg/m^2).²¹ Waist circumference (WC) was measured with a non-elastic tape (Seca 200; Seca GmbH & Co.) to the nearest 0.1 cm. Waist to height ratio (WHtR) was then calculated by dividing WC (cm) by height (cm). Bicipital, tricipital, subscapular, and suprailiac skinfold thicknesses on the left side of the body were measured with a Holtain Calliper (Holtain Ltd., Crymych, Wales, UK) to the nearest 0.2 mm. All the anthropometric measurements were taken 3 times and the mean was scored.²² Percentage of body fat was calculated using the equation of Slaughter et al.,²³ which has proven to be the most suitable equation for use with adolescents.²⁴ Fat mass (kg) was estimated by multiplying body fat percent by weight (kg) and dividing by 100, and fat-free mass (kg) was calculated as the difference between body weight and fat mass. Fat mass index (FMI) was determined by dividing fat mass (kg) by height squared (m^2). Identification of pubertal maturation (Stages I–V) was assessed by direct observation by a medical doctor, according to Tanner and Whitehouse.²⁵ SBP was measured (OMRON[®]M6, HEM 70001; Omron, Kyoto, Japan) following the recommendations for adolescent populations,²⁶

with participants seated in a separate quiet room for 10 min with their backs supported and feet on the ground. Two SPB readings were taken at 10-min intervals, and the lowest measure was recorded.

2.2. Blood samples

Blood samples were collected as previously described.²⁷ TG, TC, and HDL-C were measured using enzymatic methods (Dade Behring, Schwalbach, Germany). All blood parameters were measured after an overnight fast. HOMA index calculation was used as a measure of insulin resistance²⁸ using the formula $\text{HOMA-index} = (\text{insulin } (\mu\text{UI/mL}) \times \text{glucose } (\text{mg/dL})) / 405$. The ratio of TC/HDL-C was also calculated. Leptin concentrations were measured by the RayBio (RayBiotech Inc., GA, USA) Human Leptin enzyme-linked immunosorbent assay. Leptin assay sensitivity was set at less than 6 pg/mL, with intraclass and interclass coefficients of variation of less than 10% and less than 12%, respectively.²⁹

2.3. CRF

CRF was measured by the progressive 20-m shuttle run test. The test has high validity and reliability in adolescents (test–retest reliability coefficients were 0.89; standard error of estimation was 5.9 mL/kg/min).^{30,31} In the shuttle run test, participating adolescents ran between 2 lines 20 m apart, while keeping the pace with audio signals emitted from a pre-recorded CD with an initial speed of 8.5 km/h and increasing by 0.5 km/h every minute (1 min equals 1 stage). The test ended either when the adolescent failed to reach the end line concurrent with the audio signals on 2 consecutive occasions or when he or she stopped owing to fatigue. The last stage completed (precision of 0.5 stages) was used to calculate the maximal oxygen consumption from the equation developed by Léger et al.³⁰

2.4. CMRs

A summative cardiometabolic risk index that includes several factors has proven to be a better marker of cardiovascular health in children than an index that only includes a single risk factor.²⁹ CMRs were created from the sum of SBP, TG, TC/HDL-C ratio, HOMA, and leptin z-scores. The standardized value (z-scores) of each variable was calculated as follows: $(\text{value} - \text{mean}) / \text{SD}$, separately for boys and girls, and for each 1-year age group. Lower values are indicative of a better profile.³² This risk profile has been used previously in a cross-sectional study of 1732 randomly selected 9-year-old and 15-year-old school children from 3 European countries and in 2015.³³ Andersen et al.³² confirmed that the composite risk score improved substantially in a sample of 15,794 youths aged 6–18 years that included the HOMA index rather than fasting glucose, leptin, the sum-of-4-skinfolds (instead of BMI or WC), and CRF. Both the sum-of-4-skinfolds and CRF were removed from this CMRs because these were part of the mediation analysis and could have generated a methodological bias.

2.5. Statistical analysis

The normality of the distribution of the variables was tested using both statistical (Kolmogorov–Smirnov test) and graphical methods (normal probability plot). Adolescents' characteristics are presented as the mean \pm SD and the frequency (%) for continuous and categorical variables, respectively. The Student *t* test and the Pearson's χ^2 test were used to test sex differences in relation to adolescents' CRF, fatness (BMI, WHtR, and FMI), and cardiometabolic biomarkers (TC/HDL-C, TG, HOMA, SBP, and leptin) and pubertal stage. Partial correlation coefficients (*r*), adjusted for age, sex, pubertal stage, and center, were used as a preliminary analysis to examine the associations between CRF, cardiometabolic, and fatness biomarkers.

Multicollinearity was tested before completing the mediation analysis through tolerance value and variance inflation factors.^{34,35} We analyzed the mediation role of CRF on the association of 3 fatness markers (BMI, WHtR, and FMI) with CMRs through bootstrapped (10,000 samples) linear regression analyses³⁶ using the PROCESS SPSS script,¹⁶ adjusted for age, sex, pubertal stage, fat-free mass, and center. Each model included 3 equations.

Equation 1 regressed the mediator (CRF) on the independent variable (BMI, WHtR, and FMI). Equation 2 regressed the dependent variable (CMRs) on the independent variable (BMI, WHtR, and FMI). Equation 3 regressed the dependent variable (CMRs) on both the independent (BMI, WHtR, and FMI) and the mediator variable (CRF), and Equation 3' took into account the indirect (mediating) role. We calculated the percentage of the total contribution that is accounted for by mediation using the standardized coefficients of the Equation 1 \times Equation 2 / Equation 3. A significant "indirect role" (mediation) was established when (a) the independent variable was significantly related to the mediator, (b) the independent variable was significantly related to the dependent variable, (c) the mediator was significantly related to the dependent variable, and (d) the association between the independent and dependent variable ("direct role") was attenuated when the mediator was included in the regression model.³⁷ The Sobel test was used to test the hypothesis that the indirect role was equal to 0.³⁸ Point estimates and 95% confidence intervals (95% CIs) were estimated for the indirect role. Complete mediation was established when the independent variable (fatness) was not associated with the dependent variable (CMRs) after mediator (CRF) has been controlled, making path Equation 3' zero. Partial mediation is when the path from fatness to CMRs is reduced in absolute size but is still different from 0 when the mediator is introduced. The percentage of mediation was calculated as $1 - (\text{Equation } 3' / \text{Equation } 3)$ where Equation 3 is the role for the independent variable in predicting the main outcome and Equation 3' is the role of the independent variable in predicting the dependent variable with the mediator variable. Analyses were completed using the IBM SPSS (Version 21.0; IBM Corp., Armonk, NY, USA). The level of significance was set at $p < 0.05$.

3. Results

Table 1 shows the characteristics of the sample. The partial correlations coefficients between the fitness variables, CRF, and cardiometabolic biomarkers are presented in Table 2. There was no difference in the CMRs between sex ($p = 0.992$) when both boys and girls were grouped. CRF was negatively associated with all the study measures of fatness as well as with cardiometabolic biomarkers ($p < 0.05$), except for SBP. BMI, WHtR, and FMI were associated positively with all cardiometabolic biomarkers ($p < 0.05$). We observed no multicollinearity. The tolerance value ranged between 0.829 and 0.888 for collinearity tolerance and the variance inflation factor ranged between 1127 and 1206 for collinearity.

The results from mediation analysis are shown in Fig. 1. Overall, fatness (BMI, WHtR, and FMI) was positively associated with CMRs. Mediation analysis including CRF revealed that the association between fatness and CMRs was mediated

via CRF. The role of this mediation adjusted for potential confounders accounted for around 10%. More precisely, CRF mediates 10.0% of CMRs for BMI ($z = 3.11$; indirect role = 0.058 (95%CI: 0.023–0.101); $p = 0.001$), 11.5% for WHtR ($z = 3.86$; indirect role = 4.279 (95%CI: 2.242–7.059); $p < 0.001$), and 9.4% for FMI ($z = 2.85$; indirect role = 0.060 (95%CI: 0.020–0.106); $p = 0.004$) parameters.

4. Discussion

The main findings of the current study indicate that CRF acts as a partial mediator in the association between 3 different fitness variables and clustered CMRs in European adolescents. Therefore, although it is true that CRF is an important health biomarker, it does not seem to completely counteract the negative role of fatness on cardiometabolic health. To our knowledge, this is one of the few studies that have used CRF as a mediator,^{39,40} and the first to establish the mediation role of

Table 1
Characteristics of adolescents.

	Overall ($n = 525$)	Boys ($n = 241$)	Girls ($n = 284$)
Age (year)	14.1 ± 1.1	14.2 ± 1.2	14.1 ± 1.1
Body mass (kg)	56.9 ± 12.3	59.0 ± 13.3	55.2 ± 11.2*
Height (cm)	164.7 ± 9.7	168.9 ± 10.2	161.1 ± 7.5*
BMI (kg/m ²)	20.8 ± 3.4	20.5 ± 3.3	21.1 ± 3.5*
Pubertal stage I/II/III/IV/V (%)	0.8/6.8/21.3/37.4/29.0	1.7/9.1/20.9/34.1/30.3	0.0/5.1/21.6/39.9/28.0
Fat mass (kg)	13.5 ± 8.2	11.6 ± 9.0	15.1 ± 7.2*
Fat-free mass (kg)	43.4 ± 8.1	47.4 ± 8.9	40.1 ± 5.5*
Waist circumference (cm)	71.1 ± 8.2	72.3 ± 8.1	70.0 ± 8.2*
WHtR	0.43 ± 0.04	0.43 ± 0.04	0.43 ± 0.04
CRF (mL/kg/min)	41.1 ± 7.8	46.0 ± 7.2	37.0 ± 5.6*
FMI (kg/m ²)	4.9 ± 2.9	4.0 ± 3.0	5.7 ± 2.6*
TC (mg/dL)	161.0 ± 26.9	153.1 ± 24.0	167.7 ± 27.4*
HDL-C (mg/dL)	55.6 ± 10.5	53.4 ± 9.7	57.5 ± 10.9*
Triglycerides (mg/dL)	69.1 ± 35.4	62.4 ± 29.4	74.8 ± 38.9*
HOMA	2.4 ± 2.0	2.3 ± 2.4	2.4 ± 1.6
SBP (mmHg)	117.0 ± 13.1	120.9 ± 13.8	113.8 ± 11.6*
Leptin (ng/dL)	19.3 ± 22.0	9.3 ± 14.6	27.8 ± 23.5*

Notes: Values are mean ± SD or frequency (%). Independent two-tailed t tests or χ^2 tests were applied to compare unadjusted means by sex.

* $p < 0.05$, compared with boys.

Abbreviations: BMI = body mass index; CRF = cardiorespiratory fitness; FMI = fat mass index; HDL-C = high-density lipoprotein cholesterol; HOMA = homeostasis model assessment; SBP = systolic blood pressure; TC = total cholesterol; WHtR = waist to height ratio.

Table 2
Partial correlations coefficients (r) among fatness, CRF, and cardiometabolic biomarkers.

	CRF	BMI	WHtR	FMI	SBP	TC/HDL-C	TG	HOMA	Leptin
CRF	—	—	—	—	—	—	—	—	—
BMI	-0.318*	—	—	—	—	—	—	—	—
WHtR	-0.323*	0.892*	—	—	—	—	—	—	—
FMI	-0.409*	0.909*	0.852*	—	—	—	—	—	—
SBP	-0.030	0.356*	0.339*	0.288*	—	—	—	—	—
TC/HDL-C	-0.161*	0.233*	0.253*	0.219*	0.026	—	—	—	—
TG	-0.148*	0.196*	0.218*	0.216*	0.011	0.449*	—	—	—
HOMA	-0.240*	0.336*	0.313*	0.344*	0.170*	0.141*	0.381*	—	—
Leptin	-0.322*	0.635*	0.632*	0.683*	0.185*	0.149*	0.227*	0.320*	—

Note: Model was adjusted for age, sex, pubertal stage, and center.

* $p < 0.05$, significant association.

Abbreviations: BMI = body mass index; CRF = cardiorespiratory fitness; FMI = fat mass index; HOMA = homeostasis model assessment; SBP = systolic blood pressure; TC/HDL-C = total cholesterol to high-density lipoprotein cholesterol ratio; TG = triglycerides; WHtR = waist to height ratio.




Fig. 1. Mediation analysis. Contribution of fitness on CMRs through CRF, adjusting for potential confounders (age, sex, pubertal stage, center, and fat-free mass). BMI = body mass index; CMRs = cardiometabolic risk scores; CRF = cardiorespiratory fitness; FMI = fat mass index; WHtR = waist to height ratio.

CRF on the relationship between fatness and cardiometabolic risk in European adolescents.

The evidence shows a close relationship among fatness, low CRF, and cardiometabolic health in youths.^{41–43} Thus, CRF and obesity as predictors of CVD should be monitored to identify children and adolescents with potential CVD risk.^{44,45} Despite the importance of both markers on cardiometabolic health, mediation analysis studies have frequently used fatness as a mediator,^{17,18,46} assuming a possible unique association way (physical inactivity as causal of fatness).

However, there is also literature showing otherwise.^{6,7} Thus, looking at the 2 sides of the coin seems to be adequate for achieving a broader and objective understanding of this problem to improve the effectiveness of intervention programs in this population.

The results of the present mediation analysis show that CRF is capable of partially attenuating the association between fatness and a cluster of cardiometabolic risk factors in adolescents. In this sense, the magnitude of the mediation of CRF could support—up to a point—the fat-but-fit hypothesis,^{19,47}

because a high level of CRF would not fully protect the independent action of fatness (BMI, WHtR, and FMI) on cardiometabolic risk in adolescents. However, the independent and beneficial role of high CRF may indeed offset more of the fatness risk than indicated by the mediator analysis—just by mechanisms not related to fatness. It should be noted that this mediation role is around 10% for the 3 fatness variables studied in this approach, which gives it an important role both in public health strategies and for clinical purposes.

This outcome is in line with previous prospective research reporting that in schoolchildren (aged 7–11 years) the association of CRF with change in CVD risk factor levels after adjustment for adiposity (total body fat, BMI, and WC) was less strong and concluding that the association of CRF with CVD risk factors was largely explained by adiposity.⁴⁸ Another study of schoolchildren (aged 10–14 years) showed that children with low fitness levels had increased odds of presenting both individual and clustered cardiometabolic risk factors, but these associations no longer remained after adjusting for abdominal adiposity.⁴⁹ The authors suggested that abdominal adiposity may be a more important determinant of adverse cardiometabolic health in this age group,⁴⁹ and especially in obese children.^{15,50} Notwithstanding the foregoing, childhood CRF seems to decrease the long-term (20-year follow-up of 1792 adults) cardiometabolic risks associated with childhood obesity.⁵¹ Therefore, more studies exploring the symbiotic interaction between fatness and CRF on cardiometabolic health are warranted to establish insight into the bidirectional causality between these variables.^{6,7}

The physiological interpretation of these findings is complex because fatness and CRF are separately and independently associated with cardiometabolic risk factors,⁵² possibly exerting their effects through different causal pathways.^{53,54} For example, obesity may result in the development of a chronic, low-grade inflammatory state caused by an increased expression of pro-inflammatory adipokines and diminished expression of anti-inflammatory adipokines.⁵⁴ Moreover, obesity has been associated with elevated oxidative stress, which can lead to endothelial dysfunction.⁵⁵ It has also been related to a decrease in oxygen respiration in skeletal muscle associated with an impairment of mitochondrial function.⁵⁶

Each condition mentioned (chronic low-grade inflammatory state, diminished antioxidant capacity, and decrease in oxygen respiration in skeletal muscle) can be improved or even reversed, to some extent, by adequate levels of physical activity and exercise^{54–56} and by enhancing CRF.^{19,47,57} A meta-analysis showed that high-intensity interval training seems to be more effective for improving cardiometabolic risk and CRF than other forms of exercise in overweight and obese youth.⁵⁸ High-intensity activities, particularly among adolescents, seems to be more effective in reducing a clustered metabolic syndrome for both fit and unfit individuals.³⁹ However, our results contribute to the existing evidence by suggesting that if younger individuals maintain adequate CRF levels, this can play a protective role that partially counteracts the harmful role of obesity.^{6,7,48–50}

This work has some important strengths, such as the size of the European sample used (multicentric study) and the use of high-quality harmonization methods. It also seems to be the first study that has used CRF as a mediator to analyze CRF's role in relation to a cluster of cardiometabolic risk factors in adolescents. However, this study also has certain limitations. The present mediation analysis should be interpreted as exploratory given the cross-sectional nature of our study; thus, it does not allow conclusions about the causality relationships to be drawn. Future longitudinal and experimental studies are needed to confirm our results.

Although it is true that CRF was measured with a validated and reliable test, this variable depends heavily on body weight. This situation could lead to underestimations of CRF values in people with overweight and obesity⁵⁹ and underestimate the benefit of the fat-but-fit hypothesis.⁶⁰ Moreover, the use of clustered CMRs is specific to the study sample, and each factor is equally weighted in predicting future disease. Finally, the sample used in this study was made up of relatively healthy youths, which could hamper the sensitivity in detecting associations.

5. Conclusion

CRF in adolescents seems to act as a significant and partial mediator of the relationship between fatness and CMRs, which supports, to a certain extent, the fat-but-fit paradigm. These results suggest that the independent role of CRF is associated with other mechanisms that are not directly linked to fatness. However, these findings also indicate that maintaining adequate CRF levels among adolescents can counteract the development of harmful obesity-related CVDs and thus is an important public health strategy.

Acknowledgments

We thank the adolescents who participated in the study and their parents and teachers for their collaboration. We also acknowledge the HELENA study members involved in fieldwork for their efforts. The HELENA project was supported by the European Community 6th Framework Programme for Research and Technological Development (contract FOOD-CT-2005-007034). The data for this study were gathered under the aegis of the HELENA project, and further analysis was additionally supported by the Spanish Ministry of Economy and Competitiveness (Grants RYC-2010-05957 and RYC-2011-09011), the Spanish Ministry of Health: Maternal, Child Health and Development Network (Grants RD08/0072 and RD16/0022), the Fondo Europeo de Desarrollo Regional (MICINN-FEDER), and the University of Granada, Plan Propio de Investigación 2016, Excellence Actions: Units of Excellence; Unit of Excellence on Exercise and Health (UCEES). The content of this article reflects the authors' views alone, and the European Community is not liable for any use that may be made of the information contained herein.

Authors' contributions

CCM and JRR had full access to all of the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis; CCM and JCI conceived and designed the study and drafted the manuscript; FBO, JCP, ASP, AP, JV, AM, LMM, and JRR carried out critical revisions of the manuscript for important intellectual content. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

References

- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study 2013. *The Lancet* 2014;**384**:766–81.
- Arellano-Ruiz P, García-Hermoso A, Martínez-Vizcaíno V, Salcedo-Aguilar F, Garrido-Miguel M, Solera-Martínez M. Trends in cardiometabolic parameters among Spanish children from 2006 to 2010: The Cuenca study. *Am J Hum Biol* 2017;**29**:e22970. doi:10.1002/ajhb.22970.
- Ajala O, Mold F, Boughton C, Cooke D, Whyte M. Childhood predictors of cardiovascular disease in adulthood. A systematic review and meta-analysis. *Obes Rev* 2017;**18**:1061–70.
- World Health Organization. *Global status report on noncommunicable diseases*. Geneva: World Health Organization; 2014.
- Vilallonga R, Moreno Villares JM, Yeste Fernández D, et al. Initial approach to childhood obesity in Spain. A multisociety expert panel assessment. *Obes Surg* 2017;**27**:997–1006.
- Metcalf BS, Hosking J, Jeffery AN, Voss LD, Henley W, Wilkin TJ. Fatness leads to inactivity, but inactivity does not lead to fatness: A longitudinal study in children (EarlyBird 45). *Arch Dis Child* 2011;**96**:942–7.
- Hjorth MF, Chaput JP, Ritz C, et al. Fatness predicts decreased physical activity and increased sedentary time, but not vice versa: Support from a longitudinal study in 8- to 11-year-old children. *Int J Obes* 2014;**38**:959–65.
- Lima RA, Pfeiffer KA, Bugge A, Møller NC, Andersen LB, Stodden DF. Motor competence and cardiorespiratory fitness have greater influence on body fatness than physical activity across time. *Scand J Med Sci Sport* 2017;**27**:1638–47.
- Lima RA, Bugge A, Ersbøll AK, Stodden DF, Andersen LB. The longitudinal relationship between motor competence and measures of fatness and fitness from childhood into adolescence. *J Pediatr (Rio J)* 2019;**95**:482–8.
- Stodden DF, Goodway JD, Langendorfer SJ, et al. A developmental perspective on the role of motor skill competence in physical activity: An emergent relationship. *Quest* 2008;**60**:290–306.
- Santos R, Mota J, Okely AD, et al. The independent associations of sedentary behaviour and physical activity on cardiorespiratory fitness. *Br J Sports Med* 2014;**48**:1508–12.
- Lagestad P, Mehui I. The importance of adolescents' participation in organized sport according to VO_{2peak} : A longitudinal study. *Res Q Exerc Sport* 2018;**89**:143–52.
- Brouwer SI, Stolk RP, Liem ET, Lemmink KA, Corpeleijn E. The role of fitness in the association between fatness and cardiometabolic risk from childhood to adolescence. *Pediatr Diabetes* 2013;**14**:57–65.
- Ross R, Blair SN, Arena R, et al. Importance of assessing cardiorespiratory fitness in clinical practice: A case for fitness as a clinical vital sign: A scientific statement from the American Heart Association. *Circulation* 2016;**134**:e653–99.
- Cummings DM, DuBose KD, Imai S, Collier DN. Fitness versus fatness and insulin resistance in U.S. Adolescents. *J Obes* 2010;**2010**: 195729. doi:10.1155/2010/195729.
- Hayes A. *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. New York, NY: Guilford Press; 2013.
- Diez-Fernández A, Sánchez-López M, Mora-Rodríguez R, Notario-Pacheco B, Torrijos-Niño C, Martínez-Vizcaíno V. Obesity as a mediator of the influence of cardiorespiratory fitness on cardiometabolic risk: A mediation analysis. *Diabetes Care* 2014;**37**:855–62.
- García-Hermoso A, Agostinis-Sobrinho C, Mota J, Santos RM, Correa-Bautista JE, Ramírez-Vélez R. Adiposity as a full mediator of the influence of cardiorespiratory fitness and inflammation in schoolchildren: The FUPRECOL study. *Nutr Metab Cardiovasc Dis* 2017;**27**:525–33.
- Ortega FB, Ruiz JR, Labayen I, Lavie CJ, Blair SN. The fat but fit paradox: what we know and don't know about it. *Br J Sports Med* 2017;**52**:151–3.
- Moreno LA, De Henauw S, González-Gross M, et al. Design and implementation of the Healthy Lifestyle in Europe by Nutrition in Adolescence cross-sectional study. *Int J Obes* 2008;**32**(Suppl. 5):S4–11.
- Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* 2012;**7**:284–94.
- Nagy E, Vicente-Rodríguez G, Manios Y, et al. Harmonization process and reliability assessment of anthropometric measurements in a multicenter study in adolescents. *Int J Obes* 2008;**32**(Suppl. 5):S58–65.
- Slaughter MH, Lohman TG, Boileau RA, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol* 1988;**60**:709–23.
- Rodríguez G, Moreno LA, Blay MG, et al. Body fat measurement in adolescents: comparison of skinfold thickness equations with dual-energy X-ray absorptiometry. *Eur J Clin Nutr* 2005;**59**:1158–66.
- Tanner JM, Whitehouse RH. Clinical longitudinal standards for height, weight, height velocity, weight velocity, and stages of puberty. *Arch Dis Child* 1976;**51**:170–9.
- National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004;**114**:555–76.
- González-Gross M, Breidenassel C, Gómez-Martínez S, et al. Sampling and processing of fresh blood samples within a European multicenter nutritional study: Evaluation of biomarker stability during transport and storage. *Int J Obes* 2008;**32**(Suppl. 5):S66–75.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: Insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;**28**:412–9.
- Labayen I, Ruiz J, Ortega F, et al. Association between the FTO rs9939609 polymorphism and leptin in European adolescents: A possible link with energy balance control. The HELENA study. *Int J Obes* 2010;**35**:66–71.
- Léger LA, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci* 1988;**6**:93–101.
- Olds T, Tomkinson G, Léger L, Cazorla G. Worldwide variation in the performance of children and adolescents: An analysis of 109 studies of the 20-m shuttle run test in 37 countries. *J Sports Sci* 2006;**24**:1025–38.
- Andersen LB, Harro M, Sardinha LB, et al. Physical activity and clustered cardiovascular risk in children: A cross-sectional study (The European Youth Heart Study). *The Lancet* 2006;**368**:299–304.
- Andersen LB, Laursen JB, Brønd JC, et al. A new approach to define and diagnose cardiometabolic disorder in children. *J Diabetes Res* 2015;**2015**: 539835. doi:10.1155/2015/539835.
- Menard S. *Applied logistic regression analysis: Sage university series on quantitative applications in the social sciences*. California, CA: Sage Publications Inc; 2002.
- Myers RH. *Classical and modern regression with applications*. Boston, MA: Duxbury Press; 1990.
- Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 2008;**40**:879–91.

37. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986;**51**:1173–82.
38. Sobel ME. Asymptotic confidence intervals for indirect effects in structural equation models. *Sociol Methodol* 1982;**13**:290–312.
39. Segura-Jiménez V, Parrilla-Moreno F, Fernández-Santos JR, et al. Physical fitness as a mediator between objectively measured physical activity and clustered metabolic syndrome in children and adolescents: The UP&DOWN study. *Nutr Metab Cardiovasc Dis* 2016;**26**:1011–9.
40. Muntaner-Mas A, Pere P, Vidal-Conti J, Esteban-Cornejo I. A mediation analysis on the relationship of physical fitness components, obesity, and academic performance in children. *J Pediatr* 2018;**198**:90–7.
41. Steele RM, Brage S, Corder K, Wareham NJ, Ekelund U. Physical activity, cardiorespiratory fitness, and the metabolic syndrome in youth. *J Adolesc* 2008;**105**:342–51.
42. Pozuelo-Carrascosa DP, Sánchez-López M, Cavero-Redondo I, Torres-Costoso A, Bermejo-Cantarero A, Martínez-Vizcaíno V. Obesity as a mediator between cardiorespiratory fitness and blood pressure in pre-schoolers. *J Pediatr* 2017;**182**:114–9.
43. Countryman AJ, Saab PG, Llabre MM, Penedo FJ, McCalla JR, Schneiderman N. Cardiometabolic risk in adolescents: Associations with physical activity, fitness, and sleep. *Ann Behav Med* 2013;**45**:121–31.
44. Castro-Piñero J, Perez-Bey A, Segura-Jiménez V, et al. Cardiorespiratory fitness cutoff points for early detection of present and future cardiovascular risk in children. *Mayo Clin Proc* 2017;**92**:1753–62.
45. Seo YG, Choi MK, Kang JH, et al. Cardiovascular disease risk factor clustering in children and adolescents: A prospective cohort study. *Arch Dis Child* 2018;**103**:968–73.
46. Díez-Fernández A, Sánchez-López M, Gullías-González R, et al. BMI as a mediator of the relationship between muscular fitness and cardiometabolic risk in children: A mediation analysis. *PLoS One* 2015;**10**:e0116506. doi:10.1371/journal.pone.0116506.
47. Ortega FB, Lavie CJ, Blair SN. Obesity and cardiovascular disease. *Circ Res* 2016;**118**:1752–70.
48. Klakk H, Grøntved A, Møller NC, Heidemann M, Andersen LB, Wedderkopp N. Prospective association of adiposity and cardiorespiratory fitness with cardiovascular risk factors in healthy children. *Scand J Med Sci Sport* 2014;**24**:e275–82.
49. Bailey DP, Savory LA, Denton SJ, Kerr CJ. The association between cardiorespiratory fitness and cardiometabolic risk in children is mediated by abdominal adiposity: The HAPPY study. *J Phys Act Heal* 2015;**12**:1148–52.
50. Nyström CD, Henriksson P, Martínez-Vizcaíno V, et al. Does cardiorespiratory fitness attenuate the adverse effects of severe/morbid obesity on cardiometabolic risk and insulin resistance in children? A pooled analysis. *Diabetes Care* 2017;**40**:1580–7.
51. Schmidt MD, Magnussen CG, Rees E, Dwyer T, Venn AJ. Childhood fitness reduces the long-term cardiometabolic risks associated with childhood obesity. *Int J Obes* 2016;**40**:1134–40.
52. LaMonte MJ, Blair SN. Physical activity, cardiorespiratory fitness, and adiposity: contributions to disease risk. *Curr Opin Clin Nutr Metab Care* 2006;**9**:540–6.
53. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: Skeletal muscle as a secretory organ. *Nat Rev Endocrinol* 2012;**8**:457–65.
54. Nakamura K, Fuster JJ, Walsh K. Adipokines: A link between obesity and cardiovascular disease. *J Cardiol* 2014;**63**:250–9.
55. Huang CJ, McAllister MJ, Slusher AL, Webb HE, Mock JT, Acevedo EO. Obesity-related oxidative stress: The impact of physical activity and diet manipulation. *Sport Med Open* 2015;**1**:32. doi:10.1186/s40798-015-0031-y.
56. Safdar A, Khrapko K, Flynn JM, et al. Exercise-induced mitochondrial p53 repairs mtDNA mutations in mutator mice. *Skelet Muscle* 2016;**6**:7. doi:10.1186/s13395-016-0075-9.
57. Ruiz JR, Huybrechts I, Cuenca-Garcia M, et al. Cardiorespiratory fitness and ideal cardiovascular health in European adolescents. *Heart* 2015;**101**:766–73.
58. García-Hermoso A, Cerrillo-Urbina AJ, Herrera-Valenzuela T, Cristi-Montero C, Saavedra JM, Martínez-Vizcaíno V. Is high-intensity interval training more effective on improving cardiometabolic risk and aerobic capacity than other forms of exercise in overweight and obese youth? A meta-analysis. *Obes Rev* 2016;**17**:531–40.
59. Krachler B, Savonen K, Komulainen P, Hassinen M, Lakka TA, Rauramaa R. Cardiopulmonary fitness is a function of lean mass, not total body weight: The DRs EXTRA study. *Eur J Prev Cardiol* 2015;**22**:1171–9.
60. Silva DR, Werneck AO, Collings PJ, et al. Cardiorespiratory fitness effect may be under-estimated in “fat but fit” hypothesis studies. *Ann Hum Biol* 2017;**44**:237–42.