

The "authentic subjective experience" of memory in Alzheimer's disease

Mohamad El Haj, Ahmed A. Moustafa, Jean Roche, Florence Pasquier, Dimitrios Kapogiannis, Karim Gallouj, Pascal Antoine

► **To cite this version:**

Mohamad El Haj, Ahmed A. Moustafa, Jean Roche, Florence Pasquier, Dimitrios Kapogiannis, et al.. The "authentic subjective experience" of memory in Alzheimer's disease. *Translational Neuroscience*, 2020, *Translational neuroscience*, 11, pp.201-207. 10.1515/tnsci-2020-0123 . hal-03367625

HAL Id: hal-03367625

<https://hal.univ-lille.fr/hal-03367625>

Submitted on 6 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Mini-Review

Mohamad El Haj*, Ahmed A. Moustafa, Jean Roche, Florence Pasquier, Dimitrios Kapogiannis, Karim Gallouj, Pascal Antoine

The “authentic subjective experience” of memory in Alzheimer’s disease

<https://doi.org/10.1515/tnsci-2020-0123>
received March 18, 2020; accepted May 4, 2020

Abstract: Most research has mainly focused on the decline of the subjective experience in Alzheimer’s disease (AD). However, few attempts have been made to evaluate whether subjective experience may be maintained in AD. In this narrative review, we attempt to provide a positive view, according to which patients with AD can enjoy, to some extent, subjective experience during memory retrieval. Memory and expression difficulties (e.g., aphasia) limit the ability of patients with AD to describe their memories, resulting in a little specificity of reported memories. However, according to the “authentic subjective experience” view, we propose in this study that the ability to mentally relive these memories could be preserved in the patients. By proposing the authentic subjective experience view, we attempt to provide an alternative view to the general consideration that the patients suffer a diminished subjective experience. This view can contribute to a larger clinical framework that gives a positive meaning to the subjective experience of

patients with AD. Furthermore, several clinical and empirical implications can be drawn from the authentic subjective experience view, including the possibility to evaluate behavioral correlates of the subjective experience in AD.

Keywords: Alzheimer’s disease, autobiographical memory, consciousness, subjective experience

1 Introduction

Memory decline is widely considered as the cognitive hallmark of typical Alzheimer’s disease (AD) [1]. Psychological research has been mainly concerned with the diminished ability of the patients to retrieve and relive past experiences [2]. In this study, we argue that patients with AD may still be able to enjoy an authentic subjective experience of memory, that is, they can experience an authentic sense of reliving during memory retrieval. At the clinical level, this view is worth considering because by defending the view that patients with AD possess an “authentic subjective experience” of the past, we attempt to offer a positive view of their potential to introspect into their memory functioning and cognitive performance in general.

This review begins by providing definitions of basic terms (e.g., subjective experience and lack of insight). We then emphasize empirical support to the authentic subjective experience view that we defend in this study. Finally, we discuss the implications of the view, as well as venues for future empirical research. Note that, while this review is narrative, we also performed a literature search for papers dealing with the subjective experience of autobiographical memory in AD. In this literature search, we used MEDLINE (via PubMed) and Google Scholar databases using a combination of the following search terms and key words: Alzheimer’s disease, anosognosia, autobiographical memory, auto-noetic consciousness, consciousness, denial, and insight. The search resulted in 464 papers, and we screened these

* **Corresponding author: Mohamad El Haj**, Univ Nantes, Laboratoire de psychologie des Pays de la Loire, LPPL, EA 4638, F-44000 Nantes, France; Unité de Gériatrie, Centre Hospitalier de Tourcoing, Tourcoing, France; Institut Universitaire de France, Paris, France, e-mail: mohamad.elhaj@univ-nantes.fr

Ahmed A. Moustafa: School of Social Sciences and Psychology & Marcs Institute for Brain and Behaviour, Western Sydney University, Sydney, Australia; Department of Human Anatomy and Physiology, the Faculty of Health Sciences, University of Johannesburg, Johannesburg, South Africa

Jean Roche: CHU de Lille, Unité de Psychogériatrie, Pôle de gérontologie, 59037 Lille, France

Florence Pasquier: University Lille Nord de France UDSL and Memory Clinic, CHU, Lille, France

Dimitrios Kapogiannis: Laboratory of Neurosciences, National Institute on Aging, Baltimore, MD, USA

Karim Gallouj: Unité de Gériatrie, Centre Hospitalier de Tourcoing, Tourcoing, France

Pascal Antoine: Univ. Lille, CNRS, CHU Lille, UMR 9193 – SCALab – Sciences Cognitives et Sciences Affectives, F-59000 Lille, France

papers to identify those dealing with the subjective experience of autobiographical memory in patients with AD, resulting in the below-discussed papers.

1.1 Basic definitions

The subjective experience of episodic memory can be defined with reference to “subjective experience” as proposed by Tulving [3] who associates episodic recall with a feeling of re-experiencing (i.e., autooetic consciousness), which refers to the ability to mentally project one's existence into a subjective time. Impairment in mental time travel can be associated with memory distortions in AD. According to the Temporal Consciousness Model [4], patients with confabulations (including those with AD) have a disturbed sense of chronology, such that they are able to retrieve the content of events but not their order of occurrence. Consequently, they misattribute the features of events that occurred at one time to other events that occurred at another time. Regardless of memory, the subjective experience in patients with AD is a complex and non-unitary phenomenon [5–8]. The same thing can be said for the concept of consciousness of general. Although consciousness can be defined as the awareness of the self and the environment [9], it also includes several phenomenological components (e.g., the subjective experience) and cognitive aspects (e.g., perception, affect, and control).

The literature proposes several terms (e.g., “denial,” “lack of insight,” and “anosognosia”) to describe the decline of the subjective experience in AD. The terms “denial” and “lack of insight” are typically used to describe a lack of introspective ability and metacognitive functioning, whereas the term “anosognosia” is typically used to describe a failure to acknowledge a particular cognitive deficit [10,11]. Theoretical accounts on the mechanism of anosognosia in AD have associated a lack of insight with a disconnection of the “conscious awareness system.” The conscious awareness system is a cognitive system that detects the changes in cognitive functioning based on the inputs from various brain regions [12,13]. Any disconnection between the conscious awareness system and the changes in a given cognitive domain (e.g., memory) results in a lack of awareness within that cognitive domain. A similar suggestion was made by the Cognitive Awareness Model [14,15]. This model is composed of components receiving information from autobiographical memory

and comparator components monitoring performances. All these components provide an output that feeds into awareness about the actual performances. According to the Cognitive Awareness Model, patients with unawareness of memory deficits have difficulties to encode those failures into memory, relying on outdated representations of their memory functioning. Research on anosognosia emphasizes the ability of patients with AD to estimate their own memory functioning and introspect into their own subjective experience during recall. While this research explains how the patients estimate and experience their own memory functioning, it runs the risk of denying the patients any level of awareness of their memories (e.g., “lack of awareness” and “anosognosia”). However, the concept of awareness in AD, at least the subjective experience of memory, is not necessarily negative since patients may be able to enjoy an authentic subjective experience of the past. We here discuss the research demonstrating how patients with AD can enjoy an authentic subjective experience of memory.

2 The authentic subjective experience view

In a previous study, we investigated whether patients with AD may report a high subjective experience of autobiographical memories [16]. We asked patients with mild AD and the control participants to retrieve three autobiographical memories. For each memory, the participants were invited to rate its subjective characteristics on items assessing travel in time, remembering, reliving, realness, visual imagery, auditory imagery, emotion, and importance. Besides this subjective evaluation, we analyzed the specificity of the retrieved memories (i.e., whether memories described general events or specific events situated in time and space). AD participants showed lower general autobiographical recall than the controls, and poorer reliving, travel in time, remembering, realness, visual imagery, auditory imagery, language, rehearsal, and spatial detail – a decrease that was especially pronounced for visual imagery. Yet, AD participants showed high rating for emotion and importance. Critically, a discrepancy (i.e., higher levels of subjective reliving than objective recall) was observed in patients with AD but not in the control participants. In other words, despite a compromise in their (objective) autobiographical performance, patients with AD attributed a high value to their subjective

experience, especially to emotion and importance of memory. This discrepancy could be due to an overestimation and alternatively to a high consciousness experience during which patients with mild AD possess authentic subjective experiences of the past. This interpretation is further supported by a study demonstrating that despite a compromise in some subjective features of recall (e.g., in the ability to create a visual image of the retrieved event), other subjective features (i.e., emotion and importance of the recall) can be fairly experienced by patients with mild AD [17].

We propose the authentic subjective experience view to explain the discrepancy between subjective reliving and objective recall in patients with AD. This is different from the anosognosia account. While the anosognosia account also deals with the subjective experience of remembering, this account attributes the failure to introspect into one’s own performance to a discrepancy between the expected and actual inputs. Our view emphasizes how patients with AD, and patients with amnesia in general [18], attribute a high value to their sense of reliving as a result of their authentic ability to experience the retrieved event rather than as a result of their failures to experience it.

As illustrated in Figure 1, the authentic subjective experience view is summarized as follows: when asked to retrieve and report autobiographical memories, patients with AD may struggle to describe them, probably due to difficulties in both retrieving and expressing their memories. However, these patients may be able to enjoy an authentic subjective experience of their memories (with corresponding emotions), as this experience does not require linguistic processing. The difficulty to describe specific autobiographical events in AD is in agreement with the large body of research demonstrating a diminished ability of the patients to retrieve and report specific autobiographical memories [19–32]. In this body of research, the patients are typically asked to describe verbally the retrieved memories. This verbal description may limit the patients’ ability to detail retrieved memories, which is supported by a research demonstrating significant correlations

between autobiographical retrieval and verbal fluency [33]. This suggestion can be also supported by the fact that AD affects several aspects of language, typically starting with semantics before proceeding to syntax and phonology and eventually leads to frank aphasia, a multimodal clinically significant disorder of understanding and producing language [34]. Moreover, these linguistic difficulties may limit the ability of patients with AD to describe memories and their subjective experience. The authentic subjective experience view can be supported by a research on normal aging, demonstrating that narrative style can hinder the ability of healthy older adults to retrieve specific memories [35,36]. The authentic subjective experience view can also be supported by a study demonstrating that narrative abilities influence the ability of patients with amnesia to describe pictures and imagined scenarios [37].

Several theoretical and clinical implications can be drawn from the authentic subjective experience view. At the theoretical level, this view contributes to the understanding of consciousness in patients with AD, especially to the understanding of how these patients relive and have a sense of ownership of memories. The authentic subjective experience view focuses on the ability of the patients to travel back in time to re-experience the past. The authentic subjective experience view can be also considered as an attempt to demonstrate how patients with AD can benefit from an authentic subjective experience when retrieving personal memories.

Another theoretical implication of the authentic subjective experience view is that it emphasizes the heterogeneity of consciousness experience in AD. Research on the subjective experience of memory in AD has typically evaluated different forms of this experience, such as sense of self [38], metamemory [39], anosognosia [15], or even denial [40], suggesting a heterogeneity of the consciousness experience in AD. The authentic subjective experience view extends this research by suggesting how, despite a compromise of anosognosia, other components can be relatively spared

- Mnemonic difficulties and expressive aphasia limit the ability of patients with Alzheimer’s Disease to describe their autobiographical memories, resulting in a little specificity of reported memories.
- However, the ability to mentally relive their memories, which is not subject to linguistic processing expressions, may be, to some extent, preserved in patients.

Figure 1: The “authentic subjective experience” account.

in AD (e.g., the subjective experience of autobiographical memory). It is noteworthy that this mixed picture can also be observed for specific consciousness experiences in AD. For instance, individual differences among AD patients in anosognosia can be observed depending on the assessment tool, the stage of dementia, and the type of memory or other cognitive function in question.

Regarding normal aging, the subjective experience during autobiographical retrieval was evaluated by Piolino *et al.* [41] who invited healthy older adults to retrieve recent and remote autobiographical memories. After memory retrieval, the participants had to provide a “Remember” response, if they were able to recover a specific event with its encoding context, and a “Know” response, if they just knew this event had happened to them but could not recall any contextual detail. Hence, Remember responses referred to auto-noetic consciousness, whereas Know responses referred to noetic consciousness. The results demonstrated increased Know responses with age for recent but not for remote memories, supporting the view that auto-noetic consciousness is relatively better preserved in healthy older adults with respect to the distant past than with respect to the more recent past. Note that, using the Remember/Know paradigm, a research has demonstrated increased Remember responses in patients with AD [42,43], suggesting a decreased auto-noetic consciousness in AD. However, this research has been concerned with memory for simple stimuli (e.g., words and pictures) rather than with autobiographical memory.

One theoretical implication of the authentic subjective experience view is its potential effect on the patients’ sense of ownership of past experiences (i.e., the feeling that these experiences belong to them). This speculation can be compared with that of James [44] who distinguished between the self as an agent (the I) and the self as an object of reflection (the Me). According to James [44], personal ownership, as dependent on the “I”, is accompanied by a subjective experience imbued with a sense of warmth and intimacy. This colorful reliving can be associated with the subjective experience of the past in patients with AD, an assumption supported by our previous research demonstrating that these patients attribute high personal importance to their recall [16,17].

Altogether, the authentic subjective experience view can be considered as the outcome of the interaction between representational content (i.e., the retrieved memory), conscious apprehension, and sense of ownership resulting into the warmth and intimacy that characterizes the subjective experience of the past.

3 Toward a positive view of the consciousness experience in patients

As illustrated in the previous section, one implication of the authentic subjective experience view is that whereas some aspects of the consciousness experience can be affected by AD (e.g., anosognosia), other aspects (e.g., the ability to relive autobiographical memories) can be fairly experienced by patients with AD. At the clinical level, this implication is important, as it emphasizes how patients with AD can generate a subjective experience. Another implication is that by enjoying a fair subjective experience of autobiographical memories, patients with AD may be able to experience a rich sense of self and identity unlike what is generally assumed. This speculation can be supported by a research demonstrating an intimate relationship between the autobiographical performance and the sense of self in AD [33,45].

The authentic subjective experience view is also an attempt to alleviate the social stigma that can be associated with the disease. Needless to say, AD is mainly associated with dysfunction and poor cognitive abilities, and the social stigma tends to depict the disease as the “death sentence” and the patients as “empty shells” who experience a “death of the mind” [46,47]. The authentic subjective experience view can be considered as an alternative that identifies probably partially spared components of consciousness (i.e., the subjective experience of past) in patients with AD. We hope that this positive signal may prompt other researchers and clinicians to investigate, or at least consider, remaining intact cognitive functions in AD. We also hope that this view contributes to care models that give positive meaning to the consciousness experience of the patients and their ability to introspect into their cognitive functioning and, consequently, into their ability to judge, think, plan, and decide.

3.1 Perspectives: behavioral evaluation of the authentic subjective experience

One major challenge for autobiographical retrieval in AD, besides mnemonic difficulties, is aphasia. The degree to which an authentic subjective experience occurs may be partially beyond language-based description. Nevertheless, the validity of this view can be subjected to rigorous research to delineate the behavioral correlates of autobiographical memory in AD using non-linguistic means. This future research may be based on facial expression techniques. Due to

advances in the technology allowing a reliable analysis of facial expressions, there is a burgeoning interest in facial expressions that can be activated during autobiographical retrieval [48,49]. A study reported that emotional memories trigger the corresponding basic facial expressions (i.e., high happy facial expressions for happy memories and sad facial expressions for sad memories) [50]. Another study reported higher happy emotional facial expressions during imagining the future than during remembering the past [51]. It would be of interest to evaluate whether emotional facial expressions are triggered by retrieval of autobiographical memories in AD, as such expressions would provide behavioral evidence of the hypothesis that the patients subjectively experience emotional characteristics of their past.

Any behavioral evidence of the hypothesis that the patients possess authentic subjective experiences of their past would be of valuable interest to understand the subjective experience of the patients in the advanced stages of AD. The study supporting the authentic subjective experience hypothesis was conducted in patients with mild AD [16,17]; therefore, research is needed to investigate this experience in patients with advanced AD. This issue is important because advanced AD is characterized by severe language impairment and apathy [52], limiting the ability of the patients to describe memories and their subjective experience. We believe that behavioral studies, such as those on facial expressions, would provide valuable evidence of the hypothesis that patients with advanced AD may possess authentic subjective experiences of their past. Another suggestion for future research would be to examine neural basis of the authentic subjective experience hypothesis. While there is a lack of research on this hypothesis, a research has shown involvement of the hippocampus in the decline of autobiographical memory [53,54]. It would be of interest to investigate whether, despite disruption of the hippocampus, the authentic subjective experience can be supported by other brain regions that are less affected by the disease, at least during the early stages of the AD, such as the visual cortices [55,56].

4 Summary

While research on the consciousness experience in AD has been mainly concerned with the compromise of this experience, the authentic subjective experience view attempts to provide an alternative view according to

which patients with AD can enjoy, to some extent, an authentic subjective experience during memory retrieval. Although it is important to acknowledge that some components of consciousness experience are affected by AD (e.g., anosognosia), it is equally important to consider whether patients can positively experience other components such as emotion. By doing this, we hope that this narrative review will contribute to a care model that acknowledges and honors the patients’ subjective experiences, provides positive meaning to them, and uses them as a tool in managing the disease.

Acknowledgments: This work was supported by the LABEX (excellence laboratory, program investment for the future), DISTALZ (Development of Innovative Strategies for a Transdisciplinary Approach to Alzheimer Disease), and the EU Interreg CASCADE 2 Seas Programme 2014–2020 (co-funded by the European Regional Development Fund). This research was supported in part (for the author DK) by the Intramural Research Program of the National Institute on Aging, NIH.

Conflicts of interest: Authors state no conflict of interest.

References

- [1] McKhann G, Knopman DS, Chertkow H, Hyman BT, Jack Jr. CR, Kawas CH, et al. The diagnosis of dementia due to Alzheimer’s disease: recommendations from the National Institute on Aging-Alzheimer’s Association workgroups on diagnostic guidelines for Alzheimer’s disease. *Alzheimer’s Dement: J Alzheimer’s Assoc.* 2011;7(3):263–9.
- [2] El Haj M, Antoine P, Nandrino JL, Kapogiannis D. Autobiographical memory decline in Alzheimer’s disease, a theoretical and clinical overview. *Ageing Res Rev.* 2015;23(Pt B):183–92.
- [3] Tulving E. Episodic memory: from mind to brain. *Annu Rev Psychol.* 2002;53:1–25.
- [4] Dalla Barba G, Cappelletti JY, Signorini M, Denes G. Confabulation: remembering ‘another’ past, planning ‘another’ future. *Neurocase.* 1997;3(6):425–36.
- [5] Vasterling JJ, Seltzer B, Carpenter BD, Thompson KA. Unawareness of social interaction and emotional control deficits in Alzheimer’s disease. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn.* 1997;4(4):280–9.
- [6] Starkstein SE, Sabe L, Chemerinski E, Jason L, Leiguarda R. Two domains of anosognosia in Alzheimer’s disease. *J Neurol Neurosurg Psychiatr.* 1996;61(5):485–90.
- [7] Avondino E, Antoine P. Heterogeneity of cognitive anosognosia and its variation with the severity of dementia in patients with Alzheimer’s disease. *J Alzheimer’s Dis.* 2015;50(1):89–99.
- [8] Antoine P, Nandrino JL, Billiet C. Awareness of deficits in Alzheimer’s disease patients: analysis of performance

- prediction discrepancies. *Psychiatry Clin Neurosci.* 2013;67(4):237–44.
- [9] Jennett B, Plum F. Persistent vegetative state after brain damage. A syndrome in search of a name. *Lancet.* 1972;1(7753):734–7.
- [10] Amanzio M, Torta DM, Sacco K, Cauda F, D'Agata F, Duca S, et al. Unawareness of deficits in Alzheimer's disease: role of the cingulate cortex. *Brain.* 2011;134(Pt 4):1061–76.
- [11] Babinski J. Contribution a l'étude des troubles mentaux dans l'hémiplégie organique cérébrale (anosognosie). *Revue Neurologique.* 1914;27:845–7.
- [12] Grut M, Jorm AF, Fratiglioni L, Forsell Y, Viitanen M, Winblad B. Memory complaints of elderly people in a population survey: variation according to dementia stage and depression. *J Am Geriatr Soc.* 1993;41(12):1295–300.
- [13] McGlynn SM, Schacter DL. Unawareness of deficits in neuropsychological syndromes. *J Clin Exp Neuropsychol.* 1989;11(2):143–205.
- [14] Agnew SK, Morris RG. The heterogeneity of anosognosia for memory impairment in Alzheimer's disease: a review of the literature and a proposed model. *Aging Mental Health.* 1998;2(1):7–19.
- [15] Morris RG, Mograbi DC. Anosognosia, autobiographical memory and self knowledge in Alzheimer's disease. *Cortex.* 2013;49(6):1553–65.
- [16] El Haj M, Antoine P. Discrepancy between subjective autobiographical reliving and objective recall: the past as seen by Alzheimer's disease patients. *Consciousness Cognit.* 2017;49:110–6.
- [17] El Haj M, Kapogiannis D, Antoine P. Phenomenological reliving and visual imagery during autobiographical recall in Alzheimer's disease. *J Alzheimer's Dis.* 2016;52(2):421–31.
- [18] El Haj M, Nandrino JL. Seeing life through rose-colored spectacles: autobiographical memory as experienced in Korsakoff's syndrome. *Consciousness Cognit.* 2018;60:9–16.
- [19] Barnabe A, Whitehead V, Pilon R, Arsenault-Lapierre G, Chertkow H. Autobiographical memory in mild cognitive impairment and Alzheimer's disease: a comparison between the Levine and Kopelman interview methodologies. *Hippocampus.* 2012;22(9):1809–25.
- [20] Graham KS, Hodges JR. Differentiating the roles of the hippocampal complex and the neocortex in long-term memory storage: evidence from the study of semantic dementia and Alzheimer's disease. *Neuropsychology.* 1997;11(1):77–89.
- [21] Greene JD, Hodges JR, Baddeley AD. Autobiographical memory and executive function in early dementia of Alzheimer type. *Neuropsychologia.* 1995;33(12):1647–70.
- [22] Irish M, Lawlor BA, O'Mara SM, Coen RF. Impaired capacity for auto-note-taking during autobiographical event recall in mild Alzheimer's disease. *Cortex.* 2011;47(2):236–49.
- [23] Ivanoiu A, Cooper JM, Shanks MF, Venneri A. Patterns of impairment in autobiographical memory in the degenerative dementias constrain models of memory. *Neuropsychologia.* 2006;44(10):1936–55.
- [24] Leyhe T, Muller S, Milian M, Eschweiler GW, Saur R. Impairment of episodic and semantic autobiographical memory in patients with mild cognitive impairment and early Alzheimer's disease. *Neuropsychologia.* 2009;47(12):2464–9.
- [25] Muller S, Saur R, Greve B, Melms A, Hautzinger M, Fallgatter AJ, et al. Similar autobiographical memory impairment in long-term secondary progressive multiple sclerosis and Alzheimer's disease. *Mult Scler.* 2013;19(2):225–32.
- [26] Seidl U, Lueken U, Thomann PA, Geider J, Schroder J. Autobiographical memory deficits in Alzheimer's disease. *J Alzheimer's Dis.* 2011;27(3):567–74.
- [27] El Haj M, Postal V, Le Gall D, Allain P. Directed forgetting of autobiographical memory in mild Alzheimer's disease. *Memory.* 2011;19(8):993–1003.
- [28] El Haj M, Fasotti L, Allain P. The involuntary nature of music-evoked autobiographical memories in Alzheimer's disease. *Conscious Cogn.* 2012;21(1):238–46.
- [29] El Haj M, Antoine P, Nandrino JL, Gely-Nargeot MC, Raffard S. Self-defining memories during exposure to music in Alzheimer's disease. *Int Psychogeriatr.* 2015;27(10):1719–30.
- [30] De Simone MS, Fadda L, Perri R, Aloisi M, Caltagirone C, Carlesimo GA. Does retrieval frequency account for the pattern of autobiographical memory loss in early Alzheimer's disease patients? *Neuropsychologia.* 2016;80:194–200.
- [31] Martinelli P, Anssens A, Sperduti M, Piolino P. The influence of normal aging and Alzheimer's disease in autobiographical memory highly related to the self. *Neuropsychology.* 2013;27(1):69–78.
- [32] Kirk M, Berntsen D. A short cut to the past: cueing via concrete objects improves autobiographical memory retrieval in Alzheimer's disease patients. *Neuropsychologia.* 2018;110:113–22.
- [33] El Haj M, Antoine P. Describe yourself to improve your autobiographical memory: a study in Alzheimer's disease. *Cortex.* 2017;88:165–72.
- [34] Kirshner HS. Primary progressive aphasia and Alzheimer's disease: brief history, recent evidence. *Curr Neurol Neurosci Rep.* 2012;12(6):709–14.
- [35] Gaesser B, Sacchetti DC, Addis DR, Schacter DL. Characterizing age-related changes in remembering the past and imagining the future. *Psychol Aging.* 2011;26(1):80–4.
- [36] Madore KP, Schacter DL. An episodic specificity induction enhances means-end problem solving in young and older adults. *Psychol Aging.* 2014;29(4):913–24.
- [37] Dewar M, Neroni MA, Zeman A, Beschin N, Della Sala S. Visual complexity accentuates picture-description deficit in amnesia. *Neuropsychology.* 2017;31(6):689–96.
- [38] Prebble SC, Addis DR, Tippett LJ. Autobiographical memory and sense of self. *Psychol Bull.* 2013;139(4):815–40.
- [39] Souchay C. Metamemory in Alzheimer's disease. *Cortex.* 2007;43(7):987–1003.
- [40] Sevush S, Leve N. Denial of memory deficit in Alzheimer's disease. *Am J Psychiatr.* 1993;150(5):748–51.
- [41] Piolino P, Desgranges B, Clarys D, Guillery-Girard B, Tacconat L, Isingrini M, et al. Autobiographical memory, auto-note-taking, and self-perspective in aging. *Psychol Aging.* 2006;21(3):510–25.
- [42] Hudon C, Belleville S, Gauthier S. The assessment of recognition memory using the Remember/Know procedure in amnesic mild cognitive impairment and probable Alzheimer's disease. *Brain Cogn.* 2009;70(1):171–9.
- [43] Ally BA. Using pictures and words to understand recognition memory deterioration in amnesic mild cognitive impairment

- and Alzheimer's disease: a review. *Curr Neurol Neurosci Rep.* 2012;12(6):687–94.
- [44] James W. *Psychology: the briefer course.* New York, NY: Holt; 1892.
- [45] Addis DR, Tippett LJ. Memory of myself: autobiographical memory and identity in Alzheimer's disease. *Memory.* 2004;12(1):56–74.
- [46] Van Gorp B, Vercruyssen T. Frames and counter-frames giving meaning to dementia: a framing analysis of media content. *Soc Sci Med.* 2012;74(8):1274–81.
- [47] Beard RL, Neary TM. Making sense of nonsense: experiences of mild cognitive impairment. *Sociol Health Illn.* 2013;35(1):130–46.
- [48] El Haj M, Daoudi M, Gallouj K, Moustafa AA, Nandrino J-L. When your face describes your memories: facial expressions during retrieval of autobiographical memories. *Rev Neurosci.* 2018;29(8):861–72.
- [49] Gandolphe MC, Nandrino JL, Delelis G, Ducro C, Lavallee A, Saloppe X, et al. Positive facial expressions during retrieval of self-defining memories. *J Integr Neurosci.* 2018;17(3–4):367–76.
- [50] El Haj M, Antoine P, Nandrino JL. More emotional facial expressions during episodic than during semantic autobiographical retrieval. *Cognit Affect Behav Neurosci.* 2016;16(2):374–81.
- [51] El Haj M, Antoine P, Nandrino JL. Facial expressions triggered by imagining the future. *J Integr Neurosci.* 2017;16(4):483–92.
- [52] Klimova B, Maresova P, Valis M, Hort J, Kuca K. Alzheimer's disease and language impairments: social intervention and medical treatment. *Clin Interv Aging.* 2015;10:1401–7.
- [53] Philippi N, Noblet V, Botzung A, Despres O, Renard F, Sfikas G, et al. MRI-based volumetry correlates of autobiographical memory in Alzheimer's disease. *PLoS One.* 2012;7(10):e46200.
- [54] Philippi N, Botzung A, Noblet V, Rousseau F, Despres O, Cretin B, et al. Impaired emotional autobiographical memory associated with right amygdalar-hippocampal atrophy in Alzheimer's disease patients. *Front Aging Neurosci.* 2015;7:21.
- [55] Markowitsch HJ, Vandekerckhove MM, Lanfermann H, Russ MO. Engagement of lateral and medial prefrontal areas in the ephory of sad and happy autobiographical memories. *Cortex.* 2003;39(4–5):643–65.
- [56] Piefke M, Weiss PH, Zilles K, Markowitsch HJ, Fink GR. Differential remoteness and emotional tone modulate the neural correlates of autobiographical memory. *Brain.* 2003;126(Pt 3):650–68.