



**HAL**  
open science

## Effect of Waters Enriched in O<sub>2</sub> by Injection or Electrolysis on Performance and the Cardiopulmonary and Acid-Base Response to High Intensity Exercise.

F. Daussin, François Péronnet, Antoine Charton, Evelyne Lonsdorfer, Stéphane Doutreleau, Bernard Geny, Ruddy Richard

### ► To cite this version:

F. Daussin, François Péronnet, Antoine Charton, Evelyne Lonsdorfer, Stéphane Doutreleau, et al.. Effect of Waters Enriched in O<sub>2</sub> by Injection or Electrolysis on Performance and the Cardiopulmonary and Acid-Base Response to High Intensity Exercise.. *Nutrients*, 2021, *Nutrients*, 13 (12), pp.4320. 10.3390/nu13124320 . hal-03522530

**HAL Id: hal-03522530**

<https://hal.univ-lille.fr/hal-03522530>

Submitted on 12 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Article

# Effect of Waters Enriched in O<sub>2</sub> by Injection or Electrolysis on Performance and the Cardiopulmonary and Acid–Base Response to High Intensity Exercise

Frédéric N. Daussin<sup>1,\*</sup>, François Péronnet<sup>2</sup>, Antoine Charton<sup>3,4</sup>, Evelyne Lonsdorfer<sup>4,5</sup>, Stéphane Doutreleau<sup>6</sup>, Bernard Geny<sup>4,5</sup> and Ruddy Richard<sup>7</sup>

<sup>1</sup> ULR 7369–URePSSS—Unité de Recherche Pluridisciplinaire Sport Santé Société, Univ. Lille, Univ. Artois, Univ. Littoral Côte d’Opale, F-59000 Lille, France

<sup>2</sup> École de Kinésiologie et des Sciences de L’activité Physique, Faculté de Médecine, Université de Montréal, Montréal, QC H3C 3J7, Canada; francois.peronnet@umontreal.ca

<sup>3</sup> Department of Anaesthesia and Critical Care, EA 3072, University of Strasbourg, Hautepierre University Hospital, F-67091 Strasbourg, France; acharton@hotmail.fr

<sup>4</sup> Physiology Department, EA 3072, Faculty of Medicine, University of Strasbourg, F-67091 Strasbourg, France; Evelyne.lonsdorfer@chru-strasbourg.fr (E.L.); bernard.geny@chru-strasbourg.fr (B.G.)

<sup>5</sup> Department of Physiology and Functional Explorations, CHRU Strasbourg, F-67091 Strasbourg, France

<sup>6</sup> Department of Physiology, INSERM U1042, F-38000 Grenoble, France; sdoutreleau@chu-grenoble.fr

<sup>7</sup> CRNH Auvergne, INRA UMR1019 and Department of Sport Medicine and Functional Explorations, CHU Clermont-Ferrand, F-63003 Clermont-Ferrand, France; ruddy.richard@uca.fr

\* Correspondence: frederic.daussin@univ-lille.fr



**Citation:** Daussin, F.N.; Péronnet, F.; Charton, A.; Lonsdorfer, E.; Doutreleau, S.; Geny, B.; Richard, R. Effect of Waters Enriched in O<sub>2</sub> by Injection or Electrolysis on Performance and the Cardiopulmonary and Acid–Base Response to High Intensity Exercise. *Nutrients* **2021**, *13*, 4320. <https://doi.org/10.3390/nu13124320>

Academic Editor: Antoni Sureda

Received: 25 October 2021

Accepted: 26 November 2021

Published: 29 November 2021

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Several brands of water enriched with O<sub>2</sub> (O<sub>2</sub>-waters) are commercially available and are advertised as wellness and fitness waters with claims of physiological and psychological benefits, including improvement in exercise performance. However, these claims are based, at best, on anecdotal evidence or on a limited number of unreliable studies. The purpose of this double-blind randomized study was to compare the effect of two O<sub>2</sub>-waters (~110 mg O<sub>2</sub>·L<sup>-1</sup>) and a placebo (10 mg O<sub>2</sub>·L<sup>-1</sup>, i.e., close to the value at sea level, 9–12 mg O<sub>2</sub>·L<sup>-1</sup>) on the cardiopulmonary responses and on performance during high-intensity exercise. One of the two O<sub>2</sub>-waters and the placebo were prepared by injection of O<sub>2</sub>. The other O<sub>2</sub>-water was enriched by an electrolytic process. Twenty male subjects were randomly allocated to drink one of the three waters in a crossover study (2 L·day<sup>-1</sup> × 2 days and 15 mL·kg<sup>-1</sup> 90 min before exercise). During each exercise trial, the subjects exercised at 95.9 ± 4.7% of maximal workload to volitional fatigue. Exercise time to exhaustion and the cardiopulmonary responses, arterial lactate concentration and pH were measured. Oxidative damage to proteins, lipids and DNA in blood was assessed at rest before exercise. Time to exhaustion (one-way ANOVA) and the responses to exercise (two-way ANOVA [Time; Waters] with repeated measurements) were not significantly different among the three waters. There was only a trend (*p* = 0.060) for a reduction in the time constant of the rapid component of VO<sub>2</sub> kinetics with the water enriched in O<sub>2</sub> by electrolysis. No difference in oxidative damage in blood was observed between the three waters. These results suggest that O<sub>2</sub>-water does not speed up cardiopulmonary response to exercise, does not increase performance and does not trigger oxidative stress measured at rest.

**Keywords:** performance; reactive oxygen species; VO<sub>2</sub> kinetics; endurance exercise

## 1. Introduction

At equilibrium with partial pressure at sea level (PO<sub>2</sub> ~160 mmHg) and for temperature ranging between 10 and 25 °C, the amount of O<sub>2</sub> dissolved in water ranges respectively between 12 and 9 mg·L<sup>-1</sup> [1] (Box 1). However, several commercially available brands of water are purportedly enriched in O<sub>2</sub> (O<sub>2</sub>-waters) at 7 and 40 times this basal value [2–4] and are advertised as wellness and fitness waters with a large range of physiological and

psychological benefits. As discussed by several authors [2,3,5,6], these claims are based at best on anecdotal evidences or on a limited number of unreliable studies.

**Box 1.** Relationship between PO<sub>2</sub> and temperature, and O<sub>2</sub> content of water according to Henry's law [1] expressed in various units: mol·L<sup>-1</sup>, mL·L<sup>-1</sup>, mg·L<sup>-1</sup> and ppm, at sea level, 10 °C and dry air.

The amount of O<sub>2</sub> dissolved in water (mol·L<sup>-1</sup>) depends on the temperature (T in K = T[°C] + 273) and PO<sub>2</sub> (in atmosphere: atm) and is described by Henry's law where 769.23 and 1700 are Henry's constants for O<sub>2</sub>:

$$\text{O}_2 \text{ content (mol}\cdot\text{L}^{-1}) = \text{PO}_2 \text{ (atm)} / \{769.23 \times \exp[-1700 \times (1/T - 1/298)]\}$$

For example, at sea level (PO<sub>2</sub> = 160 mmHg or 0.21 atm in dry air) and 10 °C (283 K):

$$\text{O}_2 \text{ content (mol}\cdot\text{L}^{-1}) = 0.21 \text{ (atm)} / \{769.23 \times \exp[-1700 \times (1/283 - 1/298)]\} = 0.000370 \text{ mol}\cdot\text{L}^{-1}$$

Since the molar mass of O<sub>2</sub> is 32 g and since 1 mol of O<sub>2</sub> occupies 22.4 L, the conversion of this O<sub>2</sub> content in various other units is as follows:

$$\text{O}_2 \text{ content in mL}\cdot\text{L}^{-1} = \text{O}_2 \text{ content in mol}\cdot\text{L}^{-1} \times 22.4 \times 1000 = 0.000370 \times 22.4 \times 1000 = 8.29 \text{ mL}\cdot\text{L}^{-1}$$

$$\text{O}_2 \text{ content in mg}\cdot\text{L}^{-1} = \text{O}_2 \text{ content in mol}\cdot\text{L}^{-1} \times 32/1000 = 0.000370 \times 32/1000 = 11.8 \text{ mg}\cdot\text{L}^{-1}$$

Finally, since the molar mass of water is 18 g, there are 1000/18 = 55.5 mol of water·L<sup>-1</sup>. The O<sub>2</sub> content in ppm, thus is:

$$\text{O}_2 \text{ content in ppm} = \text{O}_2 \text{ content in mol}\cdot\text{L}^{-1} / \text{H}_2\text{O content in mol}\cdot\text{L}^{-1} = (0.000370/55.5) \times 10^6 = 6.67 \text{ ppm}$$

Consumption of O<sub>2</sub>-waters is promoted in active subjects, and a dozen studies have been conducted to document their effects, if any, on the response to exercise and on performance (Table 1). However, in some of these studies, the characteristics of the water ingested, including the O<sub>2</sub>-content, were not precisely described, and the volume ingested was not indicated. In only two studies [2,7], the O<sub>2</sub> content measured and/or reported appears reliable and in the range of values for which benefits are claimed by the manufacturers. In these two studies, O<sub>2</sub>-waters consumption did not improve maximal oxygen consumption (VO<sub>2</sub>max) [2,7] and in the study by Leibetseder et al. [7], the only significant difference when the O<sub>2</sub>-water was ingested was an increase in the ventilatory equivalent of oxygen (pulmonary ventilation/O<sub>2</sub> consumption or VE/VO<sub>2</sub>) at sub-maximal exercise and in plasma lactate concentration at maximal exercise. As discussed by the authors, it is difficult to speculate about the physiological significance of these changes, their possible effects on performance and how they could be related to ingestion of the O<sub>2</sub>-water, but these findings suggest that O<sub>2</sub>-waters could modify the cardiopulmonary response to exercise.

As already discussed [2,4,8], ingestion of water enriched in O<sub>2</sub> is unlikely to directly modify oxygen consumption (VO<sub>2</sub>) at the mouth during exercise. In addition, in arterial blood, the amount of dissolved O<sub>2</sub> is very small compared to the amount carried by hemoglobin, and in healthy subjects exercising at sea-level or a low altitude (<1000 m), arterial hemoglobin saturation remains close to normal values. It is, thus, very unlikely that increasing O<sub>2</sub> delivery to the blood through the gut will markedly increase PO<sub>2</sub> and O<sub>2</sub> content in arterial blood and, thus, O<sub>2</sub> delivery to any organ or tissue, including exercising muscles. However, we have previously studied the effects of a water enriched in O<sub>2</sub>, using an electrolytic process [9–11]. When compared to injection of O<sub>2</sub>, this process could generate water superstructures called clathrates, which could trap solutes, such as O<sub>2</sub>, and which can facilitate O<sub>2</sub> diffusion along PO<sub>2</sub> gradients [12–14]. This hypothesis is consistent with the higher tissue oxygenation observed in anaesthetized pigs [11] and the higher mitochondrial respiration observed at low PO<sub>2</sub> in permeabilized rat muscle fibers [10] with water enriched by electrolysis vs. injection.

It has shown that the fast component of VO<sub>2</sub> kinetics at the mouth in response to constant load exercise reflects O<sub>2</sub> diffusion from the blood to muscle fibers [15]. Therefore, the aim of this study was to investigate the effect of O<sub>2</sub>-waters on the cardiopulmonary and acid–base responses during near maximal exercise continued to exhaustion. We hypothesized that the fast component of the kinetics of VO<sub>2</sub> in response to high-intensity exercise could be faster following ingestion of the water enriched in O<sub>2</sub> by electrolysis vs. injection or a control water with a low O<sub>2</sub> content. A faster adjustment of VO<sub>2</sub> at the onset of exercise could increase the time to exhaustion. Finally, it has been suggested that ingestion of O<sub>2</sub>-waters can be a safety issue because of production of reactive oxygen

species (ROS) [5,6,16,17]. We, thus, also investigated the presence of ROS induces damages, including DNA damages, using the comet assay.

**Table 1.** Studies of the effects of O<sub>2</sub>-water ingestion on performance and on the response to exercise.

Reference	Water Ingested	Reported (1) O <sub>2</sub> Content or PO <sub>2</sub>	Computed (2) O <sub>2</sub> Content (mg·L <sup>-1</sup> ) and PO <sub>2</sub> at 10 °C	Observations
Askew et al. 2001 [17]	Stabilized O <sub>2</sub> in water Ingestion for 8 days Volume not reported	30,000 ppm	~53,700 mg·L <sup>-1</sup> ~725,000 mmHg ~950 atm	No significant difference between the O <sub>2</sub> -water and the placebo in response to a graded exercise to VO <sub>2</sub> max
Duncan 1997 [18]	Oxygen-enhanced water Volume not reported	Not reported	?	5 km run (min:s) Placebo: 21:18 O <sub>2</sub> -water: 20:47 (not significant)
Fleming et al. 2017 [8]	Activate Stabilized Water (ASO): 0.9 L during and after exercise	35 g in 62 g of water	565 mg·L <sup>-1</sup> ~7,600,000 mmHg ~10,000 atm	5-km run: Lactate clearance (t <sub>1/2</sub> in seconds) Placebo: 1223 O <sub>2</sub> -water: 1127 ( <i>p</i> < 0.05)
Fuller 2010 [19]	Activate Stabilized Water (ASO): 0.5 L during and after exercise	5 mg·L <sup>-1</sup>	5 mg·L <sup>-1</sup> 68 mmHg 0.09 atm	Trend for a longer time to exhaustion during a graded test to VO <sub>2</sub> max with the O <sub>2</sub> -water than the placebo (451 vs. 429 s)
Hampson et al. 2003 [2]	Oxygenated water 355 mL 5 min before exercise	1,184 mmHg (3) 226 mL·L <sup>-1</sup>	88 mg·L <sup>-1</sup> (3) 61.4 mL·L <sup>-1</sup> 1.56 atm	No significant difference between the O <sub>2</sub> -water and the placebo in response to a graded exercise to VO <sub>2</sub> max
Jenkins et al. 2001 [20]	Oxygenated water 0.45 L 10 min before and after exercise	Not reported	?	Higher hemoglobin saturation in arterial blood at the end of exercise at 100%VO <sub>2</sub> max with the O <sub>2</sub> -water than the placebo (94 vs. 87%)
Leibetseder et al. 2006 [7]	Oxygenated water 1.5 L·day <sup>-1</sup> for 2 weeks	160 mg·L <sup>-1</sup>	160 mg·L <sup>-1</sup> 2150 mmHg 2.83 atm	Higher VE/VO <sub>2</sub> at submaximal workload and higher lactate concentration at maximal workload with the O <sub>2</sub> -water than the placebo
McNaughton et al. 2007 [21]	Superoxygenated water (Oxyshot) 15 mL 30 min before exercise	150,000 ppm (4)	266,000 mg·L <sup>-1</sup> ~ 3,600,000 mmHg ~ 4750 atm	No significant difference for a 45-min exercise at 70%VO <sub>2</sub> max followed by a 15-min time trial to exhaustion between the O <sub>2</sub> -water and the placebo
Mielke et al. 2004 [22]	Oxygenated water 1.2 L·day <sup>-1</sup> for 3 days and 0.6 L 15 min before exercise	13.1 mg·L <sup>-1</sup>	13.1 mg·L <sup>-1</sup> 177 mmHg 0.23 atm	No significant difference in response to a graded exercise to VO <sub>2</sub> max or in exercise time to exhaustion at 90%VO <sub>2</sub> max between the O <sub>2</sub> -water and the placebo
Willmert et al. 2002 [4]	Super oxygenated water 0.5 L 15 min before exercise	13.5 mL·L <sup>-1</sup>	19.3 mg·L <sup>-1</sup> 260 mmHg 0.34 atm	No significant difference between the O <sub>2</sub> -water and the placebo in response to a graded exercise to VO <sub>2</sub> max
Wing-Gaia et al. 2005 [23]	Purified oxygen water 35 mL·kg <sup>-1</sup> ·day <sup>-1</sup> for 3 days 0.5 L 2 h before exercise	Not reported	?	No significant difference in performance or in response to a time trial at 57–59%VO <sub>2</sub> max in hypoxic condition (~76 min) between the O <sub>2</sub> -water and the placebo

Table 1. Cont.

Reference	Water Ingested	Reported (1) O <sub>2</sub> Content or PO <sub>2</sub>	Computed (2) O <sub>2</sub> Content (mg·L <sup>-1</sup> ) and PO <sub>2</sub> at 10 °C	Observations
Zhang et al. 2005 [24]	Hyperoxia solution 0.25 L before exercise	170 mL·0.5 L <sup>-1</sup>	481 mg·L <sup>-1</sup> 6500 mmHg 8.55 atm	Lower plasma lactate concentration in response to a 5 km run at altitude (2000 and 4000 m) with the O <sub>2</sub> -water than the placebo

(1) Values reported by the authors. (2) Values computed from the data reported by the authors using Henry's law [1] at 10 °C. (3) The O<sub>2</sub> contents computed in the study by Hampson et al. [2] from the PO<sub>2</sub> measured in tap water and five brands of O<sub>2</sub>-waters (e.g., 226 mL·L<sup>-1</sup> for 1184 mmHg in the brand 5 studied at exercise) are all in error. The correct value of O<sub>2</sub>-content for a PO<sub>2</sub> = 1184 mmHg and at 10 °C is 61 mL·L<sup>-1</sup> corresponding to 88 mg·L<sup>-1</sup>. The O<sub>2</sub> content in «well-stirred» tap water with a PO<sub>2</sub> of 127 mmHg is 6.6 mL·L<sup>-1</sup> at 10 °C and 4 mL·L<sup>-1</sup> at 37 °C (i.e., well below the value reported of 25 mL·L<sup>-1</sup>) which is in line with the textbook value of 3 mL·L<sup>-1</sup> dissolved in arterial blood at a PO<sub>2</sub> of ~100 mmHg and 37 °C [14]. (4) Not reported by the authors but found at website (<https://www.reachforlife.com.au/Equine/EQ-Product.php>, consulted 22 November 2021).

## 2. Methods

### 2.1. Subjects

Twenty active and healthy male subjects, recruited through advertisements in the hospital and university communities, volunteered for this study, which was approved by the local ethic committee (Comité de Protection des Personnes Est IV, Strasbourg, France: Eudract No.2008-A01051-54). Their age, height, body mass and percent body fat were respectively  $22.8 \pm 4.1$  years,  $178.7 \pm 5.9$  cm,  $70.0 \pm 5.7$  kg and  $13.0 \pm 1.6\%$  (mean  $\pm$  SD). In accordance with the declaration of Helsinki, all participants signed an informed-consent form. The sample size was determined based on the significant difference in plasma lactate concentration at maximal exercise between the control and O<sub>2</sub>-water by Leibetseder et al. [7] ( $9.6$  vs.  $11.1$  mmol·L<sup>-1</sup> with SD =  $1.6$  mmol·L<sup>-1</sup>). These data indicated that 18 subjects were needed to detect a difference with an 80% power and  $p = 0.05$  [25].

### 2.2. Study Design

The participants visited the laboratory on four separate occasions. All experimental conditions were conducted in a climate-controlled laboratory at Strasbourg Hospital. The first session was used to determine the VO<sub>2</sub>max and maximal power output by an incremental test to exhaustion. On the subsequent three visits separated by at least 5 days, the subjects performed an exercise trial to volitional fatigue. The study was randomized and double blind. The cardiopulmonary and acid–base response and time to exhaustion were measured.

### 2.3. Determination of VO<sub>2</sub>max and Pmax

VO<sub>2</sub>max and the corresponding workload (Pmax) were measured before the first experimental trial on cycle ergometer (Ergoline 900, Ergoline, Schiller, France), using an incremental test to volitional fatigue, as previously described [26]. The VO<sub>2</sub> and carbon dioxide production (VCO<sub>2</sub>) were computed from breath-by-breath measurement of gas exchanges at the mouth (Sensor Medics, Yorba Linda, CA, USA) and the heart rate was measured from a 12-lead ECG (Cardiovit CS200, Schiller, Baar, Switzerland). The Pmax was considered to be the lowest workload, eliciting VO<sub>2</sub>max =  $3.63 \pm 0.39$  L O<sub>2</sub>·min<sup>-1</sup>, reached at  $276 \pm 31$  W, with heart rate (HR) =  $187 \pm 7$  bpm and plasma lactate concentration =  $12.4 \pm 3.3$  mmol·L<sup>-1</sup>.

### 2.4. O<sub>2</sub>-Waters Ingested

The waters ingested were prepared from demineralized water, which was remineralized with Na<sup>+</sup>, SO<sub>4</sub><sup>2-</sup> and PO<sub>4</sub><sup>2-</sup>. The control water was enriched in O<sub>2</sub> by injection ( $10$  mg·L<sup>-1</sup>, i.e., close to the value at equilibrium with atmospheric O<sub>2</sub> at sea level at the temperature of ingestion (5–10 °C)). The two other waters were enriched at  $\sim 110$  mg·L<sup>-1</sup> by injection or electrolysis, as previously described [11]. Their O<sub>2</sub> content was found to be

116 and 109 mg·L<sup>-1</sup>, respectively, for the water enriched in O<sub>2</sub> by electrolysis and injection. A very good stability of O<sub>2</sub> was found in the two O<sub>2</sub>-waters. Upon opening and keeping the bottle unagitated at 20 °C, the half-life of the decrease in O<sub>2</sub> content was about six days.

### 2.5. Exercises with Control and O<sub>2</sub>-Waters

The subjects took part in three similar exercise trials by 5- to 7-day intervals. For two days before each trial, the subjects ingested 2 L·day<sup>-1</sup> of the assigned water (~500 mL every 4 h during the wake-up period). The waters were kept in a refrigerator (~5 °C) and were consumed within ~15 min, following removal of the cap. The subjects also ingested 15 mL·kg<sup>-1</sup> of the same water 90 min before the beginning of the exercise trial. Over the two-day period preceding each exercise trial, the subjects refrained from exercising and were fed a standardized diet (35 kcal·kg<sup>-1</sup>·day<sup>-1</sup> with 15, 35 and 50% energy from protein, fat and carbohydrate).

During each exercise trial, following a 20-min warm-up and a 5-min rest period, the subjects exercised at P<sub>max</sub> to volitional fatigue. Breath-by-breath gas exchanges were measured for the computation of ventilation (VE), breathing frequency (f<sub>R</sub>), tidal volume (VT), VO<sub>2</sub>, VCO<sub>2</sub>/VO<sub>2</sub> (RER) and VE/VO<sub>2</sub>. The HR was monitored from a 3-lead ECG (Nihon Kohden TEC-5500, Tokyo, Japan), and cardiac output (Q<sub>c</sub>) was continuously monitored by impedancemetry (Physioflow, Manatec, France). The stroke volume and arterio-venous difference in O<sub>2</sub> were computed. Finally, 100 µL blood samples were withdrawn from the arterialized earlobe at rest before exercise and every minute during the exercise period for the measurement of plasma lactate concentration and pH, arterial oxygen and carbon dioxide pressure (PaO<sub>2</sub> and PaCO<sub>2</sub>) and arterial oxygen saturation (SaO<sub>2</sub>) (Bayer, Bayer series 800, Bayer France; Instrumentation Laboratory, Company, Bedford, MA, USA). PO<sub>2</sub> measured in blood sampled at the arterialized earlobe was corrected, as previously suggested [27] (PaO<sub>2</sub> [in mmHg] = 1.1 × earlobe PO<sub>2</sub> – 3).

### 2.6. VO<sub>2</sub> Kinetics

The time course of the VO<sub>2</sub> response at the onset of exercise was described by using a three-component model [28]. Due to methodological considerations, the cardiodynamic phase was excluded by removing the data points in the first 20 s from the analysis [29]. We used a mathematical model with two exponential functions [30]. The parameters of the model were determined using an iterative procedure which minimizes the sum of the mean squares of the differences between VO<sub>2</sub> estimates and the corresponding actual values as previously described [31]. The amplitude of the slow component was computed as suggested by Borrani et al. [32]. Aberrant breaths were excluded from the analysis, as previously described [33]. Less than 1% of the data points were excluded.

### 2.7. ROS Generation and Damages


Blood samples (6 mL) were taken at rest before exercise in an antecubital vein, following ingestion of the waters. These samples were assessed for estimating oxidative stress on lipid species (blood malondialdehyde (MDA) concentration [34]), protein species (preserved thiol groups [34]) and on endogenous antioxidant defenses (reduced and oxidized glutathione [35]; and on DNA [36]).

## 2.8. Statistical Analyses


Data are reported as mean  $\pm$  standard deviation (SD). The normality of distribution was verified by using the Shapiro–Wilk test. Comparisons were made by using ANOVA for repeated measurements (one- or two-way [Time; Water], depending on the variable; see results) and Tukey HSD at  $p < 0.05$  (SAS Software version 9.1, SAS Institute, Cary, NC, USA). The effect of time was tested by comparing the values observed before exercise to those observed at min 5:00 and the end of the exercise period, except for RER, for which the comparison was made between the final value and the peak value observed at min 3:30. A two-sided level of 5% for the type 1 error was applied. A significance level between 5 and 10% was considered as a trend.

## 3. Results

Figures 1 and 2 show the kinetics of VE,  $\dot{V}O_2$ , VE/ $\dot{V}O_2$ , lactate concentration and of the circulatory adjustments in response to exercise, respectively. Table 2 shows the values of PaO<sub>2</sub>, PaCO<sub>2</sub>, SaO<sub>2</sub> and pH observed at rest before exercise, at min 5 during the exercise and at the end of exercise.



**Figure 1.** Pulmonary responses during exercise: (a) pulmonary ventilation (VE), (b) oxygen uptake ( $\dot{V}O_2$ ), (c) ventilatory equivalent of oxygen and (d) plasma lactate concentration with the three waters (mean  $\pm$  SD; SD not shown past min 10 because  $n < 5$ ; no significant difference was observed between the three waters,  $p > 0.05$ ).



**Figure 2.** Circulatory responses during exercise: (a) heart rate, (b) stroke volume (SV), (c) cardiac output and (d) arterio-venous difference in O<sub>2</sub> (DavO<sub>2</sub>) with the three waters (mean  $\pm$  SD; SD not shown past min 10 because  $n < 5$ ; no significant difference was observed between the three waters,  $p > 0.05$ ).

**Table 2.** Blood gases and pH.

	Water	Rest	Min 5	End of Exercise
PaO <sub>2</sub> (mmHg)	Control	91.1 $\pm$ 8.8	86.4 $\pm$ 6.4 <sup>a</sup>	86.1 $\pm$ 7.4 <sup>a</sup>
	Injection	93.1 $\pm$ 9.1	84.4 $\pm$ 6.7 <sup>a</sup>	84.6 $\pm$ 8.5 <sup>a</sup>
	Electrolysis	94.9 $\pm$ 6.8	86.1 $\pm$ 4.7 <sup>a</sup>	83.5 $\pm$ 5.0 <sup>a</sup>
PaCO <sub>2</sub> (mmHg)	Control	37.5 $\pm$ 1.9	30.7 $\pm$ 3.3 <sup>a</sup>	27.3 $\pm$ 4.3 <sup>a,b</sup>
	Injection	36.7 $\pm$ 2.2	31.0 $\pm$ 3.5 <sup>a</sup>	27.6 $\pm$ 4.3 <sup>a,b</sup>
	Electrolysis	37.3 $\pm$ 2.3	30.6 $\pm$ 3.1 <sup>a</sup>	27.9 $\pm$ 4.0 <sup>a,b</sup>
SaO <sub>2</sub> (%)	Control	95.7 $\pm$ 1.2	95.0 $\pm$ 1.4	94.3 $\pm$ 1.3
	Injection	95.5 $\pm$ 1.0	94.6 $\pm$ 0.8	94.0 $\pm$ 1.5
	Electrolysis	96.1 $\pm$ 1.1	95.0 $\pm$ 0.8	93.9 $\pm$ 1.0
pH	Control	7.40 $\pm$ 0.03	7.40 $\pm$ 0.03 <sup>a</sup>	7.26 $\pm$ 0.05 <sup>a,b</sup>
	Injection	7.39 $\pm$ 0.04	7.30 $\pm$ 0.03 <sup>a</sup>	7.26 $\pm$ 0.04 <sup>a,b</sup>
	Electrolysis	7.40 $\pm$ 0.03	7.30 $\pm$ 0.03 <sup>a</sup>	7.25 $\pm$ 0.04 <sup>a,b</sup>

Partial pressure of O<sub>2</sub> and CO<sub>2</sub>, hemoglobin saturation (SaO<sub>2</sub>) and pH in arterialized blood at rest before exercise, at min 5 during the exercise and at the end of exercise with the three waters (mean  $\pm$  SD; <sup>a</sup> significantly different from rest; <sup>b</sup> significantly different from min 5,  $p < 0.05$ ; comparisons with two-way ANOVA for repeated measurements).

As expected, in response to high-intensity exercise continued to volitional fatigue, following an early response observed between min 0 and min ~5, some variables levelled off (VT [data not shown], VE/VO<sub>2</sub>, plasma lactate concentration, pH and PaO<sub>2</sub>), while others significantly drifted upwards (fR and RER [data not shown], and VCO<sub>2</sub>, VE, VO<sub>2</sub>,



HR and  $Q_c$ ) or downwards ( $PaCO_2$  and  $SaO_2$ ) (Figures 1 and 2). No significant difference was observed between the three waters for any of the variables measured at any time points.

No significant difference was observed for any of the parameters of the  $VO_2$  kinetics between the three waters. However, there was a trend for the time constant of the fast component of the kinetics of  $VO_2$  ( $\tau_1$ ) to be different with the three waters ( $p = 0.060$ ) due to the large difference between the values computed with the control water and the water enriched in  $O_2$  by electrolysis (Tukey HSD,  $p = 0.0513$ ; the corresponding  $p$ -values for the comparison between the control water and the water enriched by injection, and between the two waters enriched in  $O_2$  were much larger, at 0.668 and 0.272, respectively) (Table 3).

**Table 3.** Pulmonary  $VO_2$  kinetics responses.

	Control	Injection	Electrolysis	$p$
$td_1$ (s)	$4.0 \pm 9.8$	$4.9 \pm 9.4$	$5.1 \pm 13.1$	0.886
$\tau_1$ (s)	$46.0 \pm 15.4$	$43.5 \pm 16.1$	$38.8 \pm 16.8$	0.060
$A_1$ (mL $O_2 \cdot \text{min}^{-1}$ )	$2.98 \pm 0.44$	$2.95 \pm 0.40$	$2.91 \pm 0.362$	0.581
$td_2$ (s)	$208.5 \pm 85.0$	$209.1 \pm 82.3$	$189.9 \pm 88.2$	0.643
$\tau_2$ (s)	$285.3 \pm 96.0$	$254.6 \pm 79.8$	$281.5 \pm 101.5$	0.266
$A'_2$ (mL $O_2 \cdot \text{min}^{-1}$ )	$0.46 \pm 0.18$	$0.53 \pm 0.24$	$0.51 \pm 0.20$	0.773

Mean  $\pm$  SD and  $p$ -values for the comparisons with one-way ANOVA for repeated measurements;  $td_1$  and  $td_2$ , and  $\tau_1$  and  $\tau_2$  are respectively the time constants and the time delays for the fast and slow components of  $VO_2$  kinetics;  $A_1$  is the asymptotic amplitude of the fast component of  $VO_2$  kinetics; and  $A'_2$  is the amplitude of the slow component of  $VO_2$  kinetics computed as suggested by Borrani et al. [32].

No significant difference was observed between exercise time to exhaustion with ingestion of the three waters (in min:s:  $10:28 \pm 3:52$ ,  $9:28 \pm 3:42$ , and  $9:54 \pm 3:38$  with control water and the waters enriched in  $O_2$  by injection and the electrolytic process), and the values in the three trials were closely correlated (injection vs. control: Pearson  $r = 0.825$ ; electrolysis vs. control: Pearson  $r = 0.847$ ; electrolysis vs. injection: Pearson  $r = 0.875$ ;  $p < 0.001$  for the three correlation coefficients).

No significant difference was observed between the three waters for any of the indices of ROS generation and damages at rest before exercise (Table 4).

**Table 4.** Indices of ROS and damages in venous blood samples taken at rest.

	Control	Injection	Electrolysis	$p$
Blood MDA content ( $\mu\text{mol} \cdot \text{L}^{-1}$ )	$3.09 \pm 0.37$	$3.11 \pm 0.37$	$3.06 \pm 0.43$	0.682
Preserved thiol ( $\mu\text{mol} \cdot \text{g protein}^{-1}$ )	$6.22 \pm 0.34$	$6.24 \pm 0.42$	$6.22 \pm 0.46$	0.895
Oxidized glutathione ( $\mu\text{mol} \cdot \text{L}^{-1}$ )	$10.1 \pm 5.1$	$10.7 \pm 8.1$	$11.1 \pm 12.7$	0.758
Reduced glutathione ( $\mu\text{mol} \cdot \text{L}^{-1}$ )	$933 \pm 266$	$869 \pm 137$	$937 \pm 444$	0.856
DNA damage (% tail) with FPG	$4.69 \pm 1.09$	$4.77 \pm 1.14$	$4.71 \pm 1.19$	0.600
without FPG	$3.02 \pm 0.74$	$2.85 \pm 0.85$	$3.01 \pm 0.70$	0.140

Mean  $\pm$  SD and  $p$ -values for the comparisons with one-way ANOVA for repeated measurements). MDA, malondialdehyde; FPG, formamidopyrimidine DNA glycosylase (DNA repair enzyme).

#### 4. Discussion

The results from the present experiment indicate that ingestion of waters enriched in  $O_2$  by injection or electrolysis at  $\sim 11$  times the level in the control water did not modify the cardiopulmonary response to high-intensity exercise, the response of plasma lactate concentration and pH, or the response of arterial blood gases. No significant difference was observed between the three waters for the parameters of the kinetics of  $VO_2$ ; however, there was a trend for the fast component to be slightly faster with the water enriched in  $O_2$  by the electrolytic process. Finally, ingestion of waters enriched in  $O_2$  does not increase time

to exhaustion at high exercise intensity, but it also has no adverse effect on the production of ROS and associated damages to lipids, proteins or DNA measured at rest.

These observations are difficult to compare with data in the literature; however, a dozen studies have described the effect of O<sub>2</sub>-water consumption on the response to exercise (Table 1). In only two of these studies [2,7], the O<sub>2</sub> content, which was reported or was computed from the PO<sub>2</sub> reported (see Box 1), was higher than the minimal value for which benefits have been claimed in terms of physiological response and performance (about >7 times the value in tap water) and were credible. In three of the ten other studies listed in Table 1, the O<sub>2</sub> content of the water ingested was simply not reported [18,20,23]. In four other studies [8,17,21,24], there was no evidence that the O<sub>2</sub> content was actually measured, and the values reported are much too high to be credible: the O<sub>2</sub> content reported would require PO<sub>2</sub> in excess of ~900 to ~10,000 atmospheres. Finally, in the three studies listed, in which the O<sub>2</sub> content in commercially available O<sub>2</sub>-waters was actually measured [4,19,37], its value was found to be much lower than advertised by the manufacturer and similar to or only slightly above the value at equilibrium at sea level (see Box 1). It is also worth mentioning that, in only one of the five brands of O<sub>2</sub>-waters analyzed by Hampson et al. [2], the PO<sub>2</sub> corresponded to an O<sub>2</sub> content higher than ~4× the content at equilibrium at sea level. Taken together, results from 10 of the 12 studies available in the literature do not allow us to conclude about putative effects of O<sub>2</sub>-waters on the response to exercise and on performance, because the O<sub>2</sub>-content cannot be ascertained, is incredibly high, or is much too low.

As already discussed by several authors [3,4,7], when compared to the VO<sub>2</sub> at rest and even more during exercise, the amount of O<sub>2</sub> which could be supplied to the blood in the gut by O<sub>2</sub>-waters is extremely small, and unlikely to directly increase aerobic energy production at rest and even less during exercise. In addition, as observed by Fleming et al. [8], a putative increase in VO<sub>2</sub> in peripheral tissues because of an additional supply of O<sub>2</sub> from ingested O<sub>2</sub>-water will not be detected from VO<sub>2</sub> measurement at the mouth which only tracks pulmonary gas exchanges. It is, thus, not surprising that, in previous studies [2,7], as well as in the present experiment, for a given workload, the VO<sub>2</sub> was similar between the three waters. The only previous significant differences reported were a slightly higher VE/VO<sub>2</sub> at submaximal workload and a higher plasma lactate concentration at maximal exercise [7]. However, the differences were low, and the authors questioned their physiological significance. In the present experiment, O<sub>2</sub>-waters with an O<sub>2</sub> content well above the range of values for which manufacturers claim benefits in terms of physiological response and performance did not have any effect on any of the variables measured. Taken together, these data question the interest to use O<sub>2</sub>-water as an ergogenic aid to improve aerobic performance.

There was, however, a trend for the water enriched in O<sub>2</sub> by electrolysis to speed up the kinetic of VO<sub>2</sub> in response to high-intensity exercise when compared with the control water. Since the fast component of the kinetics of VO<sub>2</sub> reflects O<sub>2</sub> diffusion from the blood to muscle fibers [15], this observation is consistent with our previous study showing a lower apparent Km of the mitochondria for O<sub>2</sub> in a solution enriched in O<sub>2</sub> by electrolysis, without any change in the Km for ADP in rat permeabilized muscle fibers [10]. The faster diffusion of O<sub>2</sub> to the mitochondria, as well as the trend to faster adjustment of VO<sub>2</sub> at the onset of exercise observed in the present experiment, in turn, is consistent with the hypothesis that the electrolytic process could generate supramolecular water structures, similar to clathrates [12–14], which can trap O<sub>2</sub> molecules and could modify the local pressure/content relationship for O<sub>2</sub>.

Finally, it has been suggested that administering O<sub>2</sub> internally by ingesting O<sub>2</sub>-waters, which increases PO<sub>2</sub> in the portal vein in rats [38], can be a safety issue because of possible damage to the liver and generation of oxygen radicals [16]. Previous studies showed that acute ingestion of O<sub>2</sub>-waters (300 mL with O<sub>2</sub> content ranging from 30 to ~200 mg·L<sup>-1</sup>) transiently increased blood ascorbyl radical concentration [5,6]. This effect was attenuated following chronic ingestion (0.9 to 1.5 L·day<sup>-1</sup> for 21 to 28 days). Moreover, ingestion of

an O<sub>2</sub>-water for eight days significantly increased serum lipid peroxide concentration and urine alkenyl excretion [17]. However, as already discussed, the actual O<sub>2</sub> content in the water ingested in this study cannot be ascertained. In the present experiment, ingestion of O<sub>2</sub>-water did not result in ROS-induced damage at rest. In addition, the comet assay, which is very sensitive to detect oxidative DNA damages, did not show any genotoxicity of the O<sub>2</sub>-waters administered. A similar result was reported 30 and 60 min following ingestion of 500 mL of O<sub>2</sub>-waters with 70 mg O<sub>2</sub>·L<sup>-1</sup> [39]. Taken together, these results suggest that consumption of waters enriched in O<sub>2</sub> does not induce oxidative stress at rest in healthy subjects.

## 5. Conclusions

In response to high-intensity exercise, results from the present study did not show any beneficial effects of water enriched in O<sub>2</sub> by injection or electrolysis on exercise time to exhaustion, on the cardiopulmonary response and on arterial lactate concentration and pH.

**Author Contributions:** Conceptualization and methodology, F.P. and R.R.; investigation, F.N.D., F.P., A.C., E.L., S.D., B.G. and R.R.; formal analysis, F.N.D., F.P., A.C. and R.R.; data curation, F.N.D., F.P., A.C. and R.R.; resources, A.C., E.L., S.D. and B.G.; validation, F.N.D., F.P. and A.C.; visualization, F.N.D., F.P. and A.C.; writing review and editing, F.N.D., F.P., A.C., E.L., S.D., B.G. and R.R.; supervision, F.P., B.G. and R.R.; project administration, R.R.; funding acquisition, F.P. and R.R. All authors have read and agreed to the published version of the manuscript.

**Funding:** The study was financed by an unrestricted grant from Danone Research (Palaiseau, France).

**Institutional Review Board Statement:** All procedures were performed by following the Code of Ethics of the World Medical Association (Declaration of Helsinki). The protocol was approved by the local ethic committee (Comité de Protection des Personnes Est IV, Strasbourg, France, Eudract No. 2008-A01051-54 on 14 October 2008).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data that support the findings of this study are available from the corresponding author upon reasonable request.

**Acknowledgments:** The authors thank the University of Lille which paid the publication costs.

**Conflicts of Interest:** F. Péronnet and R. Richard are occasional consultants for Danone Research. Frédéric N. Daussin, Antoine Charton, Evelyne Lonsdorfer, Stéphane Doutreleau and Bernard Geny do not have any competing interest to declare.

## References

1. Sander, R. Compilation of Henry's law constants (version 4.0) for water as solvent. *Atmos. Chem. Phys.* **2015**, *15*, 4399–4981. [[CrossRef](#)]
2. Hampson, N.B.; Pollock, N.W.; Piantadosi, C.A. Oxygenated Water and Athletic Performance. *JAMA J. Am. Med. Assoc.* **2003**, *290*, 2408–2409. [[CrossRef](#)]
3. Piantadosi, C.A. "Oxygenated" water and athletic performance. *Br. J. Sports Med.* **2006**, *40*, 740–741. [[CrossRef](#)] [[PubMed](#)]
4. Willmert, N.; Porcari, J.P.; Foster, C.; Doberstein, S.; Brice, G. The effects of oxygenated water on exercise physiology during incremental exercise and recovery. *J. Exerc. Physiol.* **2002**, *5*, 16–21.
5. Gruber, R.; Axmann, S.; Schoenberg, M.H. The influence of oxygenated water on the immune status, liver enzymes, and the generation of oxygen radicals: A prospective, randomised, blinded clinical study. *Clin. Nutr.* **2005**, *24*, 407–414. [[CrossRef](#)]
6. Schoenberg, M.H.; Hierl, T.C.; Zhao, J.; Wohlgemuth, N.; Nilsson, U.A. The generation of oxygen radicals after drinking of oxygenated water. *Eur. J. Med. Res.* **2002**, *7*, 109–116.
7. Leibetseder, V.; Strauss-Blasche, G.; Marktl, W.; Ekmekcioglu, C. Does Oxygenated Water Support Aerobic Performance and Lactate Kinetics? *Int. J. Sports Med.* **2006**, *27*, 232–235. [[CrossRef](#)] [[PubMed](#)]
8. Fleming, N.; Vaughan, J.; Feeback, M. Ingestion of oxygenated water enhances lactate clearance kinetics in trained runners. *J. Int. Soc. Sports Nutr.* **2017**, *14*, 9. [[CrossRef](#)] [[PubMed](#)]
9. Lascoste, C.; Brunner, S.; Jimenez, L.; Klein, A. Method for Enriching Water with Oxygen by an Electrolytic Process, Oxygen Enriched Water or Beverage and Uses Thereof. U.S. Patent US008709231B2, 29 April 2014.

10. Zoll, J.; Bouitbir, J.; Sirvent, P.; Klein, A.; Charton, A.; Jimenez, L.; Péronnet, F.R.; Geny, B.; Richard, R. Apparent Km of mitochondria for oxygen computed from Vmax measured in permeabilized muscle fibers is lower in water enriched in oxygen by electrolysis than injection. *Drug Des. Dev. Ther.* **2015**, *9*, 3589–3597.
11. Charton, A.; Péronnet, F.; Doutreleau, S.; Lonsdorfer, E.; Klein, A.; Jimenez, L.; Geny, B.; Diemunsch, P.; Richard, R. Effect of administration of water enriched in O<sub>2</sub> by injection or electrolysis on transcutaneous oxygen pressure in anesthetized pigs. *Drug Des. Dev. Ther.* **2014**, *8*, 1161–1167. [[CrossRef](#)]
12. Chaplin, M.F. A proposal for the structuring of water. *Biophys. Chem.* **2000**, *83*, 211–221. [[CrossRef](#)]
13. Ozeki, S.; Otsuka, I. Transient oxygen clathrate-like hydrate and water networks induced by magnetic fields. *J. Phys. Chem. B* **2006**, *110*, 20067–20072. [[CrossRef](#)] [[PubMed](#)]
14. Wu, H.-F.; Chin, C.-C.; Liu, B.-M.; Chen, Y.-C.; Lin, C.-H.; Chang, K.-D.; Lee, Y.-H. Self-assembly formation of the magic ion of (H<sub>2</sub>O)<sub>200+</sub>: Observation of nanoscale cages of oxygenated water clusters induced from iron nanoparticles. *Rapid Commun. Mass Spectrom.* **2011**, *25*, 410–414. [[CrossRef](#)]
15. Wasserman, K. Coupling of external to cellular respiration during exercise: The wisdom of the body revisited. *Am. J. Physiol. Endocrinol. Metab.* **1994**, *266*, E519–E539. [[CrossRef](#)] [[PubMed](#)]
16. Arnaud, M.J. Letter to the Editor - Does oxygenated water support aerobic performance and lactate kinetics? *Int. J. Sports Med.* **2006**, *27*, 759–760. [[CrossRef](#)]
17. Askew, E.W.; Pfeiffer, J.M.; Roberts, D.E.; Reading, J.E.; Ensign, W.Y. Does “activated stabilized oxygen” dissolved in drinking water improve aerobic metabolism at moderate altitude? *Wilderness Environ. Med.* **2001**, *12*, 49.
18. Duncan, J. *Fluid Replacement During Exercise: Psychological, Physiologic and Biochemical Benefits of Oxygenated Enhanced Water*; Texas Woman’s University. Center for Research on Women’s Health: Denton, TX, USA, 1997.
19. Fuller, P.J. The Effects of Activated Stabilized Oxygen on Aerobic Endurance in Division II Collegiate Male Soccer Players. Master’s Thesis, Humboldt State University, Arcata, CA, USA, 2010.
20. Jenkins, A.; Moreland, M.; Waddell, T.B.; Fernhall, B. Effect of oxygenated water on percent oxygen saturation and performance during exercise. *Med. Sci. Sports Exerc.* **2001**, *33*, S167. [[CrossRef](#)]
21. McNaughton, L.R.; Kenney, S.; Siegler, J.; Midgley, A.W.; Lovell, R.J.; Bentley, D.J. The effect of superoxygenated water on blood gases, lactate, and aerobic cycling performance. *Int. J. Sports Physiol. Perform.* **2007**, *2*, 377–385. [[CrossRef](#)]
22. Mielke, M. *Oxygenated Water and Exercise Performance*; Florida Atlantic University: Boca Raton, FL, USA, 2004.
23. Wing-Gaia, S.L.; Subudhi, A.W.; Askew, E.W. Effects of purified oxygenated water on exercise performance during acute hypoxic exposure. *Int. J. Sport Nutr. Exerc. Metab.* **2005**, *15*, 680–688. [[CrossRef](#)] [[PubMed](#)]
24. Zhang, H.; Chunran, L.; Ling, W. A preliminary study of hyperoxic solution taken orally on fatigue induced by exercise and its mechanism. *Med. J. Chin. Peoples Lib. Army* **2005**, *6*.
25. Whitley, E.; Ball, J. Statistics review 4: Sample size calculations. *Crit. Care* **2002**, *6*, 335. [[CrossRef](#)] [[PubMed](#)]
26. Jones, N.L.; Summers, E.; Killian, K.J. Influence of age and stature on exercise capacity during incremental cycle ergometry in men and women. *Am. Rev. Respir. Dis.* **1989**, *140*, 1373–1380. [[CrossRef](#)] [[PubMed](#)]
27. Aguilaniu, B.; Maitre, J.; Diab, S.; Perrault, H.; Péronnet, F. Detection of disturbances in pulmonary gas exchanges during exercise from arterialized earlobe PO<sub>2</sub>. *Respir. Physiol. Neurobiol.* **2011**, *177*, 30–35. [[CrossRef](#)]
28. Barstow, T.J.; Molé, P.A. Linear and nonlinear characteristics of oxygen uptake kinetics during heavy exercise. *J. Appl. Physiol.* **1991**, *71*, 2099–2106. [[CrossRef](#)] [[PubMed](#)]
29. Whipp, B.J.; Ward, S.A.; Lamarra, N.; Davis, J.A.; Wasserman, K. Parameters of ventilatory and gas exchange dynamics during exercise. *J. Appl. Physiol.* **1982**, *52*, 1506–1513. [[CrossRef](#)] [[PubMed](#)]
30. Daussin, F.N.; Zoll, J.; Dufour, S.P.; Ponsot, E.; Lonsdorfer-Wolf, E.; Doutreleau, S.; Mettauer, B.; Piquard, F.; Geny, B.; Richard, R. Effect of interval versus continuous training on cardiorespiratory and mitochondrial functions: Relationship to aerobic performance improvements in sedentary subjects. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2008**, *295*, R264–R272. [[CrossRef](#)]
31. Efron, B.; Tibshirani, R.J. *An Introduction to the Bootstrap*; CRC Press LLC: Boca Raton, FL, USA, 1994.
32. Borrani, F.; Candau, R.; Millet, G.Y.; Perrey, S.; Fuchslocher, J.; Rouillon, J.D. Is the VO<sub>2</sub> slow component dependent on progressive recruitment of fast-twitch fibers in trained runners? *J. Appl. Physiol.* **2001**, *90*, 2212–2220. [[CrossRef](#)] [[PubMed](#)]
33. Lamarra, N.; Whipp, B.J.; Ward, S.A.; Wasserman, K. Effect of interbreath fluctuations on characterizing exercise gas exchange kinetics. *J. Appl. Physiol.* **1987**, *62*, 2003–2012. [[CrossRef](#)] [[PubMed](#)]
34. Richard, M.J.; Portal, B.; Meo, J.; Coudray, C.; Hadjian, A.; Favier, A. Malondialdehyde kit evaluated for determining plasma and lipoprotein fractions that react with thiobarbituric acid. *Clin. Chem.* **1992**, *38*, 704–709. [[CrossRef](#)] [[PubMed](#)]
35. Akerboom, T.P.M.; Sies, H. Assay of glutathione, glutathione disulfide, and glutathione mixed disulfides in biological samples. *Methods Enzymol.* **1981**, *77*, 373–382. [[PubMed](#)]
36. Hininger, I.; Chollat-Namy, A.; Sauvaigo, S.; Osman, M.; Faure, H.; Cadet, J.; Favier, A.; Roussel, A.-M. Assessment of DNA damage by comet assay on frozen total blood: Method and evaluation in smokers and non-smokers. *Mutat. Res./Genet. Toxicol. Environ. Mutagenesis* **2004**, *558*, 75–80. [[CrossRef](#)] [[PubMed](#)]
37. Mielke, M.; Kroy, J.O.; Torok, D.; Zoeller, R. Oxygenated Water Does Not Improve Endurance Exercise Performance. *Med. Sci. Sports Exerc.* **2005**, *37*, S42–S43. [[CrossRef](#)]

- 
38. Gelman, S.I. The effect of enteral oxygen administration on the hepatic circulation during halothane anaesthesia: Experimental investigations. *Br. J. Anaesth.* **1975**, *47*, 1253–1259. [[CrossRef](#)] [[PubMed](#)]
  39. Speit, G.; Schütz, P.; Trenz, K.; Rothfuss, A. Oxygenated water does not induce genotoxic effects in the comet assay. *Toxicol. Lett.* **2002**, *133*, 203–210. [[CrossRef](#)]