



**HAL**  
open science

## Psychiatric hospitalisations for people who are incarcerated, 2009–2019: An 11-year retrospective longitudinal study in France

Thomas Fovet, Christine Chan-Chee, Maëlle Baillet, Mathilde Horn, Marielle Wathelet, Fabien d'Hondt, Pierre Thomas, Ali Amad, Antoine Lamer

### ► To cite this version:

Thomas Fovet, Christine Chan-Chee, Maëlle Baillet, Mathilde Horn, Marielle Wathelet, et al.. Psychiatric hospitalisations for people who are incarcerated, 2009–2019: An 11-year retrospective longitudinal study in France. *EClinicalMedicine*, 2022, 46, pp.101374. 10.1016/j.eclinm.2022.101374 . hal-03665662

**HAL Id: hal-03665662**

<https://hal.univ-lille.fr/hal-03665662v1>

Submitted on 22 Jul 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Psychiatric hospitalisations for people who are incarcerated, 2009-2019: an 11-year retrospective**  
2 **longitudinal study in France**

3

---

4 Thomas Fovet<sup>1,2\*</sup>, MD, PhD, Christine Chan-Chee<sup>3</sup>, MD, Maëlle Baillet<sup>4,5</sup>, MD,  
5 Mathilde Horn<sup>1</sup>, MD, PhD, Marielle Wathelet<sup>1,2,6</sup>, MD, Fabien D'Hondt<sup>1,2</sup>, PhD,  
6 Pierre Thomas<sup>1</sup>, MD, PhD, Ali Amad<sup>1</sup>, MD, PhD, Antoine Lamer<sup>4,5</sup>, PhD

7

8 1 – Univ. Lille, Inserm, CHU Lille, U1172 - Lille Neuroscience & Cognition, F-59000 Lille, France

9 2 – Centre national de ressources et de résilience Lille-Paris (CN2R), F-59000 Lille, France

10 3 - National Public Health Agency (Santé Publique France), F-94415 Saint-Maurice, France

11 4 – Univ. Lille, Faculté Ingénierie et Management de la Santé, F-59000, Lille, France

12 5 – Univ. Lille, CHU Lille, ULR 2694 - METRICS: Évaluation des Technologies de santé et des Pratiques  
13 médicales, F-59000, Lille, France

14 6 – Fédération régionale de recherche en psychiatrie et santé mentale, Hauts-de-France

15

16

17 \* Corresponding author

18 Dr. Thomas FOVET

19 Unité hospitalière spécialement aménagée

20 Chemin du Bois de l'Hôpital

21 59113 SECLIN, FRANCE.

22

23 E-mail address: thomas.fovet@chru-lille.fr

24 Tel: +33 3 61 76 30 02

25 **WORD COUNT**

26 4778 words/36 references/3 Tables/3 Figures/3 Supplementary Figures

27

1 **Summary**

2 **Background**

3 Despite the poor mental health status of people who are incarcerated, few studies have examined  
4 the number of psychiatric hospitalisations in this population. Since 2010, France has progressively  
5 opened nine full-time inpatient psychiatric wards exclusively for people who are incarcerated, called  
6 “specially adapted hospital units” (*unités hospitalières spécialement aménagées*, UHSAs, 440 beds).  
7 This study aimed to present the annual rates of psychiatric hospitalisations and primary psychiatric  
8 diagnoses among people who are incarcerated in France from 2009 to 2019.

9 **Methods**

10 We used discharge reports from the French national hospital database to describe longitudinal  
11 retrospective administrative data of psychiatric hospitalisations for people in jail and prison between  
12 2009 and 2019, the age, sex, and principal diagnoses of these patients, the proportion of voluntary  
13 versus involuntary care, and the interactions between UHSAs and other facilities.

14 **Findings**

15 Between Jan 1, 2009, and Dec 31, 2019, 32,228 (92.2% men, n=29,721; 7.8% women, n=2 507)  
16 people who are incarcerated were hospitalised for psychiatric care (64,481 stays). The main  
17 diagnoses were psychotic disorders (27.4%), personality disorders (23.2%), and stress-related  
18 disorders (20.2%). The annual number of people who are incarcerated hospitalised in psychiatric care  
19 increased from 3263 in 2009 to 4914 in 2019. The gradual increase in the activity of UHSAs (300  
20 hospitalisations in 2010 versus 3252 in 2019) was not associated with a reduction in the rate of  
21 hospitalisation of people who are incarcerated in local psychiatric hospitals.

22 **Interpretation**

23 The creation of psychiatric hospitals specifically dedicated to the prison population has not stopped  
24 the hospitalisation of people who are incarcerated at psychiatric hospitals. These findings suggest  
25 that access to psychiatric hospitalisation remains problematic for people who are incarcerated in  
26 France.

27 **Funding**

28 There was no funding source for this study.

29 **Keywords**

30 prison; jail; psychiatric hospitalisation; forensic; psychiatric hospital; specially adapted hospital units;  
31 data reuse

32

33

34

## 1 Introduction

2 More than 11 million people are currently incarcerated around the world.<sup>1</sup> The poor mental  
3 health status of this population has been extensively documented, and access to psychiatric care in  
4 correctional facilities is therefore a major public health issue.<sup>2</sup> In France, the situation has been  
5 severely criticised in recent years.<sup>3-5</sup> Indeed, in addition to harsh overcrowding conditions, a high  
6 prevalence of psychiatric disorders has been described in French prisons. A recent study found that  
7 63.3% of 622 newly incarcerated people presented at least one psychiatric or substance use disorder  
8 identified with the Mini International Neuropsychiatric Interview (MINI).<sup>6</sup> Compared with the general  
9 population, affective disorders, anxiety disorders, and psychotic disorders were approximately three  
10 times more frequent, and substance use disorders were eight times more prevalent.<sup>6</sup> Overall, 41.6%  
11 of the sample had two or more psychiatric or substance use disorders.<sup>6</sup> Before this study, Falissard et  
12 al. had already shown that 36% of people in prison have at least one psychiatric illness considered  
13 “marked” to “severe” (i.e., a rating of 5, 6, or 7 on the *Global Clinical Impressions Scale*).<sup>7</sup> Despite this  
14 alarming situation, very few studies have investigated the number of psychiatric hospitalisations in  
15 this population.<sup>8</sup>

16 The legal regulations for psychiatric treatment of people who are incarcerated diagnosed with a  
17 psychiatric disorder are quite different throughout the world.<sup>9</sup> The French mental health care system  
18 for people in prison is based on three levels of care (see **Table 1**) that are the same regardless of the  
19 type of correctional facility. This care is fully affiliated with the public health system.<sup>10</sup> Regarding full-  
20 time psychiatric hospitalisation, only involuntary hospitalisation was possible for people in prison  
21 until recent changes in the law.<sup>11,12</sup> Almost all of these hospitalisations occurred in psychiatric  
22 hospitals without any additional security measures by the police. Occasionally, patients were  
23 referred to maximum-security wards (called *unités pour malades difficiles*, UMDs). This situation was  
24 associated with several issues in psychiatric hospitals, particularly the inappropriate use of isolation  
25 and mechanical restraint.<sup>4,13</sup> In this context, the conditions for psychiatric full-time hospitalisation of

1 people who are incarcerated have undergone significant changes, and new facilities called specially  
2 adapted hospital units (*unités hospitalières spécialement aménagées*, UHSAs) have been created in  
3 recent years.

4 The nine UHSAs that have opened progressively in France since 2010 (with a current total  
5 capacity of 440 beds) are full-time inpatient psychiatric wards exclusively for people who are  
6 incarcerated.<sup>14,15</sup> In these facilities, the prison administration ensures the security of the institution,  
7 manages entry/exit, and coordinates the transfer of patients. The UHSAs were designed both to  
8 reduce admissions of people who are incarcerated to psychiatric hospitals and to allow these people  
9 to be hospitalised with fewer restrictions of human rights. Voluntarily or involuntary hospitalisations  
10 are possible in UHSAs, unlike psychiatric hospitals, where people who are incarcerated can only be  
11 hospitalised involuntarily. Both types of hospitalisations are requested by the psychiatrist working in  
12 the prison. Involuntary hospitalisations are strictly regulated by law and are used when the individual  
13 who is incarcerated requires immediate inpatient care and constant supervision because of a mental  
14 disorder that makes it impossible for the individual to consent and that poses a danger to self or  
15 others.<sup>16</sup>

16 Since the first UHSA was built in 2010 in Lyon, only one cross-sectional study has been conducted  
17 to examine the activity of the UHSAs.<sup>17</sup> This study showed that 4392 people who are incarcerated  
18 (7027 admissions) were hospitalised in psychiatric care in 2016 in France. Importantly, 1944 people  
19 were hospitalised in UHSAs, and 1787 people were still admitted to psychiatric hospitals despite the  
20 creation of these new facilities. However, in the absence of any longitudinal data, it is extremely  
21 difficult to determine the impact of the creation of UHSAs on the proportion of people who are  
22 incarcerated admitted in psychiatric hospitals and other facilities.

23 This 11-year retrospective longitudinal study (2009-2019) aimed to estimate the number of  
24 psychiatric hospitalisations for people who are incarcerated in France, stratified by calendar year and  
25 type of facility. We hypothesised that the creation of UHSAs decreased the admissions of people who

1 are incarcerated to general psychiatric beds. The secondary objectives were to present patients'  
2 characteristics, including age, sex, and clinical diagnoses, stratified by type of facility, across the study  
3 period of 2009-2019.

## 4 **Methods**

### 5 ***Study design***

6 This retrospective study was performed in accordance with the *REporting of studies Conducted*  
7 *using Observational Routinely collected health Data* (RECORD) statement, which completes the  
8 *Strengthening the Reporting of Observational Studies in Epidemiology* (STROBE) statement when  
9 conducting retrospective observational studies.<sup>18,19</sup>

### 10 ***Database***

11 We used the *Programme de médicalisation des systèmes d'information* (PMSI) database, which  
12 includes individual-level data about the date of admission, length of stay, hospital code number,  
13 sector code, and outcome (i.e., discharge, hospital transfer, death) for all inpatient stays in  
14 psychiatric hospitals in France. The sociodemographic data from these records include sex, age, and  
15 place of residence. The principal diagnosis, defined as the main reason for admission, and the  
16 associated diagnoses related to comorbidities are documented in the PMSI database according to the  
17 French version of the International Statistical Classification of Diseases and Related Health Problems,  
18 10th Revision (ICD-10). A unique national identification number for each patient allows the data from  
19 all hospital stays for the same patient to be linked.

20 Ethical approval was not needed for the present study because we had access to an anonymous  
21 administrative database. Moreover, the national French Public Health Agency (*Santé Publique*  
22 *France*) legally allows full access to national hospital discharge databases<sup>20</sup>, including the PMSI, which  
23 is widely used for research purposes.<sup>21</sup> The authors assert that all procedures contributing to this  
24 work comply with the ethical standards of the relevant national and institutional committees on  
25 human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

1

## 2 **Study population**

3 We conducted a retrospective cohort study, and we included data related to all psychiatric  
4 hospitalisations for people who are incarcerated in France between 2009 and 2019. We did not  
5 distinguish between people incarcerated in *jails* (67%), which house people on remand or people  
6 with a residual sentence below two years, and people incarcerated in *prisons* (33%), which house  
7 people with long sentences. In this article, we used the term “prison” to refer to both prisons and  
8 jails.

9 Cases were identified through the following criteria:

- 10 - the hospital code number corresponding to UHSA, UMD, or daytime hospital psychiatric  
11 beds inside the prison (*Service medico-psychologique regional, SMPR*);
- 12 - the specific legal mode used for the involuntary hospitalization of people who are  
13 incarcerated (“D398”);
- 14 - the types of care (full-time or daytime hospitalisation);
- 15 - the specific sector code used for psychiatric facilities in prison (“P”).

16 As no data on level 1 (consultation and ambulatory care) are documented in the PMSI, only data  
17 on levels 2 (daytime hospitalisation in the *Service medico-psychologique regional, SMPRs*) and 3 (full-  
18 time hospitalisation in the psychiatric hospitals, the UHSAs, or the maximum-security units, UMDs)  
19 were included in this study (see **Table 1**). We used the term “psychiatric hospitals” to refer to both  
20 psychiatric hospitals and psychiatric wards of general hospitals.

21 Unlike SMPRs and UHSAs, which can only accommodate people who are incarcerated,  
22 maximum-security units (UMDs) and psychiatric hospitals can accommodate both incarcerated and  
23 nonincarcerated people. For this study, we only included data from people who are incarcerated. As  
24 they are not considered incarcerated in France, people who cannot stand trial due to mental illness

1 and people found not criminally responsible on account of a mental disorder were not included in  
2 the present study.

3 Finally, additional indicators of psychiatric hospitalisations were collected for  
4 nonincarcerated people from the PMSI database for comparison purposes.

## 5 **Variables**

6 We used several variables from the PMSI database: age and sex, date of admission, outcome  
7 (i.e., discharge, hospital transfer, death), principal diagnosis, type of psychiatric facility, and legal  
8 mode (voluntary versus involuntary care).

## 9 **Statistical analysis**

10 Qualitative variables are presented with absolute numbers and percentages. Quantitative  
11 variables are presented as the median and interquartile range (IQR). The results are illustrated using  
12 line charts. The regression lines result from the affine function ( $y = ax + b$ ). These lines minimize the  
13 sum of squared deviations between each point and the line.

14 Data were extracted from the database with *SAS Enterprise Guide 7.1* software, and analyses  
15 were performed with *SAS Enterprise Guide* and R version 3.6.3. We performed individual-based  
16 analyses and hospitalisation-based analyses.

### 17 1.1.1. Individual-based analyses

18 First, we graphically present (1) the annual number of people who are incarcerated hospitalised  
19 in psychiatric facilities (UHSAs, psychiatric hospitals, SMPRs, and maximum-security units) from 2009  
20 to 2019; for each year, it was calculated as the number of incarcerated people present on January 1  
21 of the year plus the number of incarcerated people entering a psychiatric facility during the year, (2)  
22 the annual number of nonincarcerated people hospitalised in psychiatric care per year in France from  
23 2009 to 2019; for each year, it was calculated as the number of nonincarcerated people present on  
24 January 1 of the year plus the number of nonincarcerated people entering a psychiatric facility during  
25 the year, and (3) the number of incarcerated people on January 1 of each year, from 2009 to 2019.



1 The annual percentage increase in the number of individuals compared to the number measured in  
2 2009 was computed for the 3 populations.

3 Second, we present age, sex, and the principal ICD-10-coded diagnoses of incarcerated people  
4 hospitalised in psychiatric care. The principal ICD-10-coded diagnosis was collected for each stay;  
5 several primary diagnoses were thus possible for individuals with multiple hospitalisations. As a  
6 result, patients with multiple stays and diverging primary diagnoses were counted more than once  
7 for the analysis of diagnoses. We indicate the number of people with multiple diagnoses.

8 Third, we describe the number of people with 1, 2, 3, and 4 or more hospitalisations. We also  
9 present the distribution of the study population in the different psychiatric facilities (UHSAs,  
10 psychiatric hospitals, SMPRs, and maximum-security units) using a Venn diagram. The latter is based  
11 on all hospitalizations that occurred during the study period (2009-2019).

#### 12 1.1.2. Hospitalisation-based analyses

13 First, we present the number of hospitalisations per year for people who are incarcerated  
14 according to the type of facility: UHSAs, psychiatric hospitals, SMPRs, and maximum-security units  
15 (UMDs) as well as the number of beds available in UHSAs. Importantly, when counting the number of  
16 stays per year, some overlapping stays were taken into account for consecutive years, resulting in a  
17 total per year higher than the number of stays for the study period (11 years). We also measured the  
18 following for the 2009-2019 period:

- 19 - annual rates of hospitalisation among people who are incarcerated (i.e., the annual number  
20 of psychiatric hospitalisations divided by the number of people who are incarcerated on  
21 January 1);
- 22 - annual rates of hospitalisation among people who are incarcerated for each facility (i.e., the  
23 annual number of psychiatric hospitalisations in each facility [UHSAs, psychiatric hospitals,  
24 SMPRs, maximum-security units] divided by the number of people who are incarcerated on  
25 January 1);

- 1 - annual rates of hospitalisation among people who are incarcerated for daytime psychiatric  
2 hospitalisation (SMPRs) (i.e., the annual number of psychiatric daytime hospitalisations  
3 [SMPRs] divided by the number of people who are incarcerated on January 1);
- 4 - annual rates of hospitalisation among people who are incarcerated for full-time psychiatric  
5 hospitalisation (i.e., the annual number of psychiatric full-time hospitalisations [UHSAs,  
6 psychiatric hospitals, maximum-security units] in each facility divided by the number of  
7 people who are incarcerated on January 1);

8 A Mann-Kendall Trend Test was performed to determine whether a trend existed in the four time  
9 series. The ratio of stays per patient was computed each year as the number of annual  
10 hospitalisations divided by the number of patients who had one or more stays during the year.

11 Second, we present the median (IQR) duration of hospitalisation in each facility. The length of  
12 stay for each hospitalisation was calculated as the date of admission to either the date of discharge  
13 or December 31, 2019, if the patient was still hospitalised. We also present the legal mode  
14 (voluntary, involuntary, or both) for hospitalisations in UHSAs. We describe the legal mode (voluntary  
15 versus involuntary) on admissions but also at discharge to identify hospitalisations that switched over  
16 in the middle.

17 Third, we focus on hospitalisations in UHSAs to determine the proportion of hospitalisations in  
18 these facilities directly preceded by a stay in another facility or directly followed by a stay in another  
19 facility.

20

### 21 ***Role of the funding source***

22 There was no funding source for this study. CCC had access to the PMSI database. All authors had  
23 access to aggregated data from this source as well as all other data used in the study. All authors  
24 decided to submit the manuscript for publication.

## 1 Results

2 Between Jan 1, 2009, and Dec 31, 2019, 32,228 people who are incarcerated were hospitalised  
3 for psychiatric care (64,481 stays).

### 4 Individual-based analyses

#### 5 *Longitudinal analysis of the number of incarcerated people hospitalised in psychiatric care*

6 The number of incarcerated people hospitalised in psychiatric care per year in France ranged  
7 from 3263 to 4914 between 2009 and 2019 (see **Supplementary Figure 1**), i.e., an increase of 50.6%  
8 (see **Figure 1**). Over the same period, the number of prisoners and the number of nonincarcerated  
9 people hospitalised in psychiatric care per year increased from 62,252 to 70,059 (i.e., an increase of  
10 12.5%) and from 405,117 to 419,794 (i.e., an increase of 3.6%), respectively (see **Figure 1** and  
11 **Supplementary Figure 1**).

#### 12 *Age, sex, and diagnoses*

13 The age, sex, and diagnoses of the study population are presented in **Table 2**. Among the 32,228  
14 incarcerated people hospitalised in psychiatric care, 92.2% (n = 29,721) were men. The median age at  
15 the first hospitalisation was 32 years (IQR: 25-40), with 31 years (IQR: 25-40) and 34 years (IQR: 26-  
16 44) for men and women, respectively (p<.0001).

17 A psychiatric diagnosis was documented for 26,813 individuals (83.2%). There was no psychiatric  
18 diagnosis for 5415 individuals (16.8%). Among them, the information was missing for 2063  
19 individuals (6.4%), the codes Z00-Z99 (*factors influencing health status and contact with health*  
20 *services*) were used for 3112 individuals (9.7%), and other diagnoses were coded for 240 individuals  
21 (0.7%).

22 The most frequent principal psychiatric diagnoses were schizophrenia and delusional disorders  
23 (27.4%), disorders of adult personality and behaviour (23.2%), and neurotic, stress-related and  
24 somatoform disorders (20.2%). 18.0% of patients had multiple diagnoses.

25

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

## ***Number and type of hospitalisations per patient***

Most incarcerated people (61.4%, n = 19,799) had only one hospitalisation during the study period. The number (%) of people with 2, 3, and 4 or more hospitalisations was 6016 (18.7%), 2585 (8.0%) and 3828 (11.9%), respectively. The maximum number of hospitalisations for a single individual was 51. While most people stayed in only one type of facility, approximately one-fifth of patients were hospitalised in different (at least two) types of facilities (n=6871, 21.4%). The exclusive distribution is described in **Supplementary Figure 2**.

## **Hospitalisation-based analyses**

### ***Longitudinal analysis of the number of hospitalisations according to the type of facility***

Overall, 64,481 stays were analysed. Approximately one-third (28.2 %) of the stays occurred in UHSAs (n = 18,187). Psychiatric hospitals, SMPRs, and maximum-security units (UMDs) accounted for 40.5 % (n = 26,131), 31.1 % (n = 20,066), and 1.6 % (n = 1042) of the other stays, respectively.

**Supplementary Figure 3** represents the number of stays per year according to each psychiatric facility (UHSAs, psychiatric hospitals, SMPRs, and maximum-security units). The number of stays in UHSAs increased from 300 (6.8% of the total number of hospitalisations) in 2010 to 3252 in 2019 (39.7% of the total). This increase corresponds to the progressive opening of these structures between 2009 and 2019 (the number of available beds in UHSAs is presented in **Figure 3**). In contrast, the number of stays in psychiatric hospitals and SMPRs remained constant (but the proportion of hospitalisations in these facilities decreased). The number of stays in maximum-security units (UMDs) remained constant.

Annual rates of hospitalisation (daytime hospitalisation, full-time hospitalisation, total) among people who are incarcerated are presented in **Figure 2**. Annual rates of hospitalisation for each psychiatric facility (UHSAs, psychiatric hospitals, SMPRs, and maximum-security units) are presented in **Figure 3**. While rates of hospitalisation in psychiatric hospitals, SMPR, and maximum-security units remained constant, rates of hospitalisation in USHAs continually increased. This was

1 confirmed by a Mann-Kendall trend test, which reported a trend for UHSA ( $p < 0.001$ ) and no trend  
2 for psychiatric hospitals ( $p = 0.061$ ), SMPRs ( $p = 0.436$ ), and maximum-security units ( $p = 0.310$ ).

3         Between 2009 and 2019, the number of annual stays increased from 4358 to 8187, i.e., an  
4 increase of 87.86%, while the number of annual patients increased from 3263 to 4914, i.e., an  
5 increase of 50.60%. The number of stays increased faster than the number of patients, with the ratio  
6 of stay per patient increasing from 1.3 in 2009 to 1.7 in 2019.

### 7 ***Characteristics of the hospitalisations***

8         The median duration of hospitalisation was 19 (IQR: 7–49) days. The median duration of full-  
9 time stays (UHSA, psychiatric hospitals, and maximum-security units) and daytime hospital stays  
10 (SMPRs) were 14 (IQR: 6-34) and 37 (IQR: 14-104) days, respectively. The median duration of  
11 hospitalisation in UHSAs, psychiatric hospitals, and maximum-security units was 26 (IQR: 14-50) days,  
12 8.8 (IQR: 4-20), and 54.5 (IQR: 8-233) days, respectively. **Table 3** presents the characteristics of the  
13 hospitalisations included in the analysis.

14         On admission to a UHSA, 64% of the stays ( $n=11,638$ ) were voluntary, and 36% ( $n=6535$ )  
15 were involuntary. At discharge, 87.5% of stays ( $n=15,917$ ) were voluntary, and 12.5% ( $n=2248$ ) were  
16 involuntary.

### 17 ***Stays at UHSAs versus other facilities***

18         Between 2009 and 2019, 3,036 stays (16.7%) in UHSAs were directly preceded by a stay in  
19 another facility, with eight (0.3%), 2270 (77.8%), 700 (23.1%) and 58 (1.9%) from another UHSA,  
20 psychiatric hospitals, SMPRs and maximum-security units (UMDs), respectively. Of note, 1340 (7.3%)  
21 stays in the UHSA were directly followed by a stay in another facility, with eight (0.6%), 359 (26.8%),  
22 908 (67.8%), and 65 (4.9%) in another UHSA, psychiatric hospitals, SMPRs and maximum-security  
23 units, respectively.

## 1 **Discussion**

2        Developing systems for tracking the use of involuntary care and psychiatric hospitalisation in  
3        correctional contexts has recently been highlighted as a key factor for understanding the psychiatric  
4        needs of people who are incarcerated.<sup>22</sup> In this study, we describe for the first time a longitudinal  
5        analysis of daytime and full-time psychiatric hospitalisations for individuals who are incarcerated in  
6        France between 2009 and 2019. Our results show that the rate of hospitalisation of people who are  
7        incarcerated substantially increased during this period. Importantly, the creation of new facilities  
8        exclusively dedicated to the full-time psychiatric hospitalisation of people who are incarcerated  
9        (UHSAs) in France did not reduce the hospitalisation of these people in psychiatric hospitals.

10        Linking the use of health care services with the need for care remains difficult. Nevertheless, the  
11        hypothesis of growth in the mental health needs of the prison population to explain the significant  
12        increase in the number of psychiatric hospitalisations for people who are incarcerated must be  
13        discussed. It is important to note that the overall prison population has also grown over this period,  
14        from 62,252 in 2009 to 70,059 in 2019. As the poor state of mental health in French prisons is well  
15        documented,<sup>6,7</sup> this increase has probably been accompanied by an escalation in the need for  
16        psychiatric care in correctional facilities. However, while the prison population has “only” increased  
17        by 13%, the number of people in prison hospitalised in psychiatric care increased by 50% between  
18        2009 and 2019. The increasing prison population cannot, therefore, be considered the sole  
19        explanation for the explosion in the number of psychiatric hospitalisations observed in our study.  
20        This period may have been marked by a deterioration in the mental health status of people in prison,  
21        although this has not been directly studied in our work. The increasing number of hospitalisations  
22        could also be due to an unmet need for psychiatric care, which has been filled by the increase in the  
23        number of beds. Particularly, the possibility for people who are incarcerated to be hospitalised  
24        voluntarily may have allowed access to psychiatric care for people who were not being treated  
25        before the beds were opened. The hypothesis of a demand induced by the development of UHSAs  
26        must also be considered. Based on the widely cited principle of Roemer’s Law, which states that

1 *“hospital beds that are built tend to be used”*<sup>23</sup>, some studies have demonstrated a positive  
2 relationship between hospital bed availability and inpatient hospitalisation rates.<sup>24–26</sup> However, some  
3 conflicting results have been shown regarding this principle<sup>27,28</sup>. In addition, measuring the overall  
4 availability of hospital beds to a specific population is a complex task. Simply counting the number of  
5 beds is not sufficient, and it is essential to consider many factors, such as distance, demand, and  
6 access-related factors<sup>26</sup>; it was not possible to rigorously test these factors in our study.

7 Planning psychiatric hospitalisations of people in prison is challenging.<sup>29</sup> On the one hand, the  
8 care provided to people who are incarcerated must be equivalent to that provided to the general  
9 population.<sup>30</sup> On the other hand, legal and security-relevant aspects should be carefully considered.<sup>31</sup>  
10 The development of UHSAs in France since 2010 aimed to meet these two objectives simultaneously.  
11 By allowing access to voluntary full-time psychiatric hospitalisation, which reached 64% of  
12 admissions in these facilities according to our study, the creation of UHSAs has contributed to  
13 optimizing the equivalence of care for people in prison. Importantly, these secure psychiatric  
14 hospitals were intended to accommodate people who are incarcerated suffering from psychiatric  
15 disorders under conditions that respected their rights.<sup>32</sup> As mentioned above, the psychiatric  
16 hospitalisation of people who are incarcerated in community psychiatric hospitals is associated with  
17 several issues. In particular, the use of unsuitable premises with a high risk of escape sometimes  
18 leads to the use of isolation and mechanical restraint even when they are not clinically justified.<sup>4,13</sup>  
19 Although the quality of care in each type of facility was not investigated in this study, the differences  
20 observed in lengths of stay may reflect the difficulty of providing optimal psychiatric care to people  
21 who are incarcerated in community psychiatric hospitals. Indeed, the median stay was 26 days in  
22 UHSAs compared with 8.8 days in psychiatric hospitals, whereas no major difference was shown in  
23 the diagnoses between the facilities, suggesting that the treatment of people who are incarcerated in  
24 psychiatric hospitals is limited to acute stabilisation. It could be hypothesized that inappropriate  
25 conditions for the full-time hospitalisation of people who are incarcerated in psychiatric hospitals  
26 lead psychiatrists to shorten the duration of stay in these facilities.<sup>32</sup> Another important factor could

1 be the pressure on the non-prison psychiatric system to discharge patients to free up beds.  
2 Importantly, our results show that despite the rapid increase in the number of UHSAs since their  
3 creation in 2010 (9 UHSAs, 440 beds), which has made it possible to accommodate more than 18,000  
4 psychiatric stays, the number of stays in psychiatric hospitals and SMPRs remained constant during  
5 this period.

6 The persistence of a high number of people who are incarcerated admitted to psychiatric  
7 hospitals, despite the creation of UHSAs, is probably a consequence of several factors. First, it could  
8 be explained by an insufficient number of available beds (440) in these new structures. Indeed, if no  
9 bed is available in the neighboring UHSAs, the incarcerated patient is referred to the local psychiatric  
10 hospital, which could explain why the number of individuals who are incarcerated admitted to  
11 general psychiatric settings remained stable over the study period. This observation has led the  
12 French government to consider the construction of three new UHSAs (140 beds) in the coming  
13 years.<sup>33</sup> Second, the referral of people in prison requiring inpatient psychiatric care to local  
14 psychiatric hospitals could also be related to the potential difficulties of access to UHSAs. Indeed, it  
15 has been recently shown that the UHSAs mainly accommodate patients from surrounding prisons<sup>17,34</sup>  
16 and that the geographical distance makes it impossible to rapidly transfer individuals incarcerated in  
17 the most isolated prisons to UHSAs (in emergencies, people requiring full-time psychiatric  
18 hospitalisations are therefore referred to psychiatric hospitals). Although this was not directly  
19 investigated in our study, it is important to note that we identified 3036 stays (16.7%) in UHSAs that  
20 were directly preceded by a stay in another facility. This type of transfer could reflect the difficulties  
21 of direct access to UHSAs. Third, the patients referred to UHSAs may have a clinical profile different  
22 than those referred to other facilities, but our results do not support this assertion, as the main  
23 diagnoses of patients in UHSAs do not differ significantly from those of patients admitted to other  
24 facilities.

25 Regarding daytime hospitals within prisons (SMPRs), their activity remained relatively stable  
26 between 2009 and 2019, apart from a slight transitory decrease at the time of the creation of the



1 first UHSAs (2013). This probably reflects a reorientation of the type of psychiatric care within SMPRs  
2 with the opening of the UHSAs. Indeed, it can be hypothesized that the patients suffering from the  
3 most severe mental illnesses, frequently admitted to SMPRs before the creation of the UHSAs, were  
4 gradually redirected to these more appropriate care facilities. Another hypothesis is that this  
5 decrease, also observed for the number of people (not incarcerated) hospitalised in psychiatric care,  
6 could reflect a change in the PMSI data coding methods. Indeed, some psychiatric care activities  
7 were no longer considered hospitalisations after 2013. The SMPRs are located within prisons and  
8 therefore allow the provision of psychiatric care without transferring the person who is incarcerated.  
9 However, no involuntary care is possible in these facilities because French law does not allow  
10 compulsory psychiatric treatment in prison.

11 Surprisingly, the number of incarcerated people hospitalised in maximum-security units  
12 (UMDs) remained low. Only 826 out of 32,228 patients were admitted to maximum-security units  
13 during the study period, from 84 stays in 2009 to 102 stays in 2019. Maximum-security units are  
14 facilities designed to care for the most difficult patients, especially those with a high risk of  
15 violence.<sup>35</sup> The small number of people who are incarcerated could be explained by the difficulties in  
16 accessing these facilities. Indeed, the ten French maximum-security units can accommodate detained  
17 patients as well as patients transferred from psychiatric hospitals. Furthermore, as observed in our  
18 study, the median length of stay is higher in maximum-security units than in other facilities (although  
19 there are very few data available on this topic).

20 Several limitations of the present work should be acknowledged. First, we only had access to  
21 health data from the PMSI with limited sociodemographic data (only age and sex were available). No  
22 information about the severity of the index offense or the criminal history was available. Second, the  
23 use of administrative data does not allow for any confirmation of the diagnosis or the actual clinical  
24 status of the patient. However, our information is derived from diagnoses based on comprehensive  
25 clinical assessments by psychiatrists in hospital settings, and although diagnoses may vary among  
26 practitioners, high levels of agreement have been found for schizophrenia and bipolar disorders in

1 France.<sup>36</sup> Third, as our study was based on a single database, we cannot exclude variations in the  
2 quality of data coding over the years. Fourth, no outpatient mental health services data were  
3 available for the present study. Fifth, we only had access to French data. As a result, caution is  
4 needed before generalizing the results for other countries.

5 This work paves the way for many avenues of research to explain the trends observed in the  
6 present study. First, future studies should explore whether hospitalisations in USHAs are appropriate  
7 and investigate the quality of care in such facilities compared with psychiatric hospitals. Qualitative  
8 studies on care pathways would help to interpret the findings. Metrics such as suicide rates in prison  
9 could also be used to assess the impact of UHSAs on the mental health of people who are  
10 incarcerated. Second, we focused on hospitalisations for the present study, with little attention given  
11 to outpatient care, which is the mainstay of psychiatric treatment in both community and  
12 correctional settings. Access to outpatient care in French jails and prisons should be examined in  
13 future studies. Third, although it has not been rigorously tested, it seems that UHSAs accommodate  
14 more women proportionally (see **Table 2**). Future work should provide a better understanding of this  
15 finding. Finally, international comparisons regarding psychiatric hospitalisation among people who  
16 are incarcerated are urgently needed to help interpret the results presented in this study.

17 We conducted the first longitudinal study investigating the number of hospitalisations in  
18 psychiatry for people who are incarcerated in France. We showed that the number of people in  
19 prison hospitalised in psychiatric care increased drastically between 2009 and 2019 in France.  
20 Importantly, the creation of psychiatric hospitals specifically dedicated to this population (USHAs)  
21 has not stopped the hospitalisation of people who are incarcerated in psychiatric hospitals. Further  
22 arrangements in the French justice system to optimize access to psychiatric care for people in prison  
23 will need to be developed and investigated in the future.

1

2 **Contributors**

3 TF, CCC, MB and AL participated in the conception and design of the study; CCC participated in the  
4 acquisition of data; CCC, MB and AL performed the analyses and verified the underlying data; and TF  
5 wrote the first draft of the manuscript. All authors participated in the writing and revision of the  
6 successive drafts of the manuscript and approved the final version. All authors had full access to all  
7 the data in the study and accepted responsibility to submit for publication.

8 **Declaration of interests**

9 We declare no competing interests.

10 **Data sharing**

11 This administrative data is only available through request from the French National System of Health  
12 Data (“Système National des Données de Santé”, SNDS), which manages this sensitive information  
13 (<https://www.snds.gouv.fr/SNDS/Accueil>), and cannot be shared.

14 **Funding**

15 There was no funding source for this study.

16 **Acknowledgments**

17 The authors would like to thank Alexis Vanhaesebrouck for his helpful discussions on this study.

## 1 References

- 2 1 Penal Reform International. Global Prison Trends 2020. Penal Reform International. 2020.  
3 <https://www.penalreform.org/resource/global-prison-trends-2020/> (accessed April 28, 2020).
- 4 2 Fazel S, Hayes AJ, Bartellas K, Clerici M, Trestman R. The mental health of prisoners: a review of  
5 prevalence, adverse outcomes and interventions. *Lancet Psychiatry* 2016; **3**: 871–81.
- 6 3 Davidson C. France’s forensic psychiatry provision--is it up to scratch? *Lancet Psychiatry* 2015; **2**:  
7 385–7.
- 8 4 Human Rights Watch. Double Punishment | Inadequate Conditions for Prisoners with Psychosocial  
9 Disabilities in France. Human Rights Watch. 2016; published online April 5.  
10 [https://www.hrw.org/report/2016/04/05/double-punishment/inadequate-conditions-prisoners-](https://www.hrw.org/report/2016/04/05/double-punishment/inadequate-conditions-prisoners-psychosocial-disabilities-france)  
11 [psychosocial-disabilities-france](https://www.hrw.org/report/2016/04/05/double-punishment/inadequate-conditions-prisoners-psychosocial-disabilities-france) (accessed July 13, 2018).
- 12 5 Lancet T. Mental health care in French prisons. *The Lancet* 2016; **387**: 1592.
- 13 6 Fovet T, Plancke L, Amarié A, *et al.* Mental disorders on admission to jail: A study of prevalence  
14 and a comparison with a community sample in the north of France. *European Psychiatry* 2020; **63**.  
15 DOI:10.1192/j.eurpsy.2020.38.
- 16 7 Falissard B, Loze J-Y, Gasquet I, *et al.* Prevalence of mental disorders in French prisons for men.  
17 *BMC Psychiatry* 2006; **6**: 33.
- 18 8 Ministère de la Justice. La mesure de l’incarcération au 1er octobre 2021. 2021  
19 [http://www.justice.gouv.fr/prison-et-reinsertion-10036/les-chiffres-clefs-10041/statistiques-de-](http://www.justice.gouv.fr/prison-et-reinsertion-10036/les-chiffres-clefs-10041/statistiques-de-la-population-detenu-et-ecrouee-33736.html)  
20 [la-population-detenu-et-ecrouee-33736.html](http://www.justice.gouv.fr/prison-et-reinsertion-10036/les-chiffres-clefs-10041/statistiques-de-la-population-detenu-et-ecrouee-33736.html).
- 21 9 Fovet T, Horn M, Lancelevée C, Thomas P, Amad A. Language bias in forensic psychiatry research:  
22 the tower of Babel problem. *Acta Psychiatrica Scandinavica* 2020; **142**: 257–8.
- 23 10 Fovet T, Thibaut F, Parsons A, Salize H-J, Thomas P, Lancelevée C. Mental health and the criminal  
24 justice system in France: A narrative review. *Forensic Science International: Mind and Law* 2020; **1**:  
25 100028.
- 26 11 Code de procédure pénale - Article D398.  
27 [https://www.legifrance.gouv.fr/codes/article\\_lc/LEGIARTI000023410936/](https://www.legifrance.gouv.fr/codes/article_lc/LEGIARTI000023410936/) (accessed Feb 8, 2019).
- 28 12 Fovet T, Bertrand M, Horn M, *et al.* [Involuntary psychiatric care for inmates in France: Only for  
29 ‘dangerous’ patients?]. *Encephale* 2018; **44**: 568–70.
- 30 13 Committee for the Prevention of Torture (CPT). Report to the Government of France on the visit  
31 carried out by the CPT from 28 November – 10 December 2010 [CPT/Inf (2012) 13]. *Yearbook of*  
32 *the European Convention for the Prevention of Torture and Inhuman or Degrading Treatment or*  
33 *Punishment/Annuaire de la convention européenne pour la prévention de la torture et des peines*  
34 *ou traitements inhumain ou dégradants* 2018; : 855–972.
- 35 14 Fovet T, Thomas P, Adins C, Amad A. France’s forensic psychiatry provision: the long and winding  
36 road. *Lancet Psychiatry* 2015; **2**: e20.
- 37 15 Moncany A-H, Dandelot D, Bouchard J-P. Entre détention et psychiatrie, les unités hospitalières  
38 spécialement aménagées (UHSA) pour prendre en charge les personnes détenues dont l’état

- 1       psychique relève d'une hospitalisation. *Annales Médico-psychologiques, revue psychiatrique* 2019;  
2       published online Feb 8. DOI:10.1016/j.amp.2019.01.004.
- 3       16 Article L3214-3 - Code de la santé publique - Légifrance.  
4       [https://www.legifrance.gouv.fr/codes/article\\_lc/LEGIARTI000024316672/2022-02-17](https://www.legifrance.gouv.fr/codes/article_lc/LEGIARTI000024316672/2022-02-17) (accessed  
5       Feb 17, 2022).
- 6       17 Fovet T, Amad A, Horn M, Thomas P, Chan-Chee C. Utilization of Hospital-Level Mental Health  
7       Care Services for Inmates in France: A Transversal Study. *Psychiatr Serv* 2020; **71**: 824–8.
- 8       18 Vandembroucke JP, von Elm E, Altman DG, *et al.* Strengthening the Reporting of Observational  
9       Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med* 2007; **4**: e297.
- 10      19 Benchimol EI, Smeeth L, Guttman A, *et al.* The REporting of studies Conducted using  
11      Observational Routinely-collected health Data (RECORD) statement. *PLoS Med* 2015; **12**:  
12      e1001885.
- 13      20 <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000027830713>. Arrêté du 19  
14      juillet 2013 relatif à la mise en œuvre du Système national d'information interrégimes de  
15      l'assurance maladie.  
16      <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000027830713> (accessed July  
17      29, 2019).
- 18      21 Boudemaghe T, Belhadj I. Data Resource Profile: The French National Uniform Hospital Discharge  
19      Data Set Database (PMSI). *International Journal of Epidemiology* 2017; **46**: 392–392d.
- 20      22 Morris NP, Binder RL. Grave Disability in U.S. Jails and Prisons. *PS* 2021; : appi.ps.202100129.
- 21      23 Shain M, Roemer MI. Hospital Costs Relate to the Supply of Beds. *Journal of Occupational and*  
22      *Environmental Medicine* 1959; **1**: 518.
- 23      24 Ginsburg PB, Koretz DM. Bed Availability and Hospital Utilization: Estimates of the “Roemer  
24      Effect”. *Health Care Financ Rev* 1983; **5**: 87–92.
- 25      25 Kroneman M, Siegers JJ. The effect of hospital bed reduction on the use of beds: A comparative  
26      study of 10 European countries. *Social Science & Medicine* 2004; **59**: 1731–40.
- 27      26 Delamater PL, Messina JP, Grady SC, WinklerPrins V, Shortridge AM. Do More Hospital Beds Lead  
28      to Higher Hospitalization Rates? A Spatial Examination of Roemer’s Law. *PLOS ONE* 2013; **8**:  
29      e54900.
- 30      27 van Doorslaer EKA, van Vliet RCJA. “A built bed is a filled bed?” An empirical re-examination. *Social*  
31      *Science & Medicine* 1989; **28**: 155–64.
- 32      28 Rohrer JE. Supply-Induced Demand for Hospital Care. *Health Serv Manage Res* 1990; **3**: 41–8.
- 33      29 Vogel T, Lanquillon S, Graf M. When and why should mentally ill prisoners be transferred to secure  
34      hospitals: a proposed algorithm. *Int J Law Psychiatry* 2013; **36**: 281–6.
- 35      30 Wilson S. The principle of equivalence and the future of mental health care in prisons. *The British*  
36      *Journal of Psychiatry* 2004; **184**: 5–7.

- 1 31 Konrad N. Ethical issues in forensic psychiatry in penal and other correctional facilities. *Curr Opin*  
2 *Psychiatry* 2010; **23**: 467–71.
- 3 32 de Labrouhe D, Plancke L, Amad A, *et al.* [Hospitalization in French forensic units: Results of a  
4 patient satisfaction survey]. *Rev Epidemiol Sante Publique* 2017; **65**: 285–94.
- 5 33 Évaluation des unités hospitalières spécialement aménagées (UHSA) pour les personnes détenues.  
6 <https://www.igas.gouv.fr/spip.php?article775> (accessed Oct 28, 2021).
- 7 34 Eck M, Plancke L, Horn M, Amad A, Thomas P, Fovet T. [Health care delivery and psychiatric  
8 hospitalizations in the prisons of the North of France: An observational study]. *Rev Epidemiol*  
9 *Sante Publique* 2020; **68**: 273–81.
- 10 35 Bouchard J-P, Brulin-Solignac D, Lodetti C. [Prisoners in units for difficult patients]. *Soins Psychiatr*  
11 2016; : 35–40.
- 12 36 Richieri R, Boyer L, Lancon C. [Analysis of the reliability of diagnostic criteria and classifications in  
13 psychiatry]. *Sante Publique* 2011; **23 Suppl 6**: S31-38.

14

## 15 **Figure Captions**

16

17 **Figure 1. Annual percentage increase (compared to the number measured in 2009) in the number**  
18 **of (1) incarcerated people hospitalised in psychiatric facilities (in blue), (2) nonincarcerated people**  
19 **hospitalised in psychiatric care (in yellow), and (3) incarcerated people (in green) in France (2009-**  
20 **2019).**

21

22 *In all three plots, regression lines [E(percent increase in individuals) = beta\_0 + beta\_1(year)] are*  
23 *depicted as dashed lines.*

24

25


26 **Figure 2. Annual rates of daytime (in yellow) and full-time (in pink) psychiatric hospitalisation**  
27 **(total, in blue) for people who are incarcerated between 2009 and 2019 in France.**


28

29 **Figure 3. Annual rates of psychiatric hospitalisation for people who are incarcerated per type of**  
30 **facility (coloured curves) and the number of beds in UHSAs (grey bars) between 2009 and 2019 in**  
31 **France.**


32 *SMPR, services médico-psychologiques régionaux (daytime hospital psychiatric beds inside the*  
33 *prison); UHSA, unité hospitalière spécialement aménagée (full-time inpatient psychiatric ward*  
34 *exclusively for incarcerated people); UMD, unité pour malades difficiles (maximum-security*  
35 *psychiatric unit); PH, psychiatric hospital.*

36









**Table 1.** Description of the 3 levels of the French mental health care system for people in prison

	Type of psychiatric care	Location	Voluntary* or involuntary** care	French designation
<b>Level 1</b>	Consultation and ambulatory care unit inside the prison	178 prisons (100% of French prisons)	Only voluntary	Health unit in prison <i>Unité sanitaire en milieu pénitentiaire (USMP)</i>
<b>Level 2</b>	Daytime hospital psychiatric beds inside the prison	26 prisons (14% of French prisons)	Only voluntary	Regional medical-psychological service <i>Service medico-psychologique regional (SMPR)</i>
<b>Level 3</b>	Psychiatric full-time hospitalisation	Local psychiatric hospitals	Only involuntary	No specific designation
		9 full-time inpatient psychiatric wards exclusively for incarcerated people (440 beds for incarcerated men, women, and juveniles)	Voluntary or involuntary	Specially adapted hospital unit <i>Unité hospitalière spécialement aménagée (UHSA)</i>
		10 maximum-security psychiatric units designed to accommodate patients (incarcerated or not) who “present such a danger to others that the necessary care, supervision and safety measures can only be carried out in a specific unit”. (620 beds for men/36 beds for women).	Only involuntary	Unit for difficult patients <i>Unité pour malades difficiles (UMD)</i>

\*care undertaken only with the consent of the person being treated

\*\*care undertaken without the consent of the person being treated, only in certain circumstances provided for by French law

**Table 2.** Demographic characteristics and principal diagnoses of individuals in prison hospitalised in psychiatry (2009-2019)

	<b>Total</b> <b>N=32,228</b>	<b>UHSA</b> <b>n=10,619</b>	<b>PH</b> <b>n=16,082</b>	<b>SMPR</b> <b>n=12,785</b>	<b>UMD</b> <b>n=826</b>
Age (median, IQR)	32.0 [25.0-40.0]	32.0 [25.0;40.0]	31.0 [24.0;39.0]	31.0 [25.0;40.0]	31.0 [24.0;39.0]
Sex, Male (n, %)	29 721 (92.2%)	9 647 (90.8%)	15 022 (93.4 %)	11973 (93.6%)	808 (97.8 %)
Sex, Female (n, %)	2 507 (7.8%)	972 (9.2%)	1 060 (6.6%)	812 (6.4%)	18 (2.2%)
<i>Principal diagnosis</i>					
<b>F00-99 Psychiatric disorder (n, %)</b>	2,6813 (83.2%)	9,831 (92.6%)	14,164 (88.1%)	9,889 (77.4%)	725 (87.8%)
F00-09 Organic mental disorders (n, %)	180 (0.6%)	88 (0.8%)	107 (0.7%)	51 (0.4%)	3 (0.4%)
F10-19 Mental and behavioural disorders due to psychoactive substance use (n, %)	3,407 (10.6%)	859 (8.1%)	1478 (9.2%)	1,991 (15.6%)	67 (8.1%)
F20-29 Schizophrenia, and delusional disorders (n, %)	8,818 (27.4%)	4,506 (42.4%)	5,227 (32.5%)	2,932 (22.9%)	463 (56.1%)
F30-39 Mood disorders (n, %)	4,707 (14.6%)	2,160 (20.3%)	2,788 (17.3%)	1,537 (12.0%)	89 (10.8%)
F40-48 Neurotic, stress-related and somatoform disorders (n, %)	6,495 (20.2%)	2,445 (23.0%)	3,627 (22.6%)	2,454 (19.2%)	75 (9.1%)
F60-69 Disorders of adult personality and behaviour (n, %)	7,462 (23.2%)	3,107 (29.3%)	4,482 (27.9%)	2,709 (21.2%)	263 (31.8%)
F70-79 Mental retardation (n, %)	331 (1.0%)	146 (1.4%)	204 (1.3%)	155 (1.2%)	13 (1.6%)
F90-98 Behavioural and emotional disorders with onset in childhood and adolescence (n, %)	389 (1.2%)	119 (1.1%)	238 (1.5%)	173 (1.4%)	15 (1.8%)
F99 Unspecified mental disorder (n, %)	519 (1.6%)	157 (1.5%)	355 (2.2%)	216 (1.7%)	27 (3.3%)

*SMPR, services médico-psychologiques régionaux (daytime hospital psychiatric beds inside the prison); UHSA, unité hospitalière spécialement aménagée (full-time inpatient psychiatric ward exclusively for incarcerated people); UMD, unité pour malades difficiles (maximum-security psychiatric unit); PH, psychiatric hospital.*

**Table 3.** Characteristics of psychiatric hospitalisations for individuals in prison between 2009 and 2019 in France

	<b>Total</b>	<b>UHSA</b>	<b>GPH</b>	<b>SMPR</b>	<b>UMD</b>
Number of stays	64,481	18,187 (28.2%)	26,131 (40.5%)	20,066 (31.1%)	1,042 (1.6%)
Median length of stay (median, IQR)	19 (7–49)	26 (14-50)	8 (4-20)	37 (14-104)	54.5 (8-233)

*SMPR, services médico-psychologiques régionaux (daytime hospital psychiatric beds inside the prison); UHSA, unité hospitalière spécialement aménagée (full-time inpatient psychiatric ward exclusively for incarcerated people); UMD, unité pour malades difficiles (maximum-security psychiatric unit); PH, psychiatric hospital.*