



HAL
open science

Diversity of sialic acids and sialoglycoproteins in gametes and at fertilization

Ingrid Fliniaux, Guillaume Marchand, Caroline Molinaro, Mathieu Decloquement, Alain Martoriati, Matthieu Marin, Jean-Francois Bodart, Anne Harduin Lepers, Katia Cailliau

► **To cite this version:**

Ingrid Fliniaux, Guillaume Marchand, Caroline Molinaro, Mathieu Decloquement, Alain Martoriati, et al.. Diversity of sialic acids and sialoglycoproteins in gametes and at fertilization. *Frontiers in Cell and Developmental Biology*, 2022, 10, pp.982931. 10.3389/fcell.2022.982931 . hal-03857419

HAL Id: hal-03857419

<https://hal.univ-lille.fr/hal-03857419>

Submitted on 17 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Matteo Avella,
Sidra Medicine, Qatar

REVIEWED BY

Keizo Tokuhira,
Kansai Medical University, Japan
Amber R. Krauchunas,
University of Delaware, United States

*CORRESPONDENCE

Katia Cailliau,
katia.maggio@univ-lille.fr

SPECIALTY SECTION

This article was submitted to Molecular and Cellular Reproduction, a section of the journal Frontiers in Cell and Developmental Biology

RECEIVED 30 June 2022

ACCEPTED 10 October 2022

PUBLISHED 20 October 2022

CITATION

Fliniaux I, Marchand G, Molinaro C, Decloquement M, Martoriati A, Marin M, Bodart J-F, Harduin-Lepers A and Cailliau K (2022), Diversity of sialic acids and sialoglycoproteins in gametes and at fertilization.

Front. Cell Dev. Biol. 10:982931.
doi: 10.3389/fcell.2022.982931

COPYRIGHT

© 2022 Fliniaux, Marchand, Molinaro, Decloquement, Martoriati, Marin, Bodart, Harduin-Lepers and Cailliau. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Diversity of sialic acids and sialoglycoproteins in gametes and at fertilization

Ingrid Fliniaux, Guillaume Marchand, Caroline Molinaro, Mathieu Decloquement, Alain Martoriati, Matthieu Marin, Jean-François Bodart, Anne Harduin-Lepers and Katia Cailliau *

Univ. Lille, CNRS, UMR 8576-UGSF-Unité de Glycobiologie Structurale et Fonctionnelle, Lille, France

Sialic acids are a family of 9-carbon monosaccharides with particular physicochemical properties. They modulate the biological functions of the molecules that carry them and are involved in several steps of the reproductive process. Sialoglycoproteins participate in the balance between species recognition and specificity, and the mechanisms of these aspects remain an issue in gametes formation and binding in metazoan reproduction. Sialoglycoproteins form a specific coat at the gametes surface and specific polysialylated chains are present on marine species oocytes. Spermatozoa are submitted to critical sialic acid changes in the female reproductive tract facilitating their migration, their survival through the modulation of the female innate immune response, and the final oocyte-binding event. To decipher the role of sialic acids in gametes and at fertilization, the dynamical changes of enzymes involved in their synthesis and removal have to be further considered.

KEYWORDS

sialic acid, glycoprotein, oocyte, sperm, egg, fertilization

Abbreviations: CAZy, Carbohydrate Active EnZymes database; CMAH, N-acetylneuraminic acid hydroxylase; CMAS or CSS: CMP-sialic acid synthetase; CMP, cytidine monophosphate; GalNAc, N-acetylgalactosamine; GlcNAc, N-acetylglucosamine; Gd, glycodelin; GH33, glycoside hydrolase 33; GNE, glucosamine UDP-GlcNAc-2-epimerase/N-acetylmannosamine kinase; Neu, neuraminic acid; GT29, glycosyltransferase family 29; Kdn, 2-keto-3-deoxy-D-glycero-D-galacto-nonulosonic acid; NANP, N-acetylneuraminic-9-phosphatase; NANS, Neu5Ac-9-P synthase; NEU, neuraminidase; Neu5Ac, N-acetylneuraminic acid; Neu5Gc, N-glycolylneuraminic acid; NCAM, neural cell adhesion molecule; polySias, polysialic acids; PB, polar body; PEP, phosphoenolpyruvate; PSGP, polysialoglycoprotein; Sd^p, antigen, (Neu5Ac_{2,3}(GalNAc β1,4)Galβ1,4GlcNAc); Sia, sialic acid; SLe^x, Sialyl-Lewis(x) (NeuA5ca_{2,3}Galβ1,4(Fuca1,3)GlcNAc); ST8SIA2, Alpha-N-Acetyl-Neuraminidase Alpha-2,8-Sialyltransferase two; ST8SIA4, Alpha-N-Acetyl-Neuraminidase Alpha-2,8-Sialyltransferase four; UDP, uridine diphosphate; WGDR, round of whole genome duplication; ZP, zona pellucida.

1 Introduction

In vertebrate and invertebrate metazoan sexual reproduction, gametes acquire competence for fertilization during oogenesis and spermatogenesis. Before fertilization, both male and female gametes have to go through a long journey in aquatic environments or into the urogenital tracts, and have therefore developed specificity in the envelopes surrounding their plasma membrane. Gametes are surrounded by a specific extracellular matrix that exerts several functions. One function is to protect them from external (freshwater or seawater) or internal (female genital tract: vagina-cervix, uterus, oviduct) environments. This cellular matrix is enriched in negatively charged glycoconjugates, ending with sialic acids (Sias) (Schauer, 2004; Schauer, 2009). Sias predispose gametes to hydration protection, necessary for gametes interaction and successful fertilization (Hirohashi et al., 2008; Li et al., 2011). In the oocyte jelly coat, Sias also play a role in water retention and are a reservoir for several ions including calcium (Bernier and Ingermann, 1990). By virtue of both bulk and negative charge, large acidic polymers of polysialic acids (polySias) modulate negatively cell to cell and cell to matrix interactions (Varki, 1992) and protect gametes before fertilization. Sialylated glycan chains on proteins occupy more space compared to a naked globular protein (Cyster et al., 1991; Li et al., 2011; Bhide and Colley, 2017), and form a cap impairing interaction with plasma membrane conjugates that limit species unspecific fertilization. However, sialoglycoproteins are also involved in specific species recognition between the spermatozoa and the oocyte. Moreover, a balance is maintained between species specificity and immune protection through the occurrence of a spermatozoa dynamical change in the Sias coat in vertebrates and invertebrates. For those reasons, fertilization is more than a complex series of Sias modifications and interactions compared to the general recognition mechanism found in pathogens using Sias residues on their carbohydrates (as for viruses, Glasner et al., 2012; Rosenstock and Kaufmann, 2021).

The present review highlights the diversity and properties of Sias present on glycoproteins to ensure their multiple roles in gametes and at fertilization. To get further insight into the specificity of polySias in both oocytes and spermatozoa, the structure, biosynthesis of glycoproteins, and the phylogenic filiation of sialyltransferases are considered. The presence and the species-specific composition in the surrounding membranes of the oocytes with polySias chains diversity in aquatic species and the specificity of sialylated ZP in the mammalian *zona pellucida* (ZP) are then addressed. A third part is dedicated to the coating and uncoating fine-tuned regulation of Sias in sperm. Remodeling the surrounding coat of sialoglycoproteins finalizes the spermatozoa fertility properties. In addition, specific interactions between sperm Sias and Siglec receptors (sialic acid-binding immunoglobulin-like lectins) (Teclé et al., 2019) occur in the female genital tract of mammals counteracting the

female immune response. In a last part, the final unmasking of Sias from the oocyte coat by the spermatozoa acrosomal reaction to gain accessibility when the gametes encounter is discussed at fertilization.

2 Sias structure and biosynthesis

2.1 Sias structural diversity

Gametes surfaces, like other cell surfaces, are covered by a thick dense coat known as the glycocalyx, formed by a complex network of glycoconjugates including glycoproteins and glycolipids representing up to 30 μm for echinoderm oocytes (Cohen and Varki, 2010; Schjoldager et al., 2020). The terminal position of oligosaccharide chains (i.e. glycans) is occupied by a variety of negatively charged acidic monosaccharides, the Sias, all derived from Neu5Ac (*N*-Acetylneuraminic acid) and Kdn (2-keto-3-deoxynononic acid). Interestingly, these sialylated glycoconjugates are characterized by a high structural and functional diversity, which can be analyzed at different levels (Cohen and Varki, 2010).

A first level of diversification of Sia molecules is due to various substitutions with acetyl, glycolyl, methyl, lactyl, phosphate, or sulfate groups found on Sias hydroxyl or amino groups (Figure 1A) leading to over 50 structurally different Sia derivatives in nature (Schauer, 1982). The most abundant members of the Sia family are Neu5Ac and Neu5Gc (5-*N*-glycolylneuraminic acid), followed by Kdn and Neu (neuraminic acid) (Figure 1B). However, there are marked differences in the distribution of these Sias in nature and some of these Sias completely disappear from a lineage as for Neu5Gc in human and platypus (Schauer et al., 2009; Chen and Varki, 2010). Although the biological significance of these modified Sia residues is still mostly unknown, it is interesting to note that the most Sia diversity is encountered in invertebrate deuterostomes such as echinoderms (sea urchin and starfish) and the simplest profiles are found in human (Gagneux et al., 2017). The natural occurrence of *O*-acetylated Sias is more frequently documented than other modified Sias and *O*-acetylation at C-4, C-7, or C-9 of Sias has been observed in bacteria and eukaryotes including humans (Angata and Varki, 2002; Nguyen et al., 2020). A neuraminic acid molecule may contain one to three *O*-acetyl residues, but *N*-acetyl-9-*O*-acetylneuraminic acid (Neu5,9Ac2) is the predominant form (Schauer and Kamerling, 2011). As for *O*-sulfated Sia (Sias) structures, 8-*O*-sulfation of Neu5Ac and Neu5Gc have been described in sea urchin gametes where they are involved in sperm-egg interaction and in the regulation of sperm motility, but so far almost nothing is known about the occurrence of Sias in vertebrates (Ertunc et al., 2020).

Additional levels of diversity arise from 1) the type of glycosidic α -linkage through which sialic acid moieties are attached to other glycan units, 2) the underlying sugar chains and 3) the type of glycan (Cohen and Varki, 2010). Sias can be




FIGURE 1
 Sialic acids hierarchical complexity in sialoglycoproteins: **(A)** Substitutions found in sialic acid by various types of glycosidic linkages: in R2 an H is present in free sialic acid, or an alpha-2,X linkage to Gal (X = 3/4/6), GalNAc (X = 6), GlcNAc (X = 4/6) or Sia (X = 8/9), R4 and R7 are H or O-acetyl, R5 is hydroxyl, amino, N-acetyl, or N-glycolyl, R8 is an H, O-acetyl, O-methyl, O-sulfate or sialic acid, R9 is an H, O-acetyl, O-lactyl, O-phosphate, O-sulfate or sialic acid. **(B)** Sialic acids basic forms Neu5Ac, Neu5Gc, Neu, and Kdn. **(C)** Terminal alpha 2, X-linkages and patterns. **(D)** Underlying glycoproteins. **(E)** Spatial organization.

linked in $\alpha 2,3$ - and in $\alpha 2,6$ -linkage to galactose (Gal)-containing glycans or in $\alpha 2,6$ -linkage to N-acetylgalactosamine (GalNAc) and N-acetylglucosamine (GlcNAc) residues (Figure 1C). These Sias occur as monosialyl residues at the nonreducing terminal ends of glycan chains of either N- or O-glycoproteins and glycolipids. A very unique feature of Sias is that they can be linked to each other through $\alpha 2,8$ or $\alpha 2,9$ -linkages forming oligo- and polySia chains found on a few glycoproteins that are called

disialic acid (diSia), trisialic acid (triSia), oligosialic acid (oligoSia), and polysialic acid (polySia) according to their degree of polymerization (DP) (Troy, 1992; Sato, 2013; Sato and Kitajima, 2013). These combinations of Sia structures, modifications, DP, linkages, underlying glycans and glycan types lead to a huge diversity of sialoglycans and sialoglyconjugates constituting the so-called sialome (Varki, 2017; Sato and Kitajima 2021).




FIGURE 2 (A) Sialic acids biosynthesis. (B) Sialic acid linkages performed by sialyltransferases. (C) Sialyltransferases with preferred glycoproteins substrates. (D) Counteracting neuraminidases (NEU) detected in male (blue dots) and female gonads (pink dots) and spermatozoa (green dots) of human (H) (Lassalle and Testart, 1994; Ma et al., 2012), and fishes: zebrafish (Z) (Harduin-Lepers et al., 2008; Petit et al., 2018; Yamakawa et al., 2018) and sturgeon (S) (Li et al., 2017).

A final layer of complexity stems from the various glycosylation sites within a given protein (macroheterogeneity and microheterogeneity), and the three-dimensional organization of sialoglycoconjugates on cell surfaces (Figure 1D). Sialoglycoproteins and gangliosides can assemble into micro-domains on cell surfaces leading to the formation of glyco-cluster (Figure 1E) that could mediate differential biological functional events.

2.2 Sialylation machinery

Metabolism of Sias and sialoglycoproteins schematized in Figure 2A divides into several steps diversely localized in the animal cell and a recent phylogenetic analysis reported the double origin of Sias in eukaryotic cells (Petit et al., 2018). An extrinsic Sias source in Metazoa is likely ensured by an intake of Sias from the extracellular space via the lysosomal SLC17A5 transporter also known as sialin associated with neuraminidase 1 (NEU1). The canonical biosynthetic pathway of Sias takes place within the cytosolic compartment of vertebrate cells (Figure 2A). Sias formation is catalyzed by a bifunctional

enzyme, glucosamine UDP-GlcNAc-2-epimerase/*N*-acetylmannosamine kinase (encoded by the human *GNE* gene), which converts UDP-GlcNAc (uridine diphosphate *N*-acetylglucosamine) to ManNAc-6-P (*N*-acetylmannosamine 6-phosphate) and UDP (uridine diphosphate). Then, ManNAc-6-P is condensed with phosphoenolpyruvate (PEP) by the Neu5Ac-9-P synthase (*NANS*) forming *N*-acetylneuraminic acid 9-phosphate (Neu5Ac-9-P). Neu5Ac-9-P is then dephosphorylated by the cytosolic *N*-acetylneuraminic acid 9-phosphatase (*NANP*), resulting in the release of Neu5Ac to the cytoplasm. These two enzymes are widely distributed in most eukaryotic lineages (Petit et al., 2018). In prokaryotes, the condensation of ManNAc (*N*-acetylmannosamine) and PEP is catalyzed by an aldolase. The same pathway can use Man-6-P instead of ManNAc-6-P leading to Kdn.

Subsequent use of Sias depends on their activation to the sugar-nucleotide cytidine monophosphate sialic acid (CMP-sialic acid or CMP-Sia) which is catalyzed by the cytidine monophosphate *N*-acetylneuraminic acid synthetase or CMP-sialic acid synthetase (*CMAS* or *CSS*) in the nucleus of most vertebrate cells (Figure 2A). In Metazoa, orthologous sequences of human *CMAS* gene were retrieved in the Cnidaria,




FIGURE 3

Animal sialyltransferases genes, transcripts, and/or proteins detected in male (blue) and female (pink) gonads, and male (green) and female (yellow) gametes. (1) Asahina et al., 2004 (T); (2) Asahina et al., 2006 (T); (3) Venuto M. T. et al., 2020 (G, T); (4) Kurosawa et al., 1996 (G, T); (5) Krzewinski-Recchi et al., 2003 (G, T); (6) Takashima et al., 2002 (G, T); (7) Yamakawa et al., 2018 (P); (8) Petit et al., 2018 (G); (9) Harduin-Lepers et al., 2008 (G, T); (10) Stern et al., 2000 (G, T, P); (11) Taniguchi et al., 2003 (T); (12) Harduin-Lepers et al., 2005 (G); (13) Lei et al., 2018 (T); (14) Kimura et al., 1999 (T, P); (15) Takashima et al., 2003 (G, T); (16) Simon et al., 2013 (P). Genes (G), proteins (P), related gene (r), transcripts (T), whole genome duplication round (WGDR).

deuterostomes, and Arthropods (Di et al., 2017; Petit et al., 2018). Interestingly, in the zebrafish *Danio rerio*, there are two CSS with different substrate specificity and spatial expression one


localizing in the nucleus and showing higher specificity towards Neu5Ac and the other localizing to the cytosol and showing higher specificity towards Kdn (Schaper et al., 2012).

Neu5Ac can be transformed into Neu5Gc by hydroxylation of the methyl group in the *N*-acetyl moiety of the activated CMP-Neu5Ac donor through a reaction catalyzed by cytidine monophosphate-*N*-acetylneuraminic acid hydroxylase (CMAH) (Shaw and Schauer, 1988). Orthologues of the CMAH gene were found in Porifera (Kapli et al., 2021), and chordates including tunicates, lancelets, reptiles, amphibians, most fish, and a number of mammals (Peri et al., 2018). Only a few deuterostome lineages do not have a functional CMAH gene as reported in humans, New World monkeys, ferrets, several bats, the platypus, the perciform fish *lates calcarifer* (Simakov et al., 2015; Peri et al., 2018), Sauropsids (birds and reptiles), Pinnipedia, and members of Musteloidia (Altman and Gagneux, 2019) that lack Neu5Gc.

The various activated sugar-nucleotide CMP-Sias are then transported from the cytosol to the sialylation site within the Golgi lumen (Lepers et al., 1989; Lepers et al., 1990) by the SLC35A1 antiporter (Figure 2A). SLC35A1-like sequences are found in deuterostomes from cephalochordates to human and in protists further suggesting that it is ancestral in eukaryotes (Petit et al., 2018).

Then sialylation steps and glycosidic linkage formation are carried out by specific α2,3-, α2,6- or α2,8-sialyltransferases confined to the Golgi apparatus that belong to glycosyltransferase family 29 (GT29) according to the Carbohydrate Active EnZymes database (CAZy, <http://www.cazy.org>) (Lombard et al., 2014) (Figure 2B). Four large families of sialyltransferases were described in Metazoan *i.e.* the ST6GAL, ST6GALNAC, ST3GAL, and ST8SIA (Harduin-Lepers et al., 2001; Harduin-Lepers et al., 2005; Harduin-Lepers, 2010; Teppa et al., 2016). A burst of novelties was acquired in each family after the two rounds of whole genome duplication (WGD) at the root of vertebrates leading to several subfamilies that were maintained or lost during vertebrate evolution (Figure 3) (Harduin-Lepers et al., 2008; Petit et al., 2010; Petit et al., 2015). An interesting case study is the one of teleost ST8SIAs that shows an extended repertoire in fish mostly shaped by the WGD events and a particular distribution in fish species (Chang et al., 2019; Venuto M. T. et al., 2020). Moreover, sequence-based analysis suggested that molecular evolution of polysialyltransferase-related sequences could account for the extraordinary diversity in polySias encountered in fish (Venuto M. T. et al., 2020). Additional modifications of Sias like *O*-acetylation are thought to take place in the Golgi apparatus and involve intraluminal *O*-acetyltransferase and acetyl-CoA transporter proteins (Figures 2B,C).

Sialoglycoconjugates exported to the cell surface function as ligands for sialic acid-binding proteins including selectins, a class of calcium-dependent C-type lectins, and Siglecs. Siglecs are Sia



binding immunoglobulin (Ig)-like lectins that belong to the immunoglobulin superfamily and act as transmembrane cell surface immune regulatory receptors predominantly found on hematopoietic cells including B and T cells, natural killer cells, dendritic cells, granulocytes as well as mast cells and macrophages (Varki and Angata, 2006; Crocker et al., 2007). At least 15 human Siglecs are known to be expressed in human tissues which include CD33-related Siglecs and Siglec-1 (Sialoadhesin), Siglec-2 (CD22), Siglec-4 (myelin-associated glycoprotein, MAG), and Siglec-15. Siglec-1, Siglec-2, Siglec-4, and Siglec-15 constitute a group of highly conserved and ancient molecules which could be found in the common ancestor of vertebrates (Bornhöft et al., 2018). The remaining Siglecs including the CD33-related Siglecs show marked inter-species differences in repertoire (a loss of Siglecs genes was reported in rodent), sequence, and binding preference (Angata, 2006).

Finally, sialidases, also known as neuraminidases (NEU genes) release α -linked Sias from glycoconjugates and polysaccharides. These enzymes are common in Metazoa and also in microorganisms. The human enzymes NEU1, NEU2, NEU3 and NEU4 are classified in the glycoside hydrolase family GH33 of the CAZy classification (Henrissat and Bairoch, 1996). They are specific of sialylated glycoconjugates and show characteristic tissue and cell expression pattern (Monti et al., 2010). Interestingly, NEU1 is found in the lysosome of Metazoa where it is involved in the hydrolysis of exogenous sialyloglycoconjugates (Figure 2D) and recent phylogenetic analyses described NEU1 as the most ancient neuraminidase whereas the clade NEU2/NEU3/NEU4 would have another origin in deuterostomes (Giaccopuzzi et al., 2012; Petit et al., 2018).

3 Oocyte Sias diversity and their multiple roles

Deuterostome oocytes are surrounded by a soft gelatinous vitelline envelope and an important jelly coat in echinoderms and amphibians, a thick chorionic membrane in fishes, or a *zona pellucida* in mammals, all composed of numerous different carbohydrates including a high diversity of sialylated glycoproteins (Monné and Jovine, 2011). In a single phylum, the composition of Sias varies among the different species (Figure 4).

3.1 Oocytes from marine and freshwater species

3.1.1 A high content and a variety of sialic acid derivatives surround oocytes

The surrounding transparent gelatinous layer or jelly coat of sea urchin oocytes (up to 30 μ M) is rich in polysaccharides and contains unique sialylated glycoconjugates (Kitazume et al., 1994; Kitazume et al., 1996). Mass spectral analyses determined that Neu5Gc is the major Sia while Neu5Ac is the second most common (respectively 66% and 11%) and 23% of other Sias including Neu5GcS, Neu5Gc9Ac, Neu5AcS, Neu5Gc7,9Ac2, Neu5,9Ac2, Neu5Gc8Ac, Neu5Gc7Ac, Neu5,7Ac2, Neu5Gc8,9Ac2, and Neu5,8Ac2 (Yeşilyurt et al., 2015) (Figure 4).

Amphibian oocytes thick jelly coat contains several layers: three layers in the Anurans *Xenopus laevis* and six in *Rana pipiens* (*Lithobates pipiens*, Taxonomy ID: 8404). In the pipidae *X. laevis*, oocytes Sias are localized at the interface between the first and second layers, charged by oviductal Sias secretions as they go through the oviduct (Mozingo and Hedrick, 1999). The jelly coat contains four types of sialylated glycans with three of them containing Kdn and one Neu5Ac (Plancke et al., 1995), with sulfated (Strecker et al., 1995), or doubly sulfated anionic

glycans (Guerardel et al., 2000). An alteration of the Sias content in the jelly coat leads to inhibition of sperm penetration (Gurushankara et al., 2012). In *Xenopus tropicalis* jelly coat, among the 19 anionic glycans detected, six contain Neu5Ac while *Xenopus borealis* has no detectable anionic glycans and exclusively contains neutral O-glycans (Li et al., 2011). In the urodele *Triturus alpestris* (*Ichthyosaura alpestris*, Taxonomy ID: 54263) (Florea et al., 2006) Neu5Ac, Neu5Gc, and Kdn derivatives (respectively in the ratio 75, 8, 17) are detected with a high degree of O-acetylation of Neu5Ac and Kdn: Neu5,9Ac2 represents 56% of Neu5Ac derivatives (traces of Neu4,5Ac2 were also observed), Kdn represented 42% of total Kdn derivatives with mono-O-acetylated (39% Kdn9Ac, 6% Kdn7Ac, 2% Kdn5Ac). In the vitelline envelope, among the six ZP glycoproteins also found in other species, ZPA/ZP2/gp69/64, ZPB/ZP4, ZPC/ZP3/gp41, ZPD, ZPX, and ZPY (Hedrick, 2008), only ZPA (Vo et al., 2003) and ZPX are sialylated (Barisone et al., 2003). ZPA displays 30% of the sperm binding activity, while the rest is ascribed to ZPC (Miwa, 2015). It has to be noted that ZPB and ZPC are found in all species, but ZPA is not found in teleost fish, and ZPX is mainly found in invertebrates but is not present in mammals (Feng et al., 2018). In toad *Bufo arenarum* (*Rhinella arenarum*, Taxonomy ID: 38577), Sias composition varies with high levels of Sias present in the preovulation period associated with an 80% decrease in the post-ovulation period, associated with changes in neuraminidases and sialyltransferases activity (de Martinez et al., 1975).

In teleosts, high content of Sias is present around the oocytes, in the *zona radiata* (the equivalent of the mammals ZP) of the dusky grouper *Epinephelus marginatus* (Mandich et al., 2002), the swordfish *Xiphias gladius* (Ortiz-Delgado et al., 2008), the bluefin tuna *Thunnus thynnus* (Sarasquete et al., 2002), and the trout (Kitazume et al., 1994). Kdn is the major component of the vitelline envelope of the chum *Oncorhynchus keta* and kokanee salmon *Oncorhynchus nerka* while cherry salmon *Oncorhynchus masou* contains an analogous family of mucin-type glycoproteins with ending Neu5Gc as the major component (Inoue and Inoue 1997; Gallo and Costantini, 2012). In rainbow trout *Oncorhynchus mykiss*, the vitelline envelope Kdn-glycoprotein is the unique polysialoglycoprotein (PSGP). In *D. rerio*, N- and O-glycans are predominantly bearing a single terminal Neu5Ac or Neu5Gc residue but disialylated structures exist. Monosialylated Neu5Ac and Neu5Gc and Neu5Ac-Neu5Gc or Neu5Gc-Neu5Gc disialyl unit extends from an internal Gal of in internal Lewis(x) (Le^x) structure (Neu5Aca2,3Galβ1,4(Fuca1,3)GlcNAc) (Guerardel et al., 2006).

3.1.2 Unique oocytes species-specific polysialylation contributes to sperm acrosomal reaction

Unlike other monosaccharides, Sias polymers with various degrees of polymerization (DP) are evolutionarily conserved

from sea urchin to humans (Sato and Kitajima, 2013). Sea urchin and fish oocytes display a variety of different Sias polymers in contrast to mammals that mainly contain α2,8-linked N-acetylneuraminic acid residues on a restricted number of polysialylated proteins. In mammals, only a few glycoproteins are polysialylated including: the neural cell adhesion molecule (NCAM) in the central nervous system (Schnaar et al., 2014), synaptic cell adhesion molecule (SynCAM 1), neuropilin-2 (NRP-2), the C-C chemokine receptor type 7 (CCR7), E-selectin ligand-1 (GLG1), the α subunit of the voltage-dependent sodium channel, the CD36 scavenger receptor in human milk, and polysialyltransferases themselves (James and Agnew, 1987; Close and Colley, 1998; Yabe et al., 2003; Galuska et al., 2010; Mühlenhoff et al., 2013; Kiermaier et al., 2016; Werneburg et al., 2016). PolySias are long negatively charged and linear polymers that form a strong repulsive field (Yang et al., 1994).

In sea urchins oocytes, the Sias-rich polysaccharides are composed of Neu5Gc (Valle et al., 2015). In *Hemicentrotus pulcherrimus* oocytes, polySias chains (5-O-glycolyl-Neu5Gca2)_n have a DP (n) ranging from 4 to 40 Neu5Gc residues, and these polyNeu5Gc glycans are O-glycosidically linked to threonine residues (Kitazume et al., 1994; Yeşilyurt et al., 2015). The *S. purpuratus* oocyte jelly coat contains a PSGP with 25 polySias chains O-linked to both threonine (two-thirds) and serine (one-third) residues (Kitazume et al., 1994). These polySias in the oocyte jelly are highly specific to each species of echinoderm and initiate the sperm acrosomal process through calcium entry into the sperm head. The acrosomal reaction that causes granules exocytosis and release of proteolytic enzymes digests part of the thick protective jelly coat to allow access to the oocyte surface. The oocyte jelly also possesses 8-O-sulphated (5-O-glycolyl Neu5Gca2)_n with n ranging from two to 3 at the nonreducing end of the oligo/polySia chain found on the 350 kDa sperm receptor on the oocyte membrane (Kitazume et al., 1996; Sato and Kitajima, 2013), and Neu5Gc8S (Schauer, 2012) (Figure 4).

PolySias structures on glycoproteins were first discovered among vertebrates in rainbow trout eggs and isolated from salmonid as polyNeu5Gc (Inoue and Iwasaki, 1978; Kitajima et al., 1986). These teleost fishes polySias are commonly observed with a remarkable degree of structural diversity in the α2,8-linked polySia chains. Homopolymers of α2,8-linked Neu5Ac or Neu5Gc and heteropolymers containing both Neu5Ac and Neu5Gc residues, or Kdn and O-acetyl substitution at C4, C7, C9, or O-lactyl substitution at C9 were described (Kitajima et al., 1986; Sato et al., 1993). These chains are differentially expressed in various fish species. PolyNeu5Ac and polyNeu5Gc are present in the lake trout *Salvelinus namaycush*, poly (Kdn) in the rainbow trout *O. mykiss*, polyNeu5Aca/Neu5Gc in the brook trout *Salvelinus fontinalis*, polyNeu5Ac/Neu5Gc in the brown trout *Salmo trutta fario* and the Japanese char *Salvelinus leucomaenis pluvius*, polyNeu5Gc in the chum salmon *O. keta*, the cherry salmon *O.*

masou ishikawai, the perciform fish *Sander lucioperca* and the salmonid fish *Coregonus maraena* (Venuto M. T. et al., 2020; Venuto M. T. et al., 2020). Among Osteichthyes superclass, only one polysialyltransferase is detected in perciforms (ST8SIA2), and three are present in salmonids (ST8SIA2-r1, ST8SIA2-r2, and ST8SIA4) (Venuto M. T. et al., 2020; Venuto M. T. et al., 2020)). In zebrafish eggs, diSia (Neu5Ac/Neu5Gc) is identified in N- and O-glycans and is finely regulated during embryonic development (Guérardel et al., 2006; Chang et al., 2009).

3.1.3 PolySias stored in oocytes cortical granules counteract polyspermy and pathogens

During oogenesis in most animal species, mature oocytes accumulate large amounts of glycoconjugates including sialylated glycans in intracellular secretory organelles derived from the Golgi, the so-called cortical granules/vesicles, localized in the peripheral cytoplasm (Gallo and Costantini, 2012).

Sea urchins cortical granules localized in the cortex beneath the vitelline envelope, contain Neu5Ac8S (Yamakawa et al., 2007). These granules are exocytosed at fertilization and contribute to the destruction of the sperm binding capacity of the vitelline layer. Mainly the release of trypsin-like proteases results into the vitelline layer hardening to form a barrier against polyspermy by peroxidases catalyzing tri-tyrosine crosslink.

The teleost unfertilized oocyte cortical granules, named cortical alveoli, are filled with polySias with a high molecular weight (200 kDa) and degrading enzymes. During the cortical reaction after the monospermic fertilization, cortical alveoli exocytosis into the perivitelline space results in polySias cleavage to low-molecular-weight units of around 9 kDa. The cortical alveoli content prevents polyspermy and blocks the micropyle opening in the thick chorion surrounding the oocyte through which the first spermatozoa entered at fertilization. The cortical alveoli glycoproteins are subdivided into two groups, the α 2,8-linked polySia group-containing polyanionic glycan units attached to protein through O-glycosidic linkages and the other containing bulky multiantennary N-linked glycans named hyosoporphin (Taguchi et al., 1994). The PSGP (polysialoglycoprotein) polySias display species-specific structural diversity in fish. In rainbow trout *Salmo irideus* (Inoue and Iwasaki, 1978), salmon *S. leucomaenis pluvius* (Iwasaki and Inoue, 1985), and flounder *Paralichthys olivaceus* (Seko et al., 1989) the PSGP from the hyosoporphin group are the major constituent of cortical alveoli. Rainbow trout *Salmo gairdneri* (Oncorhynchus mykiss gairdneri Taxonomy ID: 857570) oocyte PSGP occur with α 2,8-linked polysialic acid chains linked to O-glycosidic carbohydrate units with a length up to 25 sialic acid residues (Kitajima et al., 1986), and are capped at their non-reducing ends by Kdn (Nadano et al., 1986). In the rainbow trout *S. gairdneri*, PSGP Sias are Neu5Gc, while in *Salmo* and *Salvelinus* species they contain both Neu5Ac and Neu5Gc (Kitajima et al., 1986; Seko et al., 1989). In Indian medaka *Oryzias latipes* and *Fundulus heteroclitus*, a

tetraantennary sialoglycan unit is present (Taguchi et al., 1993; Taguchi et al., 1996). PSGP biosynthesis has been reported at late stages during oogenesis (Asahina et al., 2004; Asahina et al., 2006). The sialylation of PSGP is catalyzed by at least four sialyltransferases present inside the oocyte cortical granules and acting synergistically: an α 2,6-sialyltransferase (ST6GALNAC), and three α 2,8-sialyltransferases (ST8SIA2-r1, ST8SIA2-r2, and ST8SIA4) (Asahina et al., 2004; Asahina et al., 2006; Venuto M. T. et al., 2020). PolySias on PSGP also serve to block polyspermy and protect the oocyte from pathogens in the extracellular environmental medium. The highly modified polySias groups of PSGP, displaying O-acetylation of the hydroxyl-groups on C4, C7, C9, and Kdn at the non-reducing termini, confer resistance to bacterial sialidases. PolySias on PSGP have also been shown to be a Ca^{2+} and ionic storage reservoir and have been supposed to be involved in osmoregulation during early development (Shimoda et al., 1994).

3.2 Zona pellucida, Sias, and sialoglycoproteins in other vertebrates

Oocytes from non-aquatic vertebrates do not bear PolySias (Simon et al., 2013) but display unique Sias structures compared to aquatic species localized in the vitelline envelope and the *zona pellucida*. In addition, humans and several mammalian species lack Neu5Gc (Altman and Gagneux, 2019) due to a single exon deletion/mutation in the gene encoding CMAH (Varki, 2017) (Figure 4). The loss of vertebrate-specific Neu5Gc is puzzling and evolutionary scenarios include the proposal for a strong selection by various parasites, but also a selection *via* females towards paternal Neu5Gc. Humans also have circulating antibodies specifically targeting Neu5Gc which is immunogenic and hypothesized to lead to chronic inflammation and other pathologies (Padler-Karavani et al., 2008). Neu5Gc antibodies do not seem to have a biological function in gametes and reproduction (Altman and Gagneux, 2019), but it cannot be excluded that they could play a role in the female genital tract to avoid interspecies fertilization.

The transparent extracellular ZP surrounding oocytes of non-aquatic vertebrate species is a viscous fibrogranular structure at the border between the oolemma plasma membrane and the inner layer of follicle cells of the cumulus matrix (*corona radiata*) (Sinowatz et al., 2001). ZP matrix proteins have been described in granulosa cells and are supposedly involved in folliculogenesis (ZP3 in rabbit, mice, and human, and both ZP3 and ZP1 in bovine and porcine (Prasad et al., 2000; Hasegawa et al., 2007; Gook et al., 2008). ZP is also involved in the protection against physical damages as it is a thick elastic matrix composed of ZP proteins assembled into polymers, which hardens upon fertilization after a series of changes that also provide a polyspermy block (Papi et al., 2010; Monné and Jovine, 2011; Fahrenkamp et al., 2020; Wassarman

and Litscher, 2021). In some mammals, ZP proteins have been demonstrated to be involved in species-specific recognition and binding (mouse and human ZP2) to the spermatozoa (Avella et al., 2014; Bhakta et al., 2019), and probably play a role in the ZP matrix modification associated to the polyspermy block (mouse and human ZP1 crosslinker of ZP filaments (Greve and Wassarman, 1985; Nishimura et al., 2019). The composition of ZP proteins varies among species. In mammals, ZP is composed of three to eight proteins. In most mammals including humans, ZP is composed of four proteins: ZP1, ZP2/ZPA/gp69/64, ZP3/ZPC/gp41, and ZP4/ZPB, while in mice it is composed of ZP1, ZP2, and ZP3, and in cows, dogs, dolphins, and pigs, ZP2, ZP3, and ZP4. Marsupials from South America have ZP1, ZP2, ZP3-1b, and ZP3-1c or from Australia ZP1, ZP2, ZP3-1a, ZP3-1b, ZP3-1c, ZP4, ZPAX, while monotremes contain ZPY, ZP1 ZP2, ZP3-1a, ZP3-1b, ZP3-2, ZP4, ZPAX (Wassarman and Mortillo, 1991; Lefèvre et al., 2004; Moros-Nicolás et al., 2021). ZP is also composed of acidic glycoproteins with *N*-linked and *O*-linked carbohydrate chains and various sialylated oligosaccharides (Noguchi et al., 1992; Nakano and Yonezawa, 2001; Dell et al., 2003; Pang et al., 2007; Velasquez et al., 2007), (Figure 4). In mammals, sialylated ZP has also been proposed to act in immune recognition as oocytes lack major histocompatibility (MHC) class I molecules that mediate self-recognition, which makes them vulnerable to natural killer cells (Clark, 2010).

In Birds, ZP1, ZP3, and ZPD are the major egg-coat ZP proteins (Nishio et al., 2018). In chicken, ZP2 is concentrated around the germinal disk region (Nishio et al., 2014; Nishio et al., 2018). Mutation of the T168 *O*-glycosylated site reduces the sperm-binding activity of recombinant chicken ZP3 (Han et al., 2010). The highest concentration of Sias is present in the vitelline membrane, but Sias are also found in chalaza, yolk and white (Nakano et al., 1994).

Chimeric engineered mice that express human ZP3 and acquire mouse-type *O*-linked glycans deprived of SLe^x structures are capable to bind murine but not human sperm (Dell et al., 2003; Gahlay et al., 2010). In addition, murine oocytes ZP3 deficient in core 1-derived *O*-glycans (mutated on T-synthase) undergo fertilization (Williams et al., 2007). Mutated mouse oocytes lacking *N*-glycans (mutated for Mgat1, encoding *N*-acetylglucosaminyltransferase I) (Shi et al., 2004), or deficient in core 2-derived *O*-glycans support mouse sperm binding but not human sperm binding compared to wild-type mouse oocytes suggesting a glycan-independent process, and other determinants necessary for taxon-specificity (Hoodbhoy et al., 2005). The respective role of the different oligosaccharide ligands remains to be elucidated as the majority of the *O*-glycans linked to murine ZP3 are sialylated including core two *O*-glycans, LacNAc (Galβ1-4GlcNAc), LacdiNAc (GalNAcβ1-4GlcNAc), Galα1-3Gal, and Neu5Acα2-3(GalNAcβ1-4)Galβ1,4GlcNAc (Sd^a antigen), and SLe^x (Dell et al., 2003; Dall'Olio et al., 2014).

In porcine, the major *O*-linked and *N*-linked oligosaccharides found in the ZP matrix are bi-antennary complex type glycans with antennae containing sialylated polyLacNAc sequences (Dell et al., 1999; Nakano and Yonezawa, 2001). Glycoprotein ZP2 display six, ZP3 three, and ZP4 five potential *N*-glycosylation sites, while ZP4 contains three and ZP3 six potential *O*-glycosylation sites (Yurewicz et al., 1991; Yonezawa et al., 1997). *N*-glycosylation of porcine ZP glycoproteins seems crucial in sperm penetration and binding to the ZP (Lay et al., 2013). While the role of sialylated glycoproteins was not established, *N*-glycosylation of ZP3 might increase sperm binding (Lay et al., 2011). Pig Sias are shown to be synthesized by α2,6-sialyltransferases (Kimura et al., 1999).

In bovine, 77% of the carbohydrate chains of ZP glycoproteins are acidic chains containing Sias (Katsumata, 1996), with mainly Neu5Ac (84.5%) and Neu5Gc (15.5%) linked to Galβ1-4GlcNAc and GalNAc with α2,3- and α2,6-linkages, on *N*- and *O*-glycans respectively (Velasquez et al., 2007). A ZP3/ZP4 association mediates interaction with the spermatozoa and depends on *N*-glycans (Suzuki et al., 2015), with ZP4 having the strongest sperm-binding activity (Yonezawa et al., 2001). The *N*-glycosylation of ZP3 (Suzuki et al., 2015), and the Sias sequence of Neu5Ac (α2-3)Gal (β1-4)GlcNAc (Velasquez et al., 2007) are also implicated in sperm-ZP binding.

In human, mutations in ZP genes encoding ZP1, ZP2, ZP3, and ZP4 are involved in cases of female infertility (Wassarman and Litscher, 2021; Wassarman and Litscher, 2022). Recombinant studies implying glycosylation of human ZP3 revealed this glycoprotein is essential to induce acrosomal exocytosis but not for sperm binding (Chakravarty et al., 2008). Human oocytes ZP *N*- and *O*-glycans display unique SLe^x structures that were initially proposed to act in the sperm interaction with the oocyte but later experiments showed they were dispensable (Miranda et al., 1997; Oehninger et al., 1998; Pang et al., 2011; Wassarman, 2011; Clark, 2013; Gupta, 2018; Bhakta et al., 2019; Gupta, 2021; Tumova et al., 2021). ZP mutated mouse oocytes rescued by human ZP2, and deprived of SLe^x in their zona pellucida matrix were shown to bind to human sperm, and ZP2 sperm-binding domain was determined to be required for human sperm-oocyte recognition and zona pellucida penetration (Baibakov et al., 2012; Avella et al., 2014). Transgenic mouse oocytes ZP composed of mouse ZP1, ZP3, and human ZP4 do not bind mouse sperm, and transgenic mouse ZP containing human ZP1, ZP3, and ZP4 do not bind human sperm and are infertile (Avella et al., 2014; Bhakta et al., 2019), supporting ZP2 as necessary and sufficient for human and mouse sperm binding. Genetic removal of ZP2 *N*-terminus glycan does not impede sperm binding or female fertility (Tokuhira and Dean, 2018), showing that ZP2 *N*-terminus is dispensable for gamete recognition (Avella et al., 2014; Tokuhira and Dean, 2018). Chimeric mouse ZP2 proteins, with a human *N*-terminal domain, support human sperm binding evidencing that the *N*-terminus of ZP2 being involved in human species-




FIGURE 5

Dynamical changes in sialoglycoproteins during the sperm journey to reach the fertilized oocyte. Spermatozoa maturation first occurs in the epididymis and the seminal vesicle in the male genital tract with the addition of sialic acids to glycoconjugates (including sialoglycoproteins) in the outer coat (khaki dots surmounted by red diamonds) (1, numbers in circles). Sialoglycoproteins from the head and tail parts help to maintain spermatozoa uncapacitated state (2) (upper blue panel). In the female genital tract (upper pink panel) multiple steps allow sperm to gain mobility (3), escape immune response (4), form an oviductal reservoir (5), acquire fertility through capacitation (6) associated with high swimming potential (7), bind to the oocyte (8), and finally trigger the acrosomal reaction to remove the oocyte extracellular sialic acid coat (9) (upper yellow panel). Numerous Siglecs localized on epithelial oviductal cells and immune cells in the female genital tract are involved in spermatozoa protection/survival (4). In species with external fertilization the sperm coat is formed within the male genital tract (numbers in squares, 1, lower blue panel), a balance exist between species specificity and protection (2), and the specific oocyte sialic acid coat removal also requires an extracellular spermatozoa acrosome reaction (3, yellow panel) before fertilization (4). Several species-specific sialic acids acting at different levels of the sperm journey in both internal and external fertilization are highlighted in red.

specificity binding of sperm. (Avella et al., 2014). Moreover, the human sialyltransferase ST3GAL4 gene is present in ovaries but its precise role is not understood (Tanagushi et al., 2003). ZP Sias could mediate other functions related to immune recognition (Clark, 2010). Indeed, SLe^x structures are ligands for Siglec nine tyrosine-based immunoreceptors that can transmit signals to a variety of different types of leukocytes and lymphocytes to allow oocyte survival and block the female immune response (Angata and Varki 2002; Avril et al., 2004).

A clue to understanding this diversity could come from the ZP glycoproteins synthesis study that differs between several vertebrate species. Mammalian ZP glycoproteins are transcribed in human, monkey, rabbit, dog, and cow both by

the oocyte and the follicular cells, while in mice the oocyte exclusively contributes to their synthesis (Sinowitz et al., 2001). In birds, ZP1 is from the liver and both ZP3 and ZPD are from follicular granulosa cells (Nishio et al., 2018).

4 Sperm preparation through the coating/uncoating changes in Sias

The spermatozoa of several invertebrates such as sea urchins, and non-mammalian vertebrates including most fishes and amphibians possess the capacity to fertilize oocytes when they

leave the testis after they have been released for external fertilization. On the contrary, in mammals, testicular spermatozoa do not have fertilizing capacity but need to mature by passing through the epididymis where sialic acid is deposited to cap most sperm glycan chains of the membrane cell surface. From the high numbers of spermatozoa deposited in the female reproductive tract, only a few thousand enter the oviduct, and fewer reach the ampulla region of the oviduct at the appropriate time window for fertilization (Varki, 2017). If the Sias coat regulates sperm migration through the cervical mucus, and the formation of the sperm oviductal reservoir, spermatozoa remain unable to fertilize an oocyte until sialoglycoproteins partial removal exposes specific critical structures during capacitation (Varki, 2017), (Figure 5).

4.1 Sias addition to sperm is a prerequisite for their maturation

4.1.1 Sperm from aquatic species

The sea urchin sperm surface of *Lytechinus variegatus* is coated by highly sulfated polysaccharide and composed almost exclusively of Neu5Ac (Valle et al., 2015) that potentiate sperm acrosome reaction (Hirohashi and Vacquier, 2002). *H. pulcherrimus* and *Strongylocentrotus purpuratus* also present a majority of Neu5Gc residues (Kitazume et al., 1994; Kitazume et al., 1996; Miyata et al., 2004). The sperm flagella displays two distinct polySia structures, which are α 2,8- and α 2,9-linked polymers of Neu5Ac: an 8-O-sulfated (9Neu5Ac α 2)_n structure of 40–80 kDa (with a DP of 15) carried by flagelliasialin, and an α 2,8-linked polyNeu5Ac O-glycan of 190 kDa (Miyata et al., 2004; Miyata et al., 2006; Miyata et al., 2011). Flagelliasialin is involved in voltage-sensitive sodium and calcium channels that control sperm motility (Kambara et al., 2011). The flagellum and the head express a 190 kDa glycoprotein that facilitates proteins rearrangement on the spermatozoa surface upon activation (Miyata et al., 2011). As a dynamic mosaic bilayer, the plasma membrane is enriched in microdomains (lipid rafts) that serve as platforms for protein segregation (Simons and Ikonen, 1997; Hakomori, 2004; Kim et al., 2020). Spermatozoa Sias are enriched in these membrane microdomains (Chen and Varki, 2010) where flagelliasialin also tends to concentrate (Sapoń et al., 2019). Clustered patches formed in dynamic microdomains could relocate sialoglycoproteins on the sperm plasma membrane and play an active role during capacitation (Wang et al., 2021).

In *Xenopus* spermatocytes Sias are histochemically detected in O-linked oligosaccharides and N-glycans (Valbuena et al., 2012).

In teleost such as in Siluriform catfish, *Heteropneustes fossilis*, a seminal vesicle present in the male reproductive tract (Legendre et al., 1996) ensures spermatozoa storage and secretion of Sias and could play a role in sperm survival and maintenance (Senthilkumaran and Joy, 1993). In the teleost catfish *Clarias*

batrachus, Sias associated with sperm in the seminal vesicle are submitted to annual variations with high levels in the spawning phase (summer peak) and low values in the resting phase (Singh and Joy, 1999). In Acipenseriformes sturgeons, *Acipenser baerii*, and *Acipenser schrenckii*, Neu1 is a biomarker associated with sperm quality (Li et al., 2017). The concomitant expression of the teleost-specific α 2,3-sialyltransferase gene *st3gal3-r* and the two sialidase genes *neu1* and *neu3.5* in zebrafish testis has not been explained (Yamakawa et al., 2018).

4.1.2 Sperm sialome maturation and immune protection in the female genital tract of several amniotes

The mammalian sperm sialome is established during spermatogenesis in the testis. It requires afterwards an epididymal maturation and the incorporation of sialylated seminal fluid components around the sperm membrane during ejaculation. These incorporations increase the number and the diversity of the sperm Sias involved in sperm progression through the female genital tract. To limit microbial attack, the female reproductive tract is submitted to immune surveillance to promote the destruction of pathogens (Wira et al., 2011) and an uterus endometrium antimicrobial resistance (Nasu and Narahara, 2010). Sperm entry into the uterine lumen triggers an immune response (or leukocytic reaction) that destroys most of the spermatozoa (Thompson et al., 1992). To resist this massive destruction, spermatozoa (as oocytes) lack major histocompatibility (MHC) class I molecules involved in the immune system self-recognition which renders them less susceptible to class I restricted cytotoxic effect by T lymphocytes, but are vulnerable to natural killer cells (Dell et al., 2003) that spermatozoa have to bypass.

The first maturation process in the Sias coat of spermatozoa occurs in the epididymis (Holt, 1980; Hoffmann and Killian, 1981; Toowicharanont and Chulavatnatol, 1983; Ma et al., 2016). Chicken spermatozoa are extensively sialylated and contain residues of α -mannose, α -glucose, α - and β -galactose, α -fucose, α - and β -N-acetylgalactosamine, N-acetyllactosamine, and monomers and dimers of N-acetylglucosamine restricted to the acrosomal region (Peláez et al., 2011).

In mice, sperm bears α 2,3-linked Neu5Ac, with more than 20% of Sias transferred to mature sperm in the caput and corpus epididymis by sialyltransferases (Ma et al., 2016). ST8SIA3, ST6GALNAC2, and ST6GAL2 are detected in mouse testis and could probably account for sialylations on sperm (Yoshida et al., 1995; Kurosawa et al., 1996; Takashima et al., 2003) or ST6GAL transcripts in rat (Bessler et al., 1995). Interestingly, in ST8SIA2 and ST8SIA4 KO mice, the loss of polySias alters the contractile phenotype of peritubular smooth muscle in the postnatal testis (Hachem et al., 2021). A 54 kDa sialoglycoprotein binds to the immature mouse sperm during maturation in the epididymis and is implicated in the protection of sperm from phagocytosis by macrophages (Toshimori et al., 1991). Sperm phagocytosis and

antigenicity are also proposed to be partly avoided in chicken, mouse, and human by a similar mechanism based on Sias (Steele and Wishart, 1996; Yudin et al., 2005; Ma et al., 2016).

In primates, β -defensin 126 (DEFB126) formerly known as epididymal secretory protein 13.2 (ESP13.2) coats sperm surface until completion of capacitation (Yudin et al., 2005), and gp20/CD52 was also reported to coat the whole sperm with an elevated presence over the equatorial segment of the sperm head (Della Giovampaola et al., 2001). DEFB126 adsorbed on the sperm surface is necessary for progression through the cervical mucus until the upper oviduct, and the site of fertilization (Tollner et al., 2008; Szczykutowicz et al., 2019). DEFB126 and gp20/CD52 Sias moieties are responsible for cloaking and probably protect the sperm surface from immune recognition in the female reproductive tract (Zhou et al., 2004). Sialylation on spermatozoa (Teclé and Gagneux, 2015) provides negatively charged and repulsion surfaces between sperm and oviductal cells necessary for their transit through the cervix (Tollner et al., 2011).

In human and rooster, the reduction of sperm Sia content is associated with lower fertility due to a migration default in the reproductive tract (Froman and Thursam, 1994; Tollner et al., 2011). Human seminal plasma and sperm acrosome contain other components that are supposed to counteract the female immune system such as *N*-glycans with SLe^x ending structure (Pang et al., 2007). Notably, polymers with a DP > 46 Sias residues directly and significantly reduce H1, H2A, H2B, H3, and H4 histones cytotoxicity (Zlatina et al., 2017). A glycomic gradient in the male genital tract fluids (from the cauda to the caput region of the epididymis), varying from high mannose to sialylated complex *N*-glycans is associated with abundant Le^x multimeric structures, and might also play an important role in the immune protection (Cheon and Kim, 2015). The human spermatozoa are suspended in the secreted seminal plasma composed of α 2,6-sialylated branched *N*-glycans and Sias α 2,3-linked *O*-glycans (Kratz et al., 2015). Spermatozoa are covered by sialylated fibronectin, detected to lower levels in infertile men (Kratz et al., 2011). The human epithelial cells of the epididymis caput are polySia-positive and polysialylated ST8SIA2 and polysialylated NCAM are integrated into the sperm membrane of the post-acrosomal region during the epididymal transit (Simon et al., 2013). Several ST3GAL4 (Taniguchi et al., 2003) or ST3GAL2 transcripts (Takashima et al., 2002; Krzewinski-Recchi et al., 2003) are detected in human testis, and postulated to be important for sperm Sias-coat acquisition. Sperm polySias may be cytoprotective from the innate and adaptive immune system of females. PolySias could bind Siglec 11 expressed on tissue macrophages, suppress the immune response (Wang and Neumann, 2010), and decrease the extracellular histones and nucleosomes cytotoxicity after their release from the neutrophils (Neutrophils extracellular traps) that have been detected after insemination (Alghamdi and Foster, 2005; Brinkmann and Zychlinsky, 2007). Intriguingly the X chromosomal type

sperm exhibits a higher negative charge compared to the Y due to a difference in exposed Sias content (Kaneko et al., 1984).

Negatively charged Sias deliver signals through a binding to Siglecs (Li and Ding, 2019). Several Siglecs, present on the sperm surface displaying species- and regional-specific expression, are thought to be involved in sperm immune protection (Fernandez-Fuertes et al., 2018), such as Siglecs 1, 2, 5, 6, 10, and 14 expressed at the surface of, bull, ram, and human spermatozoa (Alkhodair et al., 2018), and Siglecs 11, and 16 present in human (Teclé et al., 2019). Bovine Siglecs 2, 5, 6, and 10 are expressed in the sperm neck region, and Siglecs 6 and 10, in the anterior head region (Alkhodair et al., 2018). Sperm has been found to strongly bind Siglecs present on immune cells which may facilitate sperm survival in the face of female immunity. In mice, the interaction of sialylated sperm to Siglec six predominantly present in B cells is positively correlated with protection from phagocytosis (Ma et al., 2012; Ma et al., 2016), and Siglec 3 in the endometria facilitates sperm survival (Teclé et al., 2019). In human, highly sialylated sperm binds Siglec 10 in the endometrium (Teclé et al., 2019), and a sperm Sia-containing glycoprotein, CD24 (a known activator of Siglec 10) (Chen et al., 2009)), interacts with endometrial expressed Siglec 10 and with neutrophil Siglec nine during the leukocytic reaction and increases sperm survival (Teclé et al., 2019). Glycodelin-A (GdA) secreted from the endometrial glands and the decidual glandular epithelium has two *N*-linked glycosylation sites and bears terminal α 2,6-sialylated LacNAc and LacdiNAc antennae with immunosuppressive actions on cellular and humoral immune responses by blocking Siglec 2 (CD22) binding and activated-events mediated by T and B cells (Seppälä et al., 2007), (Figure 5). In addition to the Sias and Siglecs interaction between sperm and female genital tract, other mechanisms involving Sias may be involved in maternal immune regulation (Clark, 2013). In human, the high density of *N*-glycans terminated with a SLe^x are the ligands for Siglec nine and for selectins families of ligands, which is consistent with the hypothesis that human oocyte ZP glycans function in immune recognition (Clark, 2013). In addition, the female innate recognition is probably facilitated by the prevention of the complement deposition of factor H in the female reproductive tract (Brandtzaeg, 1997), as the binding of factor H to cell surface Sias could favor a conformational change that inhibits the alternative complement pathway activation and prevent the formation of the membrane attack and death complex.

4.2 Sperm Sias coat removal and mobility acquisition

4.2.1 Sperm Sias and the storage in a female reservoir

In many animal species, sperm is stored by mated females before fertilization through several mechanisms. Some species

have specialized storage organs or reservoirs, while others collect sperm in particular areas. Sperm storage within the female reproductive tract (up to months and years) is common in fish, bird, and amphibian females in oviductal sperm storage tubules or cloacal spermathecae (Sever et al., 2001). Tailed frog *Ascaphus truei* has oviductal sperm storage tubules while salamanders possess cloacal spermathecae (Sever et al., 2001). However, little is known about Sias function in this storage process in aquatic species. In birds, sperm transport and storage into tubules relies on Sias (Steele and Wishart 1996). In mammals, the oviduct is considered to be the sperm reservoir (Neubaum and Wolfner, 1999).

In mammals, sperm move to specialized regions that prolong viability and prepare for fertilization. Mammals oviductal sperm reservoir is an essential step to coordinate the sperm capacitation process and detachment from the oviduct in coordination with the appearance of the ovulated oocyte (Scott, 2000; Suarez, 2016). A few thousands from the millions of spermatozoa deposited in the female reproductive tract reach the oviduct where they bind in a Sia-dependent manner in the isthmus part to the ciliated ductal epithelial cells. The binding of spermatozoa involves Sias in hamster (Demott et al., 1995), rat (Cortés et al., 2004), and porcine where the sperm binds to specific oviductal α 2,6-sialylated biantennary *N*-glycans (Kadirvel et al., 2012). In horse and bovine sperm-oviduct binding does not involve Sias but galactose (Sabour and Ball, 2007), and fucose (Lefebvre et al., 1997) respectively. In human, the high sialylation of DEFB126 that coats the entire sperm surface is necessary for oviductal binding, extended viability, and capacitation retardation (Tollner et al., 2012). Sperm bound in a sialic acid-dependent manner to the isthmus of the oviductal epithelium is only detached to swim toward the oocyte in the ampulla of the oviduct when capacitation is induced (Suarez, 2016).

4.2.2 Sperm capacitation unmasks Sias and triggers hyperactivation

Capacitation in aquatic species results from different types of interactions with either the surrounding external medium and the oocyte jelly, or can occur within the male genital tract. Sea urchin spermatozoa capacitation occurs in seawater due to a lower potassium concentration and a basic pH that enables the ATP hydrolysis for sperm motility (Tosti, 1994). In amphibians, the fibrous glycoconjugate network of the oocyte jelly layer containing Sias is involved in capacitation (Brun, 1974). In Fish, sperm capacitation is diverse and mediated either by seminal plasma changes in pH, HCO_3^- , Na^+ , Ca^{2+} , by release in the external medium, or after a contact with the oocyte surrounding Sias (Pérez, 2020).

In mammals, capacitation is also an important maturation process for spermatozoa (Aitken and Nixon, 2013). As ovulation occurs, spermatozoa undergoing capacitation shed their surface glycoproteins, reducing their binding to the oviduct epithelium,

and start to develop hyperactivated motility. This increased mobility aids them to detach from the oviduct to further access the superior female reproductive tract (Töpfer-Petersen et al., 2002). In mouse, sperm releases sialidases that leads to the loss of 20% of the sperm sialome (Ma et al., 2012). In bull, the total removal of sialic acid from sperm decreases the motility and the mucus penetration ability but increases ZP binding and unwanted polyspermic penetration showing that the desialylation event requires fine-tuning (Fernandez-Fuertes et al., 2018).

In human, desialylation is restricted to a limited number of proteins, performed by sialidases present on the sperm surface (Ma et al., 2012), and likely facilitated by sialidases present in the female reproductive tract (Velasquez et al., 2007). Sperm neuraminidase 1 and 3 (Neu1 and Neu3) removing α 2,3-, α 2,6-, and α 2,8-linked Sias unmask certain sperm surface antigens required for the ZP recognition and binding (Lassalle and Testart, 1994; Ma et al., 2012). These sperm membrane-associated sialidases are activated by a specific Sia-binding protein found in the uterus (Banerjee and Chowdhury, 1997). Six proteins are desialylated (endothelial lipase, serine proteases 39 and 52, testis-expressed protein 101, and zonadhesin), while one becomes sialylated (aconitase), (Villaverde et al., 2020). The Sias of the *N*-glycan found on Asparagine 162 on aconitase is involved in a switch to glycolysis in capacitated sperm (Fraser and Lane, 1987). The remaining sialoproteins are relocated in the post-acrosomal region (Focarelli et al., 1995). In mouse as in human, capacitated spermatozoa gain increased motility called hyperactivation to leave the oviductal storage reservoir (Okabe, 2013). The CatSper1 component of the CatSper channel complex is an *O*-linked glycosylated protein that contains terminal Sias residues. The channel complex is compartmentalized within the flagellar membrane where it creates linear Ca^{2+} signaling nanodomains along the sperm tail to increase the mobility (Qi et al., 2007; Okabe 2013; Ded et al., 2020). Capacitation does not occur at the same time for all spermatozoa (Buffone et al., 2012). The oviduct regulates capacitation to avoid over-capacitation and premature acrosome reaction and ensure a continuous supply of fertile sperm (Boilard et al., 2002; Fazeli et al., 2003). It has been supposed that CatSper1 Sia-protein serves to select spermatozoa most likely to undergo capacitation (Ded et al., 2020). In human, a 25 kDa protein located on the non-capacitated sperm head plasma membrane binds to a Ca^{2+} -dependent Sia-binding protein (SABP) present on the endometrium and contribute to increase sperm motility (Banerjee and Chowdhury, 1997), (Figure 5).

5 Fertilization and Sias

At fertilization, sperm binding results in acrosomal exocytosis, and sperm-ZP penetration, before gamete fusion (Mengerink and Vacquier, 2001). In several species, part of

the Sias covering gametes as “biological masks” are removed to facilitate specific recognition of underlying glycans by specific binding proteins.

5.1 Sperm binding to the oocyte and sialoglycosylation

In spawning freshwater and marine organisms, the dilution that reduces the probability of gamete encounters is counteracted by the release of diffusible chemoattractant by the oocyte jelly (Bader, et al., 1998; Olson et al., 2001; Tholl et al., 2011; Guerrero et al., 2020). Once into close contact, the spermatozoa interact with the oocyte.

Sea urchin sperm (Miyata et al., 2004) interacts specifically with the oocyte jelly coat containing polySias and sulfated oligoSias chains inducing the acrosomal reaction (Yeşilyurt et al., 2015). These oligo/polySias act not only in sperm activation but are also involved in a species-dependent recognition by the release of a sperm major species-specific recognition effector called bindin after the contact with the oocyte sperm receptor and the subsequent acrosomal reaction (Lopo et al., 1982; Kitazume et al., 1996; Sato and Kitajima, 2013). The oocyte bindin receptors are aggregated into complexes in lipid rafts and the sperm polySias receptors mediate specific interactions (Hakomori, 2004; Cohen and Varki, 2010). However, the spermatozoa receptor for polyNeu5Gc is not identified and questions remain regarding carbohydrate-based sperm-oocyte recognition. In addition, polySias modify the spermatozoa intracellular pH and have been proposed to be involved in the extension of the acrosomal protrusion formed by actin filaments (Summers and Hylander 1975; Hirohashi et al., 2008).

In amphibians, except for some Urodeles where the sperm is released in the cloaca and stored in a spermatheca (Watanabe and Onitake 2002), fertilization occurs in the external medium. In newt Urodele, spermatozoa are directly inseminated on the oocyte jelly (rich in sialic acid and cations) before motility is triggered. In anuran, sperm motility is instead triggered by the external environmental composition changes and followed by a sperm-oocyte binding involving ZP, Sias and other glycoproteins.

Fish also show diverse reproductive strategies with most oviparous species having external fertilization, and few other zygoparous displaying internal fertilization. In several fishes having external fertilization, glycans can guide the spermatozoa to the oocyte into the micropylar canal playing a role similar to cell surface receptors. In the rainbow trout, *S. gairdneri*, oocytes express a 500 kDa Kdn-glycoprotein located in their vitelline envelope around the micropyle through which the spermatozoa can penetrate (Kanamori et al., 1990; Tezuka et al., 1994). This Kdn glycoprotein contains complex N-glycans as minor components and is heavily O-glycosylated with α 2,8-

linked Sias. In the cherry salmon vitelline envelope, the major component is a Neu5Gc mucin-type glycoprotein (Yanagimachi, 2011; Yanagimachi et al., 2017). Other teleost fishes present different sperm entry mechanisms through their micropyle being simply physical by an indented chorion around the micropyle in zebrafish and loach or requiring grooves in goldfish. In the case of flounder, herring, and Alaska pollock, the micropyle is a simple funnel devoid of sialoglycoprotein (Yanagimachi et al., 2017). The spermatozoa attached to the oocyte through the micropyle triggers the cortical alveoli discharge into the perivitelline space leading to PSGP cleavage, and an increase in the osmotic pressure which causes an influx of external water that swells the perivitelline space. This further ensures the chorion hardening and the closure of the micropyle to avoid polyspermy (Mengerink and Vacquier 2001; Gallo and Constantini, 2012). In the teleost Sebastidae blackbelly rosefish *Helicolenus dactylopterus*, the male possesses a urogenital copulating organ and fertilization occurs in the female genital tract where mucous cells from the papilla secrete sialoglycoproteins, O-linked mucin-type glycans terminated with α 2,3-linked Sias, and mannose type N-linked glycans terminated with sialic acid α 2,6-linked Sias to galactose/N-acetylgalactosamine, which role remains to be clarified (Accogli et al., 2018).

In mammals, fertilization takes place in the ampulla of the fallopian tubes, and sperm binding to the oocyte occurs in the ZP made of N- and O-glycoproteins (Teclé and Gagneux, 2015). The molecular model proposed to explain sperm-oocyte interaction in mammals has changed through the years, going from ZP2 and ZP3 acting as sperm receptors (Bleil and Wassarman, 1980; Bleil and Wassarman, 1988; Bleil and Wassarman, 1990), to the possibility that sperm interaction with the ZP depends on the supramolecular structure of the oocyte coat rather than a single ZP subunit (Rankin et al., 2003), to a more recent view with one domain of ZP2 playing a central role in gamete recognition by physically interacting with counterpart molecule(s) on sperm (Avella et al., 2014). Currently available data do not fully exclude the role of ZP glycosylation and Sias in fertilization in one of its multistep processes such as the formation of the membrane block to polyspermy, or ZP hardening, but this involvement differs from what was initially suggested and varies between species.

In pig, gamete recognition depends more on N- than O-glycans (Noguchi et al., 1992; Yonezawa et al., 1997), and particularly the main ZP-neutral di-antennary N-glycans (Nakano et al., 1996; Topfer-Petersen, 1999). The N-glycosylation of ZP glycoproteins was shown necessary for sperm-ZP interaction, including sperm binding to ZP as discussed in chapter 3.2 (Lay et al., 2013).

In bovine, Sias of the ZP are physiologically implicated in the binding of the sperm to the ZP (Velasquez et al., 2007).

Mice genetically modified, lacking N- and O-glycans (and without Sias), are fertile as discussed in chapter 3.2, showing sperm binding is independent of N- and O-glycans from the ZP

(Thall et al., 1995; Shi et al., 2004; Williams et al., 2007; Raj et al., 2017; Tokuhira and Dean, 2018). N-glycosylation was shown important for the normal secretion of ZP2 (Roller and Wassarman, 1983).

For human, terminal SLe^x sequences, present on human ZP, are not essential for sperm binding as discussed in chapter 3.2 (Miranda et al., 1997; Oehninger et al., 1998; Pang et al., 2011; Bhakta et al., 2019; Tumova et al., 2021), and requires other molecular effectors (Siu et al., 2021), even if lectin-like and protein–protein interactions were initially shown to play a role in gamete interaction (Clark, 2013). Human ZP2 without SLe^x antigen, expressed in transgenic mice, is sufficient for human sperm binding and penetration in the ZP (Baibakov et al., 2012; Avella et al., 2014). Chimeric ZP2 proteins composed of the human N-terminus instead of the mouse terminal region allow human sperm binding showing a role for this region in species-specificity (Avella et al., 2014; Tokuhira and Dean, 2018).

Interestingly, in *Drosophila*, oocytes are surrounded by a thick protective chorionic eggshell that presents a cone-shaped specialized projection, or micropyle, through which the sperm enters at fertilization as in teleosts (Horne-Badovinac, 2020). While *Drosophila* polySias homopolymers are present during embryonic development, no Sias nor polySias glycoproteins were detected before fertilization in oocytes and spermatozoa (Koles et al., 2004). It is hypothesized that Sias could be unnecessary at fertilization or act in a pre-mating step with a preferential selection of the spermatozoa through transport and storage in the specialized female spermatheca. This selection would build an efficient reproductive barrier by preventing heterospecific fertilization by a sperm selection before gamete encounter (Intra et al., 2009).

5.2 Sperm acrosomal reaction, oocyte deprotection, and membrane block to polyspermy

In sperm, the outer acrosomal membrane at the apical tip of the head fuse with the plasma membrane allowing exocytosis of the acrosomal contents originating from the spermatid's Golgi complex and containing a variety of lytic enzymes and ZP-binding proteins including acrosin, hyaluronidase, and sialidases (Srivastava et al., 1988; Yanagimachi, 2011; Khosravi et al., 2016). Acrosome-reacted sperm digest the protective oocyte vestment and bind to the oocyte plasma membrane (Okabe, 2015). Upon contact with the spermatozoa, several vertebrates and invertebrates oocytes exhibit polyspermy-preventing reactions. The membrane block to polyspermy is a rapid reaction that converts the oocyte jelly coat to a form that does not support sperm interaction. Urodeles amphibians, cartilaginous fish, and birds accept polyspermy. The total content of Sias between the vitelline membrane compared to

the fertilization membrane does not vary (Wolf et al., 1976). It is not known if this absence of level changes is restricted to some species and if it reflects the sum of sialylation/desialylation processes or is simply a low level of Sias dynamical change.

In sea urchin, as seen in chapter 3.1.2, the acrosomal reaction is potentiated by fucose-sulfated polymers and also requires the oocyte Sia-rich polysaccharides found on the 350 kDa sperm receptor that belongs to the heat shock protein 110 family (Kitazume et al., 1996; Hirohashi et al., 2008; Sato and Kitajima, 2013; Yeşilyurt et al., 2015). The oocyte 350 kDa sperm-binding protein is largely localized in the vitelline layer and lipid rafts and acts for a large part in the acrosome reaction (Maehashi et al., 2003). In sea urchin and non-mammalian species, the membrane block to polyspermy is mediated by the exocytosis of oocyte cortical granules to convert and elevates the oocyte jelly coat. It is accompanied by multiple biochemical changes and by a transient depolarization of the oocyte plasma membrane potential (Jaffe and Cross 1986; Longo et al., 1986; McCulloh and Chambers, 1992).

In mammals, sperm capacitation destabilizes the acrosomal sperm head membrane for penetration in the outer layer of the oocyte in addition to the chemical changes in the tail necessary for greater mobility. Sialylated acrosin released from the sperm acrosome digests the ZP allowing part of the sperm plasma membrane to fuse with the egg plasma membrane, in the boar (Schleuning et al., 1975), or hamsters (Hirose et al., 2020). Acrosin homologs have been identified in quail bird (Sasanami et al., 2011), and *Xenopus* (Kubo et al., 2010) to bind to the oocyte. However, in mice, acrosin null mutation did not impair fertilization indicating it is not a necessary molecular effector in this process (Wassarman, 2011). The polyspermy-preventing reaction in mammals is composed of complex slow calcium-dependent reactions in the oocyte jelly and plasma membrane, requiring cortical granule exocytosis and ZP conversion (Evans, 2020). The cortical granules exocytosis allows ZP mechanical properties to change from elastic to plastic, also called hardening (Papi et al., 2010; Gupta and Bhandari, 2011; Fahrenkamp et al., 2020). However, to prevent polyspermy mammals' oocytes have been observed to preferentially use a ZP coat block or a plasma membrane block. In sheep or dog, no sperm is found in the perivitelline space, between the ZP and the plasma membrane, suggesting an effective ZP block while in some other species such as rabbit, or mole sperm is found in the perivitelline space suggesting a plasma membrane block rather than a ZP block (Evans, 2020).

In pig, the N-glycosylation of ZP glycoproteins was shown necessary for the induction of acrosomal exocytosis in the ZP-bound sperm (Lay et al., 2013). The number of oocyte Sias decreases after fertilization (Rath et al., 2005), and after the exocytosis of cortical granules (Sun, 2003). Desialylation of the oocyte decreases sperm acrosome reaction and polyspermy block (Lay et al., 2011).

In bovine, it was hypothesized that neuraminidase released from the cortical granules would participate in the polyspermy block by removing Sias from the ZP (Velasquez et al., 2007). Sias need to be removed from bovine oocyte coat to allow sperm interaction but are required to avoid abnormal fertilization (Fernandez-Fuertes et al., 2018), pointing to a fine-tuned regulation of Sias.

In mice, two sperm sialidases (neuraminidases 1 and 3), shed during capacitation, interfere with ZP binding but are not associated with the polyspermy block (Ma et al., 2012). By using a transgenic mouse line expressing GFP in the acrosome, it was determined that most sperm undergo an acrosome reaction before contacting the ZP *in vitro* (Hirohashi et al., 2011; Jin et al., 2011). The timing of the acrosome reaction before the ZP binding is flexible, and exocytosed enzymes are dispensable for sperm penetration in the ZP (Okabe, 2015). The majority of sperm has undergone acrosome exocytosis in the upper segments of the oviductal isthmus before the ampulla (La Spina et al., 2016) and the cumulus (Hino et al., 2016; Muro et al., 2016), suggesting that sperm acrosome exocytosis could take place in the female oviduct during sperm migration, and questioning about the role played by Sias in the female reproductive tract.

In human, terminal SLe^x sequences have been involved in the sperm acrosomal reaction (Miranda et al., 1997; Oehninger et al., 1998; Pang et al., 2011; Tumova et al., 2021). Recently a protein bearing terminal SLe^x, C1orf56, localized in the human acrosomal region of the spermatozoa was proposed to act in the ZP-induced acrosome reaction (Wang et al., 2021). In addition, three of the four reported glycodefin (Gd) sialylated glycoproteins (acquired by the sperm in the female genital tract) GdA, GdF, and GdC, but not GdS coating the spermatozoa, are involved in the regulation of the acrosomal reaction. GdF blocks premature acrosomal reaction during Fallopian tube migration. GdF is subsequently converted into GdC by the oocyte cumulus cells when a sperm-oocyte contact is established, removing GdF inhibition (Sawyer, 2021).

6 Conclusion

In reproduction, there is a need to allow fertilization between different individuals in a given species, with the prevention of unwanted cross-fertilization by other related species, and a limited microbial attack. Cell-surface sialoglycoproteins coating germ cells show sufficient diversity to fulfill these recognition/restriction aspects even if they are not the sole type of post-translational modifications involved in this process. Sialylated glycoproteins display a large repertoire and have a spatial organization to allow several types of complex interactions with clustering in species with internal fertilization and polySias chains in species with external fertilization. Sialoglycoproteins are involved in many steps before fertilization including sperm maturation and are part of the molecular basis of gamete binding in several species. In mammals and fish with internal fertilization, sialylated glycoproteins

have an emerging role in the passage and attachment of sperm through the female reproductive tract where they fulfill a protective action against female immunity that starts to be understood. The identification of specific molecules that mediate sperm storage would help clarify how sperm can be kept viable. In free spawning organisms submitted to a dilution in the external media, polySias chains play important roles in terms of pathogens/pollutants and as self/non-self-recognition mechanisms.

However, many important issues related to Sias terminal residues in the surrounding gametes matrix remain unresolved. For example, what is the role of ZP3 Sias in mammals other than mice, and do they act in immune recognition? Why are polySias present in aquatic species and what is their relevance compared to other shorter forms of terminal sialic acids in terrestrial species? Are Sias mainly involved in immune recognition, or do they act in membrane fusion by favoring membrane attachment to different proteins located in multimeric complexes? Defects in the formation of microdomains at the level of the human oolemma plasma membrane have been demonstrated to be related to fertilization failure (Van Blerkom and Caltrider, 2013).

To generate a sufficient level of complexity, sialylated glycoproteins present at the cell surface are synthesized in the Golgi apparatus in a process subjected to multiple sequential and competitive enzymatic pathways (Drickamer and Taylor, 1998; Esko and Selleck, 2002). The expression of sialylated glycoproteins is regulated by a limited set of conserved sialyltransferases in terms of orthologs and enzymatic activity (Harduin-Lepers et al., 2005) providing the basis for a rationale for their expression (Varki, 2006; Yamakawa et al., 2018).

More than ever, the determination of gene expression patterns and activity profiles of the enzymes involved in sialoglycoproteins biosynthesis is a requisite to better understand their biological functions and interconnections in gametes and at fertilization. In this respect, further studies on sialyltransferases in several species using newly developed technology, such as CRISPR-Cas9, linked to an analysis of the gonads could provide future valuable information.

Author contributions

Conceptualization: IF, AH-L, and KC; writing: IF, GM, CM, MD, AM, MM, J-FB, AH-L, and KC.

Funding

CM is a recipient of a doctoral fellowship from the French ministry. MD is a recipient of a doctoral fellowship from the French Ministry and Region Hauts-de-France. The contribution of the COST Action CA18103-INNOGLY supported by the European Cooperation in Science and Technology (COST) is greatly

acknowledged. This work was supported by the CNRS, the ANR-21-CE44-0032 (project PsaMar) and the University of Lille.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Accogli, G., Sion, L., Maiorano, P., Capezzuto, F., D'Onghia, G., and Desantis, S. (2018). Regional morphology and mucus composition in the urogenital papilla skin of the blackbelly rosefish *Helicolenus dactylopterus* (Delaroche, 1809). *Anim. Reprod. Sci.* 189, 119–127. doi:10.1016/j.anireprosci.2017.12.018
- Aitken, R. J., and Nixon, B. (2013). Sperm capacitation: A distant landscape glimpsed but unexplored. *Mol. Hum. Reprod.* 19, 785–793. doi:10.1093/molehr/gat067
- Alghamdi, A. S., and Foster, D. N. (2005). Seminal DNase frees spermatozoa entangled in neutrophil extracellular traps. *Biol. Reprod.* 73, 1174–1181. doi:10.1095/biolreprod.105.045666
- Alkhodair, K., Almhanna, H., McGetrick, J., Gedair, S., Gallagher, M. E., Fernandez-Fuertes, B., et al. (2018). Siglec expression on the surface of human, bull and ram sperm. *Reproduction* 155, 361–371. doi:10.1530/REP-17-0475
- Altman, M. O., and Gagneux, P. (2019). Absence of Neu5Gc and presence of anti-Neu5Gc antibodies in humans-an evolutionary perspective. *Front. Immunol.* 10, 789. doi:10.3389/fimmu.2019.00789
- Angata, T. (2006). Molecular diversity and evolution of the Siglec family of cell-surface lectins. *Mol. Divers.* 10, 555–566. doi:10.1007/s11030-006-9029-1
- Angata, T., and Varki, A. (2002). Chemical diversity in the sialic acids and related α -keto acids: An evolutionary perspective. *Chem. Rev.* 102, 439–469. doi:10.1021/cr000407m
- Asahina, S., Sato, C., and Kitajima, K. (2004). Developmental expression of a sialyltransferase responsible for sialylation of cortical alveolus glycoprotein during oogenesis in rainbow trout (*Oncorhynchus mykiss*). *J. Biochem.* 136, 189–198. doi:10.1093/jb/mvh106
- Asahina, S., Sato, C., Matsuno, M., Matsuda, T., Colley, K., and Kitajima, K. (2006). Involvement of the α 2, 8-polysialyltransferases II/STX and IV/PST in the biosynthesis of polysialic acid chains on the O-linked glycoproteins in rainbow trout ovary. *J. Biochem.* 140, 687–701. doi:10.1093/jb/mvj200
- Avella, M. A., Baibakov, B., and Dean, J. (2014). A single domain of the ZP2 zona pellucida protein mediates gamete recognition in mice and humans. *J. Cell Biol.* 205, 801–809. doi:10.1083/jcb.201404025
- Avril, T., Floyd, H., Lopez, F., Vivier, E., and Crocker, P. R. (2004). The membrane-proximal immunoreceptor tyrosine-based inhibitory motif is critical for the inhibitory signaling mediated by siglecs-7 and -9, CD33-related Siglecs expressed on human monocytes and NK cells. *J. Immunol.* 173, 6841–6849. doi:10.4049/jimmunol.173.11.6841
- Bader, A., Douglas, E., and Chandler, A. (1998). A sperm chemoattractant is released from *Xenopus* egg jelly during spawning. *Dev. Biol. (N. Y.)* 198, 366–375. doi:10.1016/S0012-1606(98)80011-0
- Baibakov, B., Boggs, N. A., Yauger, B., Baibakov, G., and Dean, J. (2012). Human sperm bind to the N-terminal domain of ZP2 in humanized zonae pellucidae in transgenic mice. *J. Cell Biol.* 197, 897–905. doi:10.1083/jcb.201203062
- Banerjee, M., and Chowdhury, M. (1997). Localization of a 25 kDa human sperm surface protein: Its role in *in-vitro* human sperm capacitation. *Mol. Hum. Reprod.* 3, 109–114. doi:10.1093/molehr/3.2.109
- Barisone, G. A., Albertali, I. E., Sánchez, M., and Cabada, M. O. (2003). The envelopes of amphibian oocytes: Physiological modifications in *Bufo arenarum*. *Reprod. Biol. Endocrinol.* 1, 18–14. doi:10.1186/1477-7827-1-18
- Berner, N. J., and Ingermann, R. L. (1990). Role of sialic acid in exogenous protein accumulation and water retention by the egg jelly of the salamander *Ambystoma macrodactylum*. *J. Exp. Zool.* 256, 38–43. doi:10.1002/jez.1402560106
- Bessler, H., Singer, R., Raanani, P., Levinsky, H., Lahav, M., and Cohen, A. M. (1995). Interferon alpha-2b modulates beta-galactoside alpha-2, 6-sialyltransferase gene expression in rat testes. *Biol. Reprod.* 53, 1474–1477. doi:10.1095/biolreprod53.6.1474
- Bhakta, H. H., Refai, F. H., and Avella, M. A. (2019). The molecular mechanisms mediating mammalian fertilization. *Development* 146, dev176966. doi:10.1242/dev.176966
- Bhide, G. P., and Colley, K. J. (2017). Sialylation of N-glycans: Mechanism, cellular compartmentalization and function. *Histochem. Cell Biol.* 147, 149–174. doi:10.1007/s00418-016-1520-x
- Bleil, J. D., and Wassarman, P. M. (1988). Galactose at the nonreducing terminus of O-linked oligosaccharides of mouse egg zona pellucida glycoprotein ZP3 is essential for the glycoprotein's sperm receptor activity. *Proc. Natl. Acad. Sci. U. S. A.* 85, 6778–6782. doi:10.1073/pnas.85.18.6778
- Bleil, J. D., and Wassarman, P. M. (1990). Identification of a ZP3-binding protein on acrosome-intact mouse sperm by photoaffinity crosslinking. *Proc. Natl. Acad. Sci. U. S. A.* 87, 5563–5567. doi:10.1073/pnas.87.14.5563
- Bleil, J. D., and Wassarman, P. M. (1980). Mammalian sperm-egg interaction: Identification of a glycoprotein in mouse egg zonae pellucidae possessing receptor activity for sperm. *Cell* 20, 873–882. doi:10.1016/0092-8674(80)90334-7
- Boilard, M., Bailey, J., Collin, S., Dufour, M., and Sirard, M. A. (2002). Effect of bovine oviduct epithelial cell apical plasma membranes on sperm function assessed by a novel flow cytometric approach. *Biol. Reprod.* 67, 1125–1132. doi:10.1095/biolreprod67.4.1125
- Bornhöft, K. F., Goldammer, T., Rebl, A., and Galuska, S. P. (2018). Siglecs: A journey through the evolution of sialic acid-binding immunoglobulin-type lectins. *Dev. Comp. Immunol.* 86, 219–231. doi:10.1016/j.dci.2018.05.008
- Brandtzaeg, P. (1997). Mucosal immunity in the female genital tract. *J. Reprod. Immunol.* 36, 23–50. doi:10.1016/S0165-0378(97)00061-2
- Brinkmann, V., and Zychlinsky, A. (2007). Beneficial suicide: Why neutrophils die to make NETs. *Nat. Rev. Microbiol.* 5, 577–582. doi:10.1038/nrmicro1710
- Brun, R. B. (1974). Studies on fertilization in *Xenopus laevis*. *Biol. Reprod.* 11, 513–518. doi:10.1095/biolreprod11.5.513
- Buffone, M. G., Ijiri, T. W., Cao, W., Merdushev, T., Aghajanian, H. K., and Gerton, G. L. (2012). Heads or tails? Structural events and molecular mechanisms that promote mammalian sperm acrosomal exocytosis and motility. *Mol. Reprod. Dev.* 79, 4–18. doi:10.1002/mrd.21393
- Chakravarty, S., Kadunganattil, S., Bansal, P., Sharma, R. K., and Gupta, S. K. (2008). Relevance of glycosylation of human zona pellucida glycoproteins for their binding to capacitated human spermatozoa and subsequent induction of acrosomal exocytosis. *Mol. Reprod. Dev.* 75, 75–88. doi:10.1002/mrd.20726
- Chang, L. Y., Harduin-Lepers, A., Kitajima, K., Sato, C., Huang, C. J., Khoo, K. H., et al. (2009). Developmental regulation of oligosialylation in zebrafish. *Glycoconj. J.* 26, 247–261. doi:10.1007/s10719-008-9161-5
- Chang, L. Y., Teppa, E., Noel, M., Gilormini, P. A., Decloquement, M., Lion, C., et al. (2019). Novel zebrafish mono- α 2, 8-sialyltransferase (ST8Sia VIII): An evolutionary perspective of α 2, 8-sialylation. *Int. J. Mol. Sci.* 20, 622. doi:10.3390/ijms20030622
- Chen, G. Y., Tang, J., Zheng, P., and Liu, Y. (2009). CD24 and siglec-10 selectively repress tissue damage - induced immune responses. *Science* 323, 1722–1725. doi:10.1126/science.1168988
- Chen, X., and Varki, A. (2010). Advances in the biology and chemistry of sialic acids. *ACS Chem. Biol.* 5, 163–176. doi:10.1021/cb900266r
- Cheon, Y. P., and Kim, C. H. (2015). Impact of glycosylation on the unpaired functions of the sperm. *Clin. Exp. Reprod. Med.* 42, 77–85. doi:10.5653/cerm.2015.42.3.77

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Clark, G. F. (2010). The mammalian zona pellucida: A matrix that mediates both gamete binding and immune recognition? *Syst. Biol. Reprod. Med.* 56, 349–364. doi:10.3109/19396360903524812
- Clark, G. F. (2013). The role of carbohydrate recognition during human sperm-egg binding. *Hum. Reprod.* 28, 566–577. doi:10.1093/humrep/des447
- Close, B. E., and Colley, K. J. (1998). *In vivo* autopolysialylation and localization of the polysialyltransferases PST and STX. *J. Biol. Chem.* 273, 34586–34593. doi:10.1074/jbc.273.51.34586
- Cohen, M., and Varki, A. (2010). The sialome—far more than the sum of its parts. *OmicS. A J. Integr. Biol.* 14, 455–464. doi:10.1089/omi.2009.0148
- Cortés, P. P., Orihuela, P. A., Zúñiga, L. M., Velásquez, L. A., and Croxatto, H. B. (2004). Sperm binding to oviductal epithelial cells in the rat: Role of sialic acid residues on the epithelial surface and sialic acid-binding sites on the sperm surface. *Biol. Reprod.* 71, 1262–1269. doi:10.1095/bioreprod.104.027474
- Crocker, P. R., Paulson, J. C., and Varki, A. (2007). Siglecs and their roles in the immune system. *Nat. Rev. Immunol.* 7, 255–266. doi:10.1038/nri2056
- Cyster, J. G., Shotton, D. M., and Williams, A. F. (1991). The dimensions of the T lymphocyte glycoprotein leukosialin and identification of linear protein epitopes that can be modified by glycosylation. *EMBO J.* 10, 893–902. doi:10.1002/j.1460-2075.1991.tb08022.x
- Dall’Olio, F., Malagolini, N., Chiricolo, M., Trincherà, M., and Harduin-Lepers, A. (2014). The expanding roles of the Sd(a)/Cad carbohydrate antigen and its cognate glycosyltransferase B4GALNT2. *Biochim. Biophys. Acta* 1, 443–453. doi:10.1016/j.bbagen.2013.09.036
- de Martínez, N. R., Méndez, B. A., and Olavarria, J. M. (1975). Topographical distribution of sialic acids and related enzymes in toad oviduct. *Comp. Biochem. Physiol. B* 50, 603–607. doi:10.1016/0305-0491(75)90097-8
- Ded, L., Hwang, J. Y., Miki, K., Shi, H. F., and Chung, J. J. (2020). 3D *in situ* imaging of the female reproductive tract reveals molecular signatures of fertilizing spermatozoa in mice. *Elife* 9, e62043–e62051. doi:10.7554/eLife.62043
- Dell, A., Chalabi, S., Easton, R. L., Haslam, S. M., Sutton-Smith, M., Patankar, M. S., et al. (2003). Murine and human zona pellucida 3 derived from mouse eggs express identical O-glycans. *Proc. Natl. Acad. Sci. U. S. A.* 100, 15631–15636. doi:10.1073/pnas.2635507100
- Dell, A., Morris, H. R., Easton, R. L., Patankar, M., and Clark, G. F. (1999). The glycobiology of gametes and fertilization. *Biochim. Biophys. Acta* 1473, 196–205. doi:10.1016/S0304-4165(99)00179-8
- Della Giovampaola, C., Flori, F., Sabatini, L., Incerti, L., La Sala, G. B., Rosati, F., et al. (2001). Surface of human sperm bears three differently charged CD52 forms, two of which remain stably bound to sperm after capacitation. *Mol. Reprod. Dev.* 60, 89–96. doi:10.1002/mrd.1065
- Demott, R. P., Lefebvre, R., and Suarez, S. S. (1995). Carbohydrates mediate the adherence of hamster sperm to oviductal epithelium. *Biol. Reprod.* 52, 1395–1403. doi:10.1095/bioreprod52.6.1395
- Di, W., Fujita, A., Hamaguchi, K., Delannoy, P., Sato, C., and Kitajima, K. (2017). Diverse subcellular localizations of the insect CMP-sialic acid synthetases. *Glycobiology* 27, 329–341. doi:10.1093/glycob/cww128
- Drickamer, K., and Taylor, M. E. (1998). Evolving views of protein glycosylation. *Trends biochem. Sci.* 23, 321–324. doi:10.1016/S0968-0004(98)01246-8
- Ertunc, N., Sato, C., and Kitajima, K. (2020). Sialic acid sulfation is induced by the antibiotic treatment in mammalian cells. *Biosci. Biotechnol. Biochem.* 84, 2311–2318. doi:10.1080/09168451.2020.1792763
- Esko, J. D., and Selleck, S. B. (2002). Order out of chaos: Assembly of ligand binding sites in heparan sulfate. *Annu. Rev. Biochem.* 71, 435–471. doi:10.1146/annurev.biochem.71.110601.135458
- Evans, J. P. (2020). Preventing polyspermy in mammalian eggs—Contributions of the membrane block and other mechanisms. *Mol. Reprod. Dev.* 87, 341–349. doi:10.1002/mrd.23331
- Fahrenkamp, E., Algarra, B., and Jovine, L. (2020). Mammalian egg coat modifications and the block to polyspermy. *Mol. Reprod. Dev.* 87, 326–340. doi:10.1002/mrd.23320
- Fazeli, A., Elliott, R. M. A., Duncan, A. E., Moore, A., Watson, P. F., and Holt, W. V. (2003). *In vitro* maintenance of boar sperm viability by a soluble fraction obtained from oviductal apical plasma membrane preparations. *Reproduction* 125, 509–517. doi:10.1530/rep.0.1250509
- Feng, J. M., Tian, H. F., Hu, Q. M., Meng, Y., and Xiao, H. B. (2018). Evolution and multiple origins of zona pellucida genes in vertebrates. *Biol. Open* 7, bio036137–10. doi:10.1242/bio.036137
- Fernandez-Fuertes, B., Blanco-Fernandez, A., Reid, C. J., Meade, K. G., Fair, S., and Lonergan, P. (2018). Removal of sialic acid from bull sperm decreases motility and mucus penetration ability but increases zona pellucida binding and polyspermic penetration *in vitro*. *Reproduction* 155, 481–492. doi:10.1530/REP-17-0429
- Florea, D., Maes, E., Guérardel, Y., Page, A., Zanetta, J. P., Cogalniceanu, D., et al. (2006). Structure elucidation of NeuAc, NeuGc and kdn-containing O-glycans released from *Triturus alpestris* oviductal mucins. Characterization of the poly LacdiNAc sequence: HSO3(4)(GalNAcβ1-4GlcNAcβ1-3)1-3GalNAcβ1-4(GlcNAcβ1-3)0-1GlcNAcβ1-6GalNAc-ol. *Glycoconj. J.* 23, 377–399. doi:10.1007/s10719-006-6126-4
- Focarelli, R., Giuffrida, A., and Rosati, F. (1995). Changes in the sialylglycoconjugate distribution on the human sperm surface during *in-vitro* capacitation: Partial purification of a 20 kDa sialylglycoprotein of capacitated spermatozoa. *Hum. Reprod.* 1, 2755–2759. doi:10.1093/oxfordjournals.humrep.a135787
- Fraser, L. R., and Lane, M. R. (1987). Capacitation- and fertilization-related alterations in mouse sperm oxygen consumption. *J. Reprod. Fertil.* 81, 385–393. doi:10.1530/jrf.0.0810385
- Froman, D. P., and Thursam, K. A. (1994). Desialylation of the rooster sperm’s glycolyx decreases sperm sequestration following intravaginal insemination of the hen. *Biol. Reprod.* 50, 1094–1099. doi:10.1095/bioreprod50.5.1094
- Gagneux, P., Aebi, M., and Varki, A. (2017). Evolution of glycan diversity. *Essentials Glycobiol.* doi:10.1101/glycobiology.3e.020
- Gahlay, G., Gauthier, L., Baibakov, B., Epifano, O., and Dean, J. (2010). Gamete recognition in mice depends on the cleavage status of an egg’s zona pellucida protein. *Sci. (New York, N.Y.)* 329 (5988), 216–219. doi:10.1126/science.1188178
- Gallo, A., and Costantini, M. (2012). Glycobiology of reproductive processes in marine animals: The state of the art. *Mar. Drugs* 10, 2861–2892. doi:10.3390/md10122861
- Galuska, S. P., Rollenhagen, M., Kaup, M., Eggers, K., Oltmann-Norden, I., Schiff, M., et al. (2010). Synaptic cell adhesion molecule SynCAM 1 is a target for polysialylation in postnatal mouse brain. *Proc. Natl. Acad. Sci. U. S. A.* 107, 10250–10255. doi:10.1073/pnas.0912103107
- Giapocuzzi, E., Bresciani, R., Schauer, R., Monti, E., and Borsani, G. (2012). New insights on the sialidase protein family revealed by a phylogenetic analysis in metazoa. *PLoS one* 7, e44193. doi:10.1371/journal.pone.0044193
- Glasner, A., Zurunic, A., Meningher, T., Lenac Rovis, T., Tsukerman, P., Bar-On, Y., et al. (2012). Elucidating the mechanisms of influenza virus recognition by Ncr1. *PLoS One* 7, e36837. doi:10.1371/journal.pone.0036837
- Gook, D. A., Edgar, D. H., Borg, J., and Martic, M. (2008). Detection of zona pellucida proteins during human folliculogenesis. *Hum. Reprod.* 23, 394–402. doi:10.1093/humrep/dem373
- Greve, J. M., and Wassarman, P. M. (1985). Mouse egg extracellular coat is a matrix of interconnected filaments possessing a structural repeat. *J. Mol. Biol.* 181, 253–264. doi:10.1016/0022-2836(85)90089-0
- Guérardel, Y., Chang, L. Y., Maes, E., Huang, C. J., and Khoo, K. H. (2006). Glycomic survey mapping of zebrafish identifies unique sialylation pattern. *Glycobiology* 16, 244–257. doi:10.1093/glycob/cwj062
- Guérardel, Y., Kol, O., Maes, E., Lefebvre, T., Boilly, B., Davril, M., et al. (2000). O-glycan variability of egg-jelly mucins from *Xenopus laevis*: Characterization of four phenotypes that differ by the terminal glycosylation of their mucins. *Biochem. J.* 352, 449–463. doi:10.1042/bj3520449
- Guerrero, A., Gadêlha, H., Ramirez-Gomez, H. V., Ramirez, R., Beltran, C., and Tuval, I. (2020). “Motility and guidance of sea urchin sperm,” in *Reproduction in aquatic animals* (Singapore: Springer), 249–276. doi:10.1007/978-981-15-2290-1_13
- Gupta, S. K., and Bhandari, B. (2011). Acrosome reaction: Relevance of zona pellucida glycoproteins. *Asian J. Androl.* 13, 97–105. doi:10.1038/aja.2010.72
- Gupta, S. K. (2021). Human zona pellucida glycoproteins: Binding characteristics with human spermatozoa and induction of acrosome reaction. *Front. Cell Dev. Biol.* 9, 619868. doi:10.3389/fcell.2021.619868
- Gupta, S. K. (2018). The human egg’s zona pellucida. *Curr. Top. Dev. Biol.* 130, 379–411. doi:10.1016/bs.ctdb.2018.01.001
- Gurushankara, H. P., Krishnamurthy, S. V., and Vasudev, V. (2012). Changes in sialic acid content of jelly coat in pesticide-exposed frog eggs and their influence on fertilization. *Adv. Exp. Med. Biol.* 749, 329–336. doi:10.1007/978-1-4614-3381-1_22
- Hachem, N. E., Humpfle, L., Simon, P., Kaese, M., Weinhold, B., Günther, J., et al. (2021). The loss of polysialic acid impairs the contractile phenotype of peritubular smooth muscle cells in the postnatal testis. *Cells* 6, 1347. doi:10.3390/cells10061347
- Hakomori, S. (2004). Carbohydrate-to-carbohydrate interaction, through glycosynapse, as a basis of cell recognition and membrane organization. *Glycoconj. J.* 21, 125–137. doi:10.1023/B:GLYC.0000044844.95878.cf

- Han, L., Monné, M., Okumura, H., Schwend, T., Cherry, A. L., Flot, D., et al. (2010). Insights into egg coat assembly and egg-sperm interaction from the X-ray structure of full-length ZP3. *Cell* 143, 404–415. doi:10.1016/j.cell.2010.09.041
- Harduin-Lepers, A. (2010). Comprehensive analysis of sialyltransferases in vertebrate genomes. *Glycobiol. Insights* 2, 29–61. doi:10.4137/gbi.s3123
- Harduin-Lepers, A., Mollicone, R., Delannoy, P., and Oriol, R. (2005). The animal sialyltransferases and sialyltransferase-related genes: A phylogenetic approach. *Glycobiology* 15, 805–817. doi:10.1093/glycob/cwi063
- Harduin-Lepers, A., Petit, D., Mollicone, R., Delannoy, P., Petit, J. M., and Oriol, R. (2008). Evolutionary history of the alpha2, 8-sialyltransferase (ST8Sia) gene family: Tandem duplications in early deuterostomes explain most of the diversity found in the vertebrate ST8Sia genes. *BMC Evol. Biol.* 8, 258. doi:10.1186/1471-2148-8-258
- Harduin-Lepers, A., Vallejo-Ruiz, V., Krzewinski-Recchi, M. A., Samyn-Petit, B., Julien, S., and Delannoy, P. (2001). The human sialyltransferase family. *Biochimie* 83, 727–737. doi:10.1016/s0300-9084(01)01301-3
- Hasegawa, A., Kanazawa, N., Koyama, K., and Koyama, K. (2007). Contribution of zona pellucida proteins to ovarian folliculogenesis. *Reproductive Immunol. Biol.* 22, 1–7. doi:10.3192/jsirib.22.1
- Hedrick, J. L. (2008). Anuran and pig egg zona pellucida glycoproteins in fertilization and early development. *Int. J. Dev. Biol.* 52, 683–701. doi:10.1387/ijdb.082580jh
- Henrissat, B., and Bairoch, A. (1996). Updating the sequence-based classification of glycosyl hydrolases. *Biochem. J.* 316, 695–696. doi:10.1042/bj3160695
- Hino, T., Muro, Y., Tamura-Nakano, M., Okabe, M., Tateno, H., and Yanagimachi, R. (2016). The behavior and acrosomal status of mouse spermatozoa *in vitro*, and within the oviduct during fertilization after natural mating. *Biol. Reprod.* 95, 50. doi:10.1095/biolreprod.116.140400
- Hirohashi, N., Gerton, G. L., and Buffone, M. G. (2011). Video imaging of the sperm acrosome reaction during *in vitro* fertilization. *Commun. Integr. Biol.* 4, 471–476. doi:10.4161/cib.4.4.15636
- Hirohashi, N., Kamei, N., Kubo, H., Sawada, H., Matsumoto, M., and Hoshi, M. (2008). Egg and sperm recognition systems during fertilization. *Dev. Growth Differ.* 50, S221–S238. doi:10.1111/j.1440-169X.2008.01017.x
- Hirohashi, N., and Vacquier, V. D. (2002). Egg fucose sulfate polymer, sialoglycan, and speract all trigger the sea urchin sperm acrosome reaction. *Biochem. Biophys. Res. Commun.* 296, 833–839. doi:10.1016/S0006-291X(02)00941-5
- Hirose, M., Honda, A., Fulka, H., Tamura-Nakano, M., Matoba, S., Tomishima, T., et al. (2020). Erratum: Acrosin is essential for sperm penetration through the zona pellucida in hamsters. *Proc. Natl. Acad. Sci. U. S. A.* 117, 2513–2518. doi:10.1073/pnas.2018022117
- Hoffmann, D. S., and Killian, G. J. (1981). Isolation of epithelial cells from the corpus epididymidis and analysis for glycylphosphorylcholine, sialic acid, and protein. *J. Exp. Zool.* 217, 93–102. doi:10.1002/jez.1402170110
- Holt, W. V. (1980). Surface-bound sialic acid on ram and bull spermatozoa: Deposition during epididymal transit and stability during washing. *Biol. Reprod.* 23, 847–857. doi:10.1095/biolreprod23.4.847
- Hoodbhoy, T., Joshi, S., Boja, E. S., Williams, S. A., Stanley, P., and Dean, J. (2005). Human sperm do not bind to rat zonae pellucidae despite the presence of four homologous glycoproteins. *J. Biol. Chem.* 280, 12721–12731. doi:10.1074/jbc.M413569200
- Horne-Badovinac, S. (2020). The *Drosophila* micropyle as a system to study how epithelia build complex extracellular structures. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 375, 20190561. doi:10.1098/rstb.2019.0561
- Inoue, S., and Inoue, Y. (1997). “Fish glycoproteins,” in *Glycoproteins II*. Editors J. Montreuil, J. F. G. Vlieghehart, and H. Schachter (Elsevier Science B.V.), 143–188.
- Inoue, S., and Iwasaki, M. (1978). Isolation of a novel glycoprotein from the eggs of rainbow trout: Occurrence of disialosyl groups on all carbohydrate chains. *Biochem. Biophys. Res. Commun.* 83, 1018–1023. doi:10.1016/0006-291x(78)91497-3
- Intra, J., Cenni, F., Pavesi, G., Pasini, M., and Perotti, M. E. (2009). Interspecific analysis of the glycosidases of the sperm plasma membrane in *Drosophila*. *Mol. Reprod. Dev.* 76, 85–100. doi:10.1002/mrd.20932
- Iwasaki, M., and Inoue, S. (1985). Structures of the carbohydrate units of polysialoglycoproteins isolated from the eggs of four species of salmonid fishes. *Glycoconj. J.* 2, 209–228. doi:10.1007/BF01049269
- Jaffe, L. A., and Cross, N. L. (1986). Electrical regulation of sperm-egg fusion. *Annu. Rev. physiology* 48, 191–200. doi:10.1146/annurev.ph.48.030186.001203
- James, W. M., and Agnew, W. S. (1987). Multiple oligosaccharide chains in the voltage-sensitive Na channel from *Electrophorus electricus*: Evidence for α -2, 8-linked polysialic acid. *Biochem. Biophys. Res. Commun.* 148, 817–826. doi:10.1016/0006-291X(87)90949-1
- Jin, M., Fujiwara, E., Kakiuchi, Y., Okabe, M., Satouh, Y., Baba, S. A., et al. (2011). Most fertilizing mouse spermatozoa begin their acrosome reaction before contact with the zona pellucida during *in vitro* fertilization. *Proc. Natl. Acad. Sci. U. S. A.* 108, 4892–4896. doi:10.1073/pnas.1018202108
- Kadirvel, G., Machado, S. A., Korneli, C., Collins, E., Miller, P., Bess, K. N., et al. (2012). Porcine sperm bind to specific 6-sialylated biantennary glycans to form the oviduct reservoir. *Biol. Reprod.* 87, 147. doi:10.1095/biolreprod.112.103879
- Kambara, Y., Shiba, K., Yoshida, M., Sato, C., Kitajima, K., and Shingyoji, C. (2011). Mechanism regulating Ca²⁺-dependent mechanosensory behaviour in sea urchin spermatozoa. *Cell Struct. Funct.* 36, 69–82. doi:10.1247/csf.10020
- Kanamori, A., Inoue, S., Iwasaki, M., Kitajima, K., Kawai, G., Yokoyama, S., et al. (1990). Deaminated neuraminic acid-rich glycoprotein of rainbow trout egg vitelline envelope. Occurrence of a novel α -2, 8-linked oligo(deaminated neuraminic acid) structure in O-linked glycan chains. *J. Biol. Chem.* 265, 21811–21819. doi:10.1016/s0021-9258(18)45812-7
- Kaneko, S., Oshio, S., Kobayashi, T., Iizuka, R., and Mohri, H. (1984). Human X- and Y-bearing sperm differ in cell surface sialic acid content. *Biochem. Biophys. Res. Commun.* 124, 950–955. doi:10.1016/0006-291x(84)91050-7
- Kapli, P., Natsidis, P., Leite, D. J., Fursman, M., Jeffrie, N., Rahman, I. A., et al. (2021). Lack of support for Deuterostomia prompts reinterpretation of the first Bilateria. *Sci. Adv.* 19, eabe2741. doi:10.1126/sciadv.abe2741
- Katsumata, T., Noguchi, S., YoNezawaN.TanokuraM.and NakanoM. (1996). Structural characterization of the N-linked carbohydrate chains of the zona pellucida glycoproteins from bovine ovarian and fertilized eggs. *Eur. J. Biochem.* 240, 448–453. doi:10.1111/j.1432-1033.1996.0448h.x
- Khosravi, F., Michel, V., Galuska, C. E., Bhushan, S., Christian, P., Schuppe, H. C., et al. (2016). Desialylation of spermatozoa and epithelial cell glycoalyx is a consequence of bacterial infection of the epididymis. *J. Biol. Chem.* 291, 17717–17726. doi:10.1074/jbc.M116.718072
- Kiermaier, E., Moussion, C., Veldkamp, C. T., Gerardy-Schahn, R., De Vries, I., Williams, L. G., et al. (2016). Polysialylation controls dendritic cell trafficking by regulating chemokine recognition. *Science* 351, 186–190. doi:10.1126/science.aad0512
- Kim, B. H., Ju, W. S., Kim, J. S., Kim, S. U., Park, S. J., Ward, S. M., et al. (2020). Effects of gangliosides on spermatozoa, oocytes, and preimplantation embryos. *Int. J. Mol. Sci.* 21, E106–E114. doi:10.3390/ijms21010106
- Kimura, Y., Manabe, N., Nishihara, S., Matsushita, H., Tajima, C., Wada, S., et al. (1999). Up-regulation of the alpha2, 6-sialyltransferase messenger ribonucleic acid increases glycoconjugates containing alpha2, 6-linked sialic acid residues in granulosa cells during follicular atresia of porcine ovaries. *Biol. Reprod.* 60, 1475–1482. doi:10.1095/biolreprod60.6.1475
- Kitajima, K., Inoue, Y., and Inoue, S. (1986). Polysialoglycoproteins of Salmonidae fish eggs. Complete structure of 200-kDa polysialoglycoprotein from the unfertilized eggs of rainbow trout (*Salmo gairdneri*). *J. Biol. Chem.* 261, 5262–5269. doi:10.1016/s0021-9258(19)57208-8
- Kitazume, S., Kitajima, K., Inoue, S., Haslam, S. M., Morris, H. R., Dell, A., et al. (1996). The occurrence of novel 9-O-sulfated N-glycolylneuraminic Acid-capped α 2–5-Oglycolyl-linked oligo/polyNeu5Gc chains in sea urchin egg cell surface glycoprotein: Identification of a new chain termination signal for polysialyltransferase. *J. Biol. Chem.* 271, 6694–6701. doi:10.1074/jbc.271.12.6694
- Kitazume, S., Kitajima, K., Inoue, S., Troy, F. A., Cho, J. W., Lennarz, W. J., et al. (1994). Identification of polysialic acid-containing glycoprotein in the jelly coat of sea urchin eggs. Occurrence of a novel type of polysialic acid structure. *J. Biol. Chem.* 269, 22712–22718. doi:10.1016/s0021-9258(17)31704-0
- Koles, K., Irvine, K. D., and Panin, V. M. (2004). Functional characterization of *Drosophila* sialyltransferase. *J. Biol. Chem.* 279, 4346–4357. doi:10.1074/jbc.M309912200
- Kratz, E. M., Faundez, R., and Kątnik-Prastowska, I. (2011). Fucose and sialic acid expressions in human seminal fibronectin and α [1]-acid glycoprotein associated with leukocytospermia of infertile men. *Dis. Markers* 31, 317–325. doi:10.3233/DMA-2011-0846
- Kratz, E. M., Kaluza, A., Zimmer, M., and Ferens-Sieczkowska, M. (2015). The analysis of sialylation, N-glycan branching, and expression of O-glycans in seminal plasma of infertile men. *Dis. Markers* 2015, 1–11. doi:10.1155/2015/941871
- Krzewinski-Recchi, M. A., Julien, S., Juliant, S., Teinturier-Lelièvre, M., Samyn-Petit, B., Montiel, M. D., et al. (2003). Identification and functional expression of a second human beta-galactoside alpha2, 6-sialyltransferase, ST6Gal II. *Eur. J. Biochem.* 270, 950–961. doi:10.1046/j.1432-1033.2003.03458.x
- Kubo, H., Shiga, K., Harada, Y., and Iwao, Y. (2010). Analysis of a sperm surface molecule that binds to a vitelline envelope component of *Xenopus laevis* eggs. *Mol. Reprod. Dev.* 77, 728–735. doi:10.1002/mrd.21211
- Kurosawa, N., Inoue, M., Yoshida, Y., and Tsuji, S. (1996). Molecular cloning and genomic analysis of mouse Galbeta1, 3GalNAc-specific GalNAc alpha2, 6-sialyltransferase. *J. Biol. Chem.* 271, 15109–15116. doi:10.1074/jbc.271.25.15109

- La Spina, F. A., Puga Molina, L. C., Romarowski, A., Vitale, A. M., Falzone, T. L., Krapf, D., et al. (2016). Mouse sperm begin to undergo acrosomal exocytosis in the upper isthmus of the oviduct. *Dev. Biol.* 411, 172–182. doi:10.1016/j.ydbio.2016.02.006
- Lassalle, B., and Testart, J. (1994). Human zona pellucida recognition associated with removal of sialic acid from human sperm surface. *J. Reprod. Fertil.* 101, 703–711. doi:10.1530/jrf.0.1010703
- Lay, K. M., Nakada, T., and Tatemoto, H. (2013). Involvement of N-glycosylation of zona glycoproteins during meiotic maturation in sperm-zona pellucida interactions of porcine denuded oocytes. *Anim. Sci. J.* 84, 8–14. doi:10.1111/j.1740-0929.2012.01027.x
- Lay, K. M., Oshiro, R., Arasaki, C., Ashizawa, K., and Tatemoto, H. (2011). Role of acidification elicited by sialylation and sulfation of zona glycoproteins during oocyte maturation in porcine sperm-zona pellucida interactions. *J. reproduction Dev.* 57, 744–751. doi:10.1262/jrd.11-057h
- Lefebvre, R., Lo, M. C., and Suarez, S. S. (1997). Bovine sperm binding to oviductal epithelium involves fucose recognition. *Biol. Reprod.* 56, 1198–1204. doi:10.1095/biolreprod56.5.1198
- Lefèvre, L., Conner, S. J., Salpekar, A., Olufowobi, O., Ashton, P., Pavlovic, B., et al. (2004). Four zona pellucida glycoproteins are expressed in the human. *Hum. Reprod.* 19, 1580–1586. doi:10.1093/humrep/deh301
- Legendre, M., Linhart, O., and Billard, R. (1996). Spawning and management of gametes, fertilized eggs and embryos in Siluroidei. *Aquat. Living Resour.* 9, 59–80. doi:10.1051/alr:1996042
- Lei, T., Moos, S., Klug, J., Aslani, F., Bhushan, S., Wahle, E., et al. (2018). Galectin-1 enhances TNF α -induced inflammatory responses in Sertoli cells through activation of MAPK signalling. *Sci. Rep.* 8, 3741. doi:10.1038/s41598-018-22135-w
- Lepers, A., Shaw, L., Cacan, R., Schauer, R., Montreuil, J., Verbert, A., et al. (1989). Transport of CMP-N-glycolylneuraminic acid into mouse liver Golgi vesicles. *FEBS Lett.* 250, 245–250. doi:10.1016/0014-5793(89)80731-8
- Lepers, A., Shaw, L., Schneckeburger, P., Cacan, R., Verbert, A., and Schauer, R. (1990). A study on the regulation of N-glycolylneuraminic acid biosynthesis and utilization in rat and mouse liver. *Eur. J. Biochem.* 193, 715–723. doi:10.1111/j.1432-1033.1990.tb19391.x
- Li, B., Russell, S. C., Zhang, J., Hedrick, J. L., and Lebrilla, C. B. (2011). Structure determination by MALDI-IRMPD mass spectrometry and exoglycosidase digestions of O-linked oligosaccharides from *Xenopus borealis* egg jelly. *Glycobiology* 21, 877–894. doi:10.1093/glycob/cwr003
- Li, F., and Ding, J. (2019). Sialylation is involved in cell fate decision during development, reprogramming and cancer progression. *Protein Cell* 10, 550–565. doi:10.1007/s13238-018-0597-5
- Li, P., Guo, W., Yue, H., Li, C., Du, H., Qiao, X., et al. (2017). Variability in the protein profiles in spermatozoa of two sturgeon species. *PLoS One* 12, e0186003–e0186020. doi:10.1371/journal.pone.0186003
- Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P. M., and Henrissat, B. (2014). The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* 42, D490–D495. doi:10.1093/nar/gkt1178
- Longo, F. J., Lynn, J. W., McCulloh, D. H., and Chambers, E. L. (1986). Correlative ultrastructural and electrophysiological studies of sperm-egg interactions of the sea urchin, *Lytechinus variegatus*. *Dev. Biol.* 118, 155–166. doi:10.1016/0012-1606(86)90083-7
- Lopo, A. C., Glabe, C. G., Lennarz, W. J., and Vacquier, V. D. (1982). Sperm-egg binding events during sea urchin fertilization. *Ann. N. Y. Acad. Sci.* 383, 405–425. doi:10.1111/j.1749-6632.1982.tb23181.x
- Ma, F., Wu, D., Deng, L., Secrest, P., Zhao, J., Varki, N., et al. (2012). Sialidases on mammalian sperm mediate deciduous sialylation during capacitation. *J. Biol. Chem.* 287, 38073–38079. doi:10.1074/jbc.M112.380584
- Ma, X., Pan, Q., Feng, Y., Choudhury, B. P., Ma, Q., Gagneux, P., et al. (2016). Sialylation facilitates the maturation of mammalian sperm and affects its survival in female uterus. *Biol. Reprod.* 94 (123), 1–10. doi:10.1095/biolreprod.115.137810
- Maehashi, E., Sato, C., Ohta, K., Harada, Y., Matsuda, T., Hirohashi, N., et al. (2003). Identification of the sea urchin 350-kDa sperm-binding protein as a new sialic acid-binding lectin that belongs to the heat shock protein 110 family: Implication of its binding to gangliosides in sperm lipid rafts in fertilization. *J. Biol. Chem.* 278, 42050–42057. doi:10.1074/jbc.M307493200
- Mandich, A., Massari, A., Bottero, S., and Marino, G. (2002). Histological and histochemical study of female germ cell development in the dusky grouper *Epinephelus marginatus* (Lowe, 1834). *Eur. J. Histochem.* 46, 87–100. doi:10.4081/1657
- McCulloh, D. H., and Chambers, E. L. (1992). Fusion of membranes during fertilization. Increases of the sea urchin egg's membrane capacitance and membrane conductance at the site of contact with the sperm. *J. Gen. Physiol.* 99, 137–175. doi:10.1085/jgp.99.2.137
- Mengerink, K. J., and Vacquier, V. D. (2001). Glycobiology of sperm-egg interactions in deuterostomes. *Glycobiology* 11, 37R–43R. doi:10.1093/glycob/11.4.37R
- Miranda, P. V., Gonzalez-Echeverria, F., Marín-Briggiler, C. I., Brandelli, A., Blaquier, J. A., and Tezón, J. G. (1997). Glycosidic residues involved in human sperm-zona pellucida binding *in vitro*. *Mol. Hum. Reprod.* 3, 399–404. doi:10.1093/molehr/3.5.399
- Miwa, N. (2015). Protein-carbohydrate interaction between sperm and the egg-coating envelope and its regulation by dicalcin, a *Xenopus laevis* zona pellucida protein-associated protein. *Molecules* 20, 9468–9486. doi:10.3390/molecules20059468
- Miyata, S., Sato, C., Kitamura, S., Toriyama, M., and Kitajima, K. (2004). A major flagellum sialoglycoprotein in sea urchin sperm contains a novel polysialic acid, an α 2, 9-linked poly-N-acetylneuraminic acid chain, capped by an 8-O-sulfated sialic acid residue. *Glycobiology* 14, 827–840. doi:10.1093/glycob/cwh100
- Miyata, S., Sato, C., Kumita, H., Toriyama, M., Vacquier, V. D., and Kitajima, K. (2006). Flagellin sialin: A novel sulfated α 2, 9-linked polysialic acid glycoprotein of sea urchin sperm flagella. *Glycobiology* 16, 1229–1241. doi:10.1093/glycob/cwl036
- Miyata, S., Yamakawa, N., Toriyama, M., Sato, C., and Kitajima, K. (2011). Co-expression of two distinct polysialic acids, α 2, 8- and α 2, 9-linked polymers of N-acetylneuraminic acid, in distinct glycoproteins and glycolipids in sea urchin sperm. *Glycobiology* 21, 1596–1605. doi:10.1093/glycob/cwr081
- Monné, M., and Jovine, L. (2011). A structural view of egg coat architecture and function in fertilization. *Biol. Reprod.* 85, 661–669. doi:10.1095/biolreprod.111.092098
- Monti, E., Bonten, E., D'Azzo, A., Bresciani, R., Venerando, B., Borsani, G., et al. (2010). Sialidases in vertebrates: A family of enzymes tailored for several cell functions. *Adv. Carbohydr. Chem. Biochem.* 64, 403–479. doi:10.1016/S0065-2318(10)64007-3
- Moros-Nicolás, C., Chevret, P., Jiménez-Movilla, M., Algarra, B., Cots-Rodríguez, P., González-Brusi, L., et al. (2021). New insights into the mammalian egg zona pellucida. *Int. J. Mol. Sci.* 22, 3276. doi:10.3390/ijms22063276
- Mozing, N. M., and Hedrick, J. L. (1999). Distribution of lectin binding sites in *Xenopus laevis* egg jelly. *Dev. Biol.* 210, 428–439. doi:10.1006/dbio.1999.9289
- Mühlenhoff, M., Rollenhagen, M., Werneburg, S., Gerardy-Schahn, R., and Hildebrandt, H. (2013). Polysialic acid: Versatile modification of NCAM, SynCAM 1 and neuropilin-2. *Neurochem. Res.* 38, 1134–1143. doi:10.1007/s11064-013-0979-2
- Muro, Y., Hasuwa, H., Isotani, A., Miyata, H., Yamagata, K., Ikawa, M., et al. (2016). Behavior of mouse spermatozoa in the female reproductive tract from soon after mating to the beginning of fertilization. *Biol. Reprod.* 94, 80. doi:10.1095/biolreprod.115.135368
- Nadano, D., Iwasaki, M., Endo, S., Kitajima, K., Inoue, S., and Inoue, Y. (1986). A naturally occurring deaminated neuraminic acid, 3-deoxy-D-glycero-D-galactono-nonulosonic acid (KDN). Its unique occurrence at the nonreducing ends of oligosialyl chains in polysialoglycoprotein of rainbow trout eggs. *J. Biol. Chem.* 261, 11550–11557. doi:10.1016/s0021-9258(18)67278-3
- Nakano, K., Nakano, T., Ahn, D. U., and Sim, J. S. (1994). Sialic acid contents in chicken eggs and tissues. *Can. J. Anim. Sci.* 74, 601–606. doi:10.4141/cjas94-088
- Nakano, M., Yonezawa, N., Hatanaka, Y., and Noguchi, S. (1996). Structure and function of the N-linked carbohydrate chains of pig zona pellucida glycoproteins. *J. Reprod. Fertil. Suppl.* 50, 25–34.
- Nakano, M., and Yonezawa, N. (2001). Localization of sperm ligand carbohydrate chains in pig zona pellucida glycoproteins. *Cells, tissues, organs* 168, 65–75. doi:10.1159/000016807
- Nasu, K., and Narahara, H. (2010). Pattern recognition via the toll-like receptor system in the human female genital tract. *Mediat. Inflamm.* 2010, 976024. doi:10.1155/2010/976024
- Neubauer, D. M., and Wolfner, M. F. (1999). 3 wise, winsome, or weird? Mechanisms of sperm storage in female animals. *Curr. Top. Dev. Biol.* 41, 67–97. doi:10.1016/S0070-2153(08)60270-7
- Nguyen, T., Lee, S., Yang, Y.-A., Ahn, C., Sim, J. H., Kei, T. G., et al. (2020). The role of 9-O-acetylated glycan receptor moieties in the typhoid toxin binding and intoxication. *PLoS Pathog.* 16, e1008336. doi:10.1371/journal.ppat.1008336
- Nishimura, K., Dioguardi, E., Nishio, S., Villa, A., Han, L., Matsuda, T., et al. (2019). Molecular basis of egg coat cross-linking sheds light on ZP1-associated female infertility. *Nat. Commun.* 10, 3086. doi:10.1038/s41467-019-10931-5
- Nishio, S., Kohno, Y., Iwata, Y., Arai, M., Okumura, H., Oshima, K., et al. (2014). Glycosylated chicken ZP2 accumulates in the egg coat of immature oocytes and remains localized to the germinal disc region of mature eggs. *Biol. Reprod.* 91, 107. doi:10.1095/biolreprod.114.119826

- Nishio, S., Okumura, H., and Matsuda, T. (2018). Egg-coat and zona pellucida proteins of chicken as a typical species of aves. *Curr. Top. Dev. Biol.* 130, 307–329. doi:10.1016/bs.ctdb.2018.02.008
- Noguchi, S., Hatanaka, Y., Tobita, T., and Nakano, M. (1992). Erratum: Structural analysis of the N-linked carbohydrate chains of the 55-kDa glycoprotein family (PZP3) from porcine zona pellucida. *Eur. J. Biochem. J. Biochem.* 204207 (1), 1130. doi:10.1111/j.1432-1033.1992.tb17151.x_3
- Oehninger, S., Patankar, M., Seppala, M., and Clark, G. F. (1998). Involvement of selectin-like carbohydrate binding specificity in human gamete interaction. *Andrologia* 30, 269–274. doi:10.1111/j.1439-0272.1998.tb01170.x
- Okabe, M. (2015). Mechanisms of fertilization elucidated by gene-manipulated animals. *Asian J. Androl.* 17, 646–652. doi:10.4103/1008-682X.153299
- Okabe, M. (2013). The cell biology of mammalian fertilization. *Development* 140, 4471–4479. doi:10.1242/dev.090613
- Olson, J. H., Xiang, X., Ziegert, T., Kittelson, A., Rawls, A., Bieber, A. L., et al. (2001). Allurin, a 21-kDa sperm chemoattractant from *Xenopus* egg jelly, is related to mammalian sperm-binding proteins. *Proc. Natl. Acad. Sci. U. S. A.* 98, 11205–11210. doi:10.1073/pnas.211316798
- Ortiz-Delgado, J. B., Porcelloni, S., Fossi, C., and Sarasquete, C. (2008). Caracterización histoquímica de ovocitos del pez espada, <i>Xiphias gladius</i>. *Sci. Mar.* 72, 549–564. doi:10.3989/scimar.2008.72n3549
- Padler-Karavani, V., Yu, H., Cao, H., Chokhawala, H., Karp, F., Varki, N., et al. (2008). Diversity in specificity, abundance, and composition of anti-Neu5Gc antibodies in normal humans: Potential implications for disease. *Glycobiology* 10, 818–830. doi:10.1093/glycob/cwn072
- Pang, P. C., Chiu, P. C. N., Lee, C. L., Chang, L. Y., Panico, M., Morris, H. R., et al. (2011). Human sperm binding is mediated by the sialyl-lewisx Oligosaccharide on the zona pellucida. *Science* 333, 1761–1764. doi:10.1126/science.1207438
- Pang, P. C., Tissot, B., Drobnis, E. Z., Sutovsky, P., Morris, H. R., Clark, G. F., et al. (2007). Expression of bisecting type and Lewis/Lewis terminated N-glycans on human sperm. *J. Biol. Chem.* 282, 36593–36602. doi:10.1074/jbc.M705134200
- Papi, M., Brunelli, R., Sylla, L., Parasassi, T., Monaci, M., Maulucci, G., et al. (2010). Mechanical properties of zona pellucida hardening. *Eur. Biophys. J.* 39, 987–992. doi:10.1007/s00249-009-0468-3
- Peláez, J., Bongalhardo, D. C., and Long, J. A. (2011). Characterizing the glycocalyx of poultry spermatozoa: III. Semen cryopreservation methods alter the carbohydrate component of rooster sperm membrane glycoconjugates. *Poult. Sci.* 90, 435–443. doi:10.3382/ps.2010-00998
- Pérez, L. M. (2020). “Fish sperm maturation, capacitation, and motility activation.” Chapter 4 in *Reproduction in aquatic animals*. Editors M. Yoshida and J. Asturiano (Singapore: Springer). doi:10.1007/978-981-15-2290-1_5
- Peri, S., Kulkarni, A., Feyertag, F., Berninson, P. M., and Alvarez-Ponce, D. (2018). Phylogenetic distribution of CMP-neu5Ac hydroxylase (CMAH), the enzyme synthesizing the proinflammatory human xenoantigen Neu5Gc. *Genome Biol. Evol.* 10, 207–219. doi:10.1093/gbe/evx251
- Petit, D., Mir, A. M., Petit, J. M., Thisse, C., Delannoy, P., Oriol, R., et al. (2010). Molecular phylogeny and functional genomics of beta-galactoside alpha2, 6-sialyltransferases that explain ubiquitous expression of st6gal1 gene in amniotes. *J. Biol. Chem.* 285, 38399–38414. doi:10.1074/jbc.M110.163931
- Petit, D., Teppa, E., Cenci, U., Ball, S., and Harduin-Lepers, A. (2018). Reconstruction of the sialylation pathway in the ancestor of eukaryotes. *Sci. Rep.* 8, 2946. doi:10.1038/s41598-018-20920-1
- Petit, D., Teppa, E., Mir, A. M., Vicogne, D., Thisse, C., Thisse, B., et al. (2015). Integrative view of α 2, 3-sialyltransferases (ST3Gal) molecular and functional evolution in deuterostomes: Significance of lineage-specific losses. *Mol. Biol. Evol.* 32, 906–927. doi:10.1093/molbev/msu395
- Plancke, Y., Wieruszkeski, J. -M., Alonso, C., Boilly, B., and Strecker, G. (1995). Structure of four acidic oligosaccharides from the jelly coat surrounding the eggs of *Xenopus laevis*. *Eur. J. Biochem.* 231, 434–439. doi:10.1111/j.1432-1033.1995.tb20716.x
- Prasad, S. V., Skinner, S. M., Carino, C., Wang, N., Cartwright, J., and Dunbar, B. S. (2000). Structure and function of the proteins of the mammalian Zona pellucida. *Cells, tissues, organs* 166, 148–164. doi:10.1159/000016730
- Qi, H., Moran, M. M., Navarro, B., Chong, J. A., Krapivinsky, G., Krapivinsky, L., et al. (2007). All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *Proc. Natl. Acad. Sci. U. S. A.* 104, 1219–1223. doi:10.1073/pnas.0610286104
- Raj, I., Sadat Al Hosseini, H., Dioguardi, E., Nishimura, K., Han, L., Villa, A., et al. (2017). Structural basis of egg coat-sperm recognition at fertilization. *Cell* 169 (7), 1315–1326. doi:10.1016/j.cell.2017.05.033
- Rankin, T. L., Coleman, J. S., Epifano, O., Hoodbhoy, T., Turner, S. G., Castle, P. E., et al. (2003). Fertility and taxon-specific sperm binding persist after replacement of mouse sperm receptors with human homologs. *Dev. Cell* 5, 33–43. doi:10.1016/s1534-5807(03)00195-3
- Rath, D., Töpfer-Petersen, E., Michelmann, H. W., Schwartz, P., and Ebeling, S. (2005). Zona pellucida characteristics and sperm-binding patterns of *in vivo* and *in vitro* produced porcine oocytes inseminated with differently prepared spermatozoa. *Theriogenology* 63, 352–362. doi:10.1016/j.theriogenology.2004.09.044
- Roller, R. J., and Wassarman, P. M. (1983). Role of asparagine-linked oligosaccharides in secretion of glycoproteins of the mouse egg's extracellular coat. *J. Biol. Chem.* 258, 13243–13249. doi:10.1016/s0021-9258(17)44107-x
- Rosenstock, P., and Kaufmann, T. (2021). Sialic acids and their influence on human NK cell function. *Cells* 10, 263. doi:10.3390/cells10020263
- Sabeur, K., and Ball, B. A. (2007). Characterization of galactose-binding proteins in equine testis and spermatozoa. *Anim. Reprod. Sci.* 101, 74–84. doi:10.1016/j.anireprosci.2006.08.028
- Sapoń, K., Janas, T., Sikorski, A. F., and Janas, T. (2019). Polysialic acid chains exhibit enhanced affinity for ordered regions of membranes. *Biochim. Biophys. Acta. Biomembr.* 1861, 245–255. doi:10.1016/j.bbamem.2018.07.008
- Sarasquete, C., Cárdenas, S., González de Canales, M. L., and Pascual, E. (2002). Oogenesis in the bluefin tuna, *Thunnus thynnus* L.: A histological and histochemical study. *Histol. Histopathol.* 17, 775–788. doi:10.14670/HH-17.775
- Sasanami, T., Yoshizaki, N., Dohra, H., and Kubo, H. (2011). Sperm acrosin is responsible for the sperm binding to the egg envelope during fertilization in Japanese quail (*Coturnix japonica*). *Reproduction* 142, 267–276. doi:10.1530/REP-11-0120
- Sato, C. (2013). In *Polysialic acid in sialobiology: Structure, biosynthesis and function*. Editors J. Tiralongo and I. Martinez-Duncker (United Arab Emirates, United States, Netherlands: Bentham Books), 33–75.
- Sato, C., and Kitajima, K. (2013). Disialic, oligosialic and polysialic acids: Distribution, functions and related disease. *J. Biochem.* 154, 115–136. doi:10.1093/jb/mvt057
- Sato, C., and Kitajima, K. (2021). Polysialylation and disease. *Mol. Asp. Med.* 79, 100892. doi:10.1016/j.mam.2020.100892
- Sato, C., Kitajima, K., Tazawa, I., Inoue, Y., Inoue, S., and Troy, F. A. (1993). Structural diversity in the α 2 \rightarrow 8-linked polysialic acid chains in salmonid fish egg glycoproteins. Occurrence of poly(Neu5Ac), poly(Neu5Gc), poly(Neu5Ac, Neu5Gc), poly(KDN), and their partially acetylated forms. *J. Biol. Chem.* 268, 23675–23684. doi:10.1016/s0021-9258(19)49515-x
- Sawyer, L. (2021). β -Lactoglobulin and glycodefin: Two sides of the same coin? *Front. Physiol.* 12, 678080. doi:10.3389/fphys.2021.678080
- Schaper, W., Bentrop, J., Ustinova, J., Blume, L., Kats, E., Tiralongo, J., et al. (2012). Identification and biochemical characterization of two functional CMP-sialic acid synthetases in *Danio rerio*. *J. Biol. Chem.* 287, 13239–13248. doi:10.1074/jbc.M111.327544
- Schauer, R. (2012). In *Sialic acids: Chemistry, metabolism, and function*. Editor R. Schauer (Wien, New York: Springer-Verlag).
- Schauer, R. (1982). Chemistry, metabolism, and biological functions of sialic acid. *Adv. Carbohydr. Chem. Biochem.* 40, 131–234. doi:10.1016/s0065-2318(08)60109-2
- Schauer, R., and Kamerling, J. P. (2011). The chemistry and biology of trypanosomal trans-sialidases: Virulence factors in chagas disease and sleeping sickness. *ChemBioChem* 12, 2246–2264. doi:10.1002/cbic.201100421
- Schauer, R. (2009). Sialic acids as regulators of molecular and cellular interactions. *Curr. Opin. Struct. Biol.* 19, 507–514. doi:10.1016/j.sbi.2009.06.003
- Schauer, R. (2004). Sialic acids: Fascinating sugars in higher animals and man. *Zoology* 107, 49–64. doi:10.1016/j.zool.2003.10.002
- Schauer, R., Srinivasan, G. V., Coddeville, B., Zanetta, J. P., and Guerardel, Y. (2009). Low incidence of N-glycolylneuraminic acid in birds and reptiles and its absence in the platypus. *Carbohydr. Res.* 344, 1494–1500. doi:10.1016/j.carres.2009.05.020
- Schjoldager, K. T., Narimatsu, Y., Joshi, H. J., and Clausen, H. (2020). Global view of human protein glycosylation pathways and functions. *Nat. Rev. Mol. Cell Biol.* 21, 729–749. doi:10.1038/s41580-020-00294-x
- Schleuning, W.-D., Hell, R., Schiessler, H., and Fritz, H. (1975). Boar acrosin, I. Modified isolation procedure, active-site titration and evidence for the presence of sialic acid. *Hoppe. Seylers. Z. Physiol. Chem.* 356, 1915–1921. doi:10.1515/bchm2.1975.356.2.1915
- Schnaar, R. L., Gerardy-Schahn, R., and Hildebrandt, H. (2014). Sialic acids in the brain: Gangliosides and polysialic acid in nervous system development, stability, disease, and regeneration. *Physiol. Rev.* 94, 461–518. doi:10.1152/physrev.00033.2013
- Scott, M. A. (2000). A glimpse at sperm function *in vivo*: Sperm transport and epithelial interaction in the female reproductive tract. *Anim. Reprod. Sci.* 61, 337–348. doi:10.1016/S0378-4320(00)00130-5

- Seko, A., Kitajima, K., Iwasaki, M., Inoue, S., and Inoue, Y. (1989). Structural studies of fertilization-associated carbohydrate-rich glycoproteins (hyosoporphin) isolated from the fertilized and unfertilized eggs of flounder, *Paralichthys olivaceus*. *J. Biol. Chem.* 264, 15922–15929. doi:10.1016/S0021-9258(18)71567-6
- Senthilkumaran, B., and Joy, K. P. (1993). Annual cyclic, and castration and cyproterone acetate-induced, changes in sialic acid content of the seminal vesicle of the catfish, *Heteropneustes fossilis* (Bloch). *Fish. Physiol. Biochem.* 10, 425–430. doi:10.1007/BF00004509
- Seppälä, M., Koistinen, H., Koistinen, R., Chiu, P. C. N., and Yeung, W. S. B. (2007). Glycosylation related actions of glycodefin: Gamete, cumulus cell, immune cell and clinical associations. *Hum. Reprod. Update* 13, 275–287. doi:10.1093/humupd/dmm004
- Sever, D. M., Moriarty, E. C., Rania, L. C., and Hamlett, W. C. (2001). Sperm storage in the oviduct of the internal fertilizing frog *Ascaphus truei*. *J. Morphol.* 248, 1–21. doi:10.1002/jmor.1017
- Shaw, L., and Schauer, R. (1988). The biosynthesis of N-glycolylneuraminic acid occurs by hydroxylation of the CMP-glycoside of NAcetylneuraminic acid. *Biol. Chem. Hoppe. Seyler.* 369, 477–486. doi:10.1151/bchm3.1988.369.1.477
- Shi, S., Williams, S. A., Seppo, A., Kurniawan, H., Chen, W., Ye, Z., et al. (2004). Inactivation of the Mgat1 gene in oocytes impairs oogenesis, but embryos lacking complex and hybrid N-glycans develop and implant. *Mol. Cell. Biol.* 24, 9920–9929. doi:10.1128/MCB.24.22.9920-9929.2004
- Shimoda, Y., Kitajima, K., Inoue, S., and Inoue, Y. (1994). Calcium ion binding of three different types of oligo/polysialic acids as studied by equilibrium dialysis and circular dichroic methods. *Biochemistry* 33, 1202–1208. doi:10.1021/bi00171a020
- Simakov, O., Kawashima, T., Marlétaz, F., Jenkins, J., Koyanagi, R., Mitros, T., et al. (2015). Hemichordate genomes and deuterostome origins. *Nature* 527, 459–465. doi:10.1038/nature16150
- Simon, P., Bäumner, S., Busch, O., Röhrich, R., Kaese, M., Richterich, P., et al. (2013). Polysialic acid is present in mammalian semen as a post-translational modification of the neural cell adhesion molecule NCAM and the polysialyltransferase ST8SiaII. *J. Biol. Chem.* 288, 18825–18833. doi:10.1074/jbc.M113.451112
- Simons, K., and Ikonen, E. (1997). Functional rafts in cell membranes. *Nature* 387, 569–572. doi:10.1038/42408
- Singh, M. S., and Joy, K. P. (1999). Annual correlative changes in some biochemical contents of seminal vesicle and testis in the catfish *Clarias batrachus* (L.). *Zool. Sci.* 16, 345–356. doi:10.2108/zsj.16.345
- Sinowatz, F., Töpfer-Petersen, E., Kölle, S., and Palma, G. (2001). Functional morphology of the zona pellucida. *Anat. Histol. Embryol.* 30, 257–263. doi:10.1046/j.1439-0264.2001.00337.x
- Siu, K. K., Serrão, V., Ziyat, A., and Lee, J. E. (2021). The cell biology of fertilization: Gamete attachment and fusion. *J. Cell Biol.* 220, e202102146. doi:10.1083/jcb.202102146
- Srivastava, P. N., Kumar, V. M., and Arbtan, K. D. (1988). Neuraminidase induces capacitation and acrosome reaction in mammalian spermatozoa. *J. Exp. Zool.* 245, 106–110. doi:10.1002/jez.1402450116
- Steele, M. G., and Wishart, G. J. (1996). Demonstration that the removal of sialic acid from the surface of chicken spermatozoa impedes their transvaginal migration. *Theriogenology* 46, 1037–1044. doi:10.1016/s0093-691x(96)00268-3
- Stern, C. A., Braverman, T. R., and Tiemeyer, M. (2000). Molecular identification, tissue distribution and subcellular localization of mST3GalV/GM3 synthase. *Glycobiology* 10, 365–374. doi:10.1093/glycob/10.4.365
- Strecker, G., Wieruszkes, J. M., Plancke, Y., and Boilly, B. (1995). Primary structure of 12 neutral oligosaccharide-alditols released from the jelly coats of the anuran *Xenopus laevis* by reductive β -elimination. *Glycobiology* 5, 137–146. doi:10.1093/glycob/5.1.137
- Suarez, S. S. (2016). Mammalian sperm interactions with the female reproductive tract. *Cell Tissue Res.* 363, 185–194. doi:10.1007/s00441-015-2244-2
- Summers, R. G., and Hylander, B. L. (1975). Species-specificity of acrosome reaction and primary gamete binding in echinoids. *Exp. Cell Res.* 96, 63–68. doi:10.1016/s0014-4827(75)80037-1
- Sun, Q. Y. (2003). Cellular and molecular mechanisms leading to cortical reaction and polyspermy block in mammalian eggs. *Microsc. Res. Tech.* 61, 342–348. doi:10.1002/jemt.10347
- Suzuki, K., Tabe, N., Kojima, S., Hamano, A., Orita, M., and Yonezawa, N. (2015). The hinge region of bovine zona pellucida glycoprotein ZP3 is involved in the formation of the sperm-binding active ZP3/ZP4 complex. *Biomolecules* 5, 3339–3353. doi:10.3390/biom5043339
- Szczykutowicz, J., Kałuża, A., Kaźmierowska-Niemczuk, M., and Ferens-Sieczkowska, M. (2019). The potential role of seminal plasma in the fertilization outcomes. *Biomed. Res. Int.* 2019, 5397804. doi:10.1155/2019/5397804
- Taguchi, T., Iwasaki, M., Muto, Y., Kitajima, K., Inoue, S., Khoo, K. H., et al. (1996). Occurrence and structural analysis of highly sulfated multiantennary N-linked glycan chains derived from a fertilization-associated carbohydrate-rich glycopeptide in unfertilized eggs of *Tribolodon hakonensis*. *Eur. J. Biochem.* 238, 357–367. doi:10.1111/j.1432-1033.1996.03572.x
- Taguchi, T., Seko, A., Kitajima, K., Inoue, S., Iwamatsu, T., Khoo, K. H., et al. (1993). Structural studies of a novel type of tetraantennary sialoglycan unit in a carbohydrate-rich glycopeptide isolated from the fertilized eggs of Indian Medaka fish, *Oryzias latipes*. *J. Biol. Chem.* 268, 2353–2362. doi:10.1016/s0021-9258(18)53783-2
- Taguchi, T., Seko, A., Kitajima, K., Muto, Y., Inoue, S., Khoo, K. H., et al. (1994). Structural studies of a novel type of pentaantennary large glycan unit in the fertilization-associated carbohydrate-rich glycopeptide isolated from the fertilized eggs of *Oryzias latipes*. *J. Biol. Chem.* 269, 8762–8771. doi:10.1016/s0021-9258(17)37034-5
- Takashima, S., Tsuji, S., and Tsujimoto, M. (2002). Characterization of the second type of human beta-galactoside alpha 2, 6-sialyltransferase (ST6Gal II), which sialylates Galbeta 1, 4GlcNAc structures on oligosaccharides preferentially. Genomic analysis of human sialyltransferase genes. *J. Biol. Chem.* 277, 45719–45728. doi:10.1074/jbc.M206808200
- Takashima, S., Tsuji, S., and Tsujimoto, M. (2003). Comparison of the enzymatic properties of mouse beta-galactoside alpha2, 6-sialyltransferases, ST6Gal I and II. *J. Biochem.* 134, 287–296. doi:10.1093/jb/mvg142
- Taniguchi, A., Hioki, M., and Matsumoto, K. (2003). Transcriptional regulation of human Galbeta1, 3GalNAc/Galbeta1, 4GlcNAc alpha2, 3-sialyltransferase (hST3Gal IV) gene in testis and ovary cell lines. *Biochem. Biophys. Res. Commun.* 301, 764–768. doi:10.1016/s0006-291x(03)00041-x
- Tecle, E., and Gagneux, P. (2015). Sugar-coated sperm: Unraveling the functions of the mammalian sperm glycocalyx. *Mol. Reprod. Dev.* 82, 635–650. doi:10.1002/mrd.22500
- Tecle, E., Reynoso, H. S., Wang, R., and Gagneux, P. (2019). The female reproductive tract contains multiple innate sialic acid-binding immunoglobulin-like lectins (Siglecs) that facilitate sperm survival. *J. Biol. Chem.* 294, 11910–11919. doi:10.1074/jbc.RA119.008729
- Teppa, R. E., Petit, D., Plechakova, O., Cogez, V., and Harduin-Lepers, A. (2016). Phylogenetic-derived insights into the evolution of sialylation in eukaryotes: Comprehensive analysis of vertebrate β -galactoside α 2, 3/6-sialyltransferases (ST3Gal and ST6Gal). *Int. J. Mol. Sci.* 17, 1286. doi:10.3390/ijms17081286
- Tezuka, T., Taguchi, T., Kanamori, A., Muto, Y., Kitajima, K., Inoue, Y., et al. (1994). Identification and structural determination of the KDN-containing N-linked glycan chains consisting of Bi- and triantennary complex-type units of KDN-glycoprotein previously isolated from rainbow trout vitelline envelopes. *Biochemistry* 33, 6495–6502. doi:10.1021/bi00187a016
- Thall, A. D., Malý, P., and Lowe, J. B. (1995). Oocyte Gala1, 3Gal epitopes implicated in sperm adhesion to the zona pellucida glycoprotein ZP3 are not required for fertilization in the mouse. *J. Biol. Chem.* 270, 21437–21440. doi:10.1074/jbc.270.37.21437
- Tholl, N., Naqvi, S., McLaughlin, E., Boyles, S., Bieber, A. L., and Chandler, D. E. (2011). Swimming of *Xenopus laevis* sperm exhibits multiple gears and its duration is extended by egg jelly constituents. *Biol. Bull.* 220, 174–185. doi:10.1086/BBLv220n3p174
- Thompson, L. A., Barratt, C. L. R., Bolton, A. E., and Cooke, I. D. (1992). The leukocytic reaction of the human uterine cervix. *Am. J. Reprod. Immunol.* 28, 85–89. doi:10.1111/j.1600-0897.1992.tb00765.x
- Tokuhiro, K., and Dean, J. (2018). Glycan-independent gamete recognition triggers egg zinc sparks and ZP2 cleavage to prevent polyspermy. *Dev. Cell* 46, 627–640. doi:10.1016/j.devcel.2018.07.020
- Tollner, T. L., Bevins, C. L., and Cherr, G. N. (2012). Multifunctional glycoprotein DEFB126—a curious story of defensin-clad spermatozoa. *Nat. Rev. Urol.* 9, 365–375. doi:10.1038/nrurol.2012.109
- Tollner, T. L., Venners, S. A., Hollox, E. J., Yudin, A. I., Liu, X., Tang, G., et al. (2011). A common mutation in the defensin DEFB126 causes impaired sperm function and subfertility. *Sci. Transl. Med.* 3, 92ra65. doi:10.1126/scitranslmed.3002289
- Tollner, T. L., Yudin, A. I., Treece, C. A., Overstreet, J. W., and Cherr, G. N. (2008). Macaque sperm coating protein DEFB126 facilitates sperm penetration of cervical mucus. *Hum. Reprod.* 23, 2523–2534. doi:10.1093/humrep/den276
- Toowicharanont, P., and Chulavatnatol, M. (1983). Characterization of sialoglycoproteins of rat epididymal fluid and spermatozoa by periodate tritiated borohydride. *J. Reprod. Fertil.* 67, 133–141. doi:10.1530/jrf.0.0670133
- Töpfer-Petersen, E. (1999). Carbohydrate-based interactions on the route of a spermatozoon to fertilization. *Hum. Reprod. Update* 5, 314–329. doi:10.1093/humupd/5.4.314
- Töpfer-Petersen, E., Wagner, A., Friedrich, J., Petrunkina, A., Ekhlesi-Hundrieser, M., Waberski, D., et al. (2002). Function of the mammalian oviductal sperm reservoir. *J. Exp. Zool.* 292, 210–215. doi:10.1002/jez.1157

- Toshimori, K., Araki, S., Öra, C., and Eddy, E. M. (1991). Loss of sperm surface sialic acid induces phagocytosis: An assay with a monoclonal antibody T21, which recognizes a 54K sialoglycoprotein. *Arch. Androl.* 27, 79–86. doi:10.3109/01485019108987656
- Tosti, E. (1994). Sperm activation in species with external fertilisation. *Zygote* 2, 359–361. doi:10.1017/s0967199400002215
- Troy, F. A. (1992). Polysialylation: From bacteria to brains. *Glycobiology* 2, 5–23. doi:10.1093/glycob/2.1.5
- Tumova, L., Zigo, M., Sutovsky, P., Sedmikova, M., and Postlerova, P. (2021). Ligands and receptors involved in the sperm-zona pellucida interactions in Mammals. *Cells* 10, 133–136. doi:10.3390/cells10010133
- Valbuena, G., Alonso, E., de Ubago, M. M., Madrid, J. F., Díaz-Flores, L., and Sáez, F. J. (2012). Histochemical identification of sialylated glycans in *Xenopus laevis* testis. *J. Anat.* 221, 318–330. doi:10.1111/j.1469-7580.2012.01548.x
- Valle, G. M., Cinelli, L. P., Todeschini, A. R., de Brito-Gitirana, L., Vilela-Silva, A. C. E. S., and Mourão, P. A. S. (2015). Sperm and egg jelly coat from sea urchin *Lytechinus variegatus* collected in Rio de Janeiro contain distinct sialic acid-rich polysaccharides. *Braz. Arch. Biol. Technol.* 58, 617–627. doi:10.1590/S1516-8913201500052
- Van Blerkom, J., and Caltrider, K. (2013). Sperm attachment and penetration competence in the human oocyte: A possible aetiology of fertilization failure involving the organization of oolemmal lipid raft microdomains influenced by the $\Delta\Psi_m$ of subplasmalemmal mitochondria. *Reprod. Biomed. Online* 27, 690–701. doi:10.1016/j.rbmo.2013.09.011
- Varki, A., and Angata, T. (2006). Siglecs—the major subfamily of I-type lectins. *Glycobiology* 16, 1R–27R. doi:10.1093/glycob/cwj008
- Varki, A. (2017). Biological roles of glycans. *Glycobiology* 27, 3–49. doi:10.1093/glycob/cww086
- Varki, A. (1992). Diversity in the sialic acids. *Glycobiology* 2, 25–40. doi:10.1093/glycob/2.1.25
- Varki, A. (2006). Nothing in glycobiology makes sense, except in the light of evolution. *Cell* 126, 841–845. doi:10.1016/j.cell.2006.08.022
- Velásquez, J. G., Canovas, S., Barajas, P., Marcos, J., Jiménez-Movilla, M., Gallego, R. G., et al. (2007). Role of sialic acid in bovine sperm-zona pellucida binding. *Mol. Reprod. Dev.* 5, 617–628. doi:10.1002/mrd.20619
- Venuto, M. T., Declouement, M., Martorell Ribera, J., Noel, M., Rebl, A., Cogeze, V., et al. (2020a). Vertebrate Alpha2, 8-sialyltransferases (ST8Sia): A teleost perspective. *Int. J. Mol. Sci.* 21, 513. doi:10.3390/ijms21020513
- Venuto, M. T., Martorell-Ribera, J., Bochert, R., Harduin-Lepers, A., Rebl, A., and Galuska, S. P. (2020b). Characterization of the polysialylation status in ovaries of the salmonid fish *Coregonus maraena* and the percid fish *Sander lucioperca*. *Cells* 9, doi:10.3390/cells9112391
- Villaverde, A. I. S. B., Ogle, R. A., Lewis, P., Carbone, V., Velkov, T., Netherton, J. K., et al. (2020). Sialylation of asparagine 612 inhibits aconitase activity during mouse sperm capacitation; a possible mechanism for the switch from oxidative phosphorylation to glycolysis. *Mol. Cell. Proteomics* 19, 1860–1875. doi:10.1074/mcp.RA120.002109
- Vo, L. H., Yen, T. Y., Macher, B. A., and Hedrick, J. L. (2003). Identification of the ZPC oligosaccharide ligand involved in sperm binding and the glycan structures of *Xenopus laevis* vitelline envelope glycoproteins. *Biol. Reprod.* 69, 1822–1830. doi:10.1095/biolreprod.103.015370
- Wang, Y., and Neumann, H. (2010). Alleviation of neurotoxicity by microglial human Siglec-11. *J. Neurosci.* 30, 3482–3488. doi:10.1523/JNEUROSCI.3940-09.2010
- Wang, Y., Zhao, W., Mei, S., Chen, P., Leung, T. Y., Lee, C. L., et al. (2021). Identification of sialyl-lewis(x)-interacting protein on human spermatozoa. *Front. Cell Dev. Biol.* 9, 700396. doi:10.3389/fcell.2021.700396
- Wassarman, P. M. (2011). Development: The sperm's sweet tooth. *Science* 333, 1708–1709. doi:10.1126/science.1212841
- Wassarman, P. M., and Litscher, E. S. (2022). Female fertility and the zona pellucida. *eLife* 11, e76106. doi:10.7554/eLife.76106
- Wassarman, P. M., and Litscher, E. S. (2021). Zona pellucida genes and proteins: Essential players in mammalian oogenesis and fertility. *Genes* 12, 1266. doi:10.3390/genes12081266
- Wassarman, P. M., and Mortillo, S. (1991). Structure of the mouse egg extracellular coat, the zona pellucida. *Int. Rev. Cytol.* 130, 85–110. doi:10.1016/S0074-7696(08)61502-8
- Watanabe, A., and Onitake, K. (2002). The urodele egg-coat as the apparatus adapted for the internal fertilization. *Zool. Sci.* 19, 1341–1347. doi:10.2108/zsj.19.1341
- Werneburg, S., Buettner, F. F. R., Erben, L., Mathews, M., Neumann, H., Mühlhoff, M., et al. (2016). Polysialylation and lipopolysaccharide-induced shedding of E-selectin ligand-1 and neuropilin-2 by microglia and THP-1 macrophages. *Glia* 64, 1314–1330. doi:10.1002/glia.23004
- Williams, S. A., Xia, L., Cummings, R. D., McEver, R. P., and Stanley, P. (2007). Fertilization in mouse does not require terminal galactose or N-acetylglucosamine on the zona pellucida glycans. *J. Cell Sci.* 120, 1341–1349. doi:10.1242/jcs.004291
- Wira, C. R., Patel, M. V., Ghosh, M., Mukura, L., and Fahey, J. V. (2011). Innate immunity in the human female reproductive tract: Endocrine regulation of endogenous antimicrobial protection against HIV and other sexually transmitted infections. *Am. J. Reprod. Immunol.* 65, 196–211. doi:10.1111/j.1600-0897.2011.00970.x
- Wolf, D. P., Nishihara, T., West, D. M., Wyrick, R. E., and Hedrick, J. L. (1976). Isolation, physicochemical properties, and the macromolecular composition of the vitelline and fertilization envelopes from *Xenopus laevis* eggs. *Biochemistry* 15, 3671–3678. doi:10.1021/bi00662a005
- Yabe, U., Sato, C., Matsuda, T., and Kitajima, K. (2003). Polysialic acid in human milk: CD36 is a new member of mammalian polysialic acid-containing glycoprotein. *J. Biol. Chem.* 278, 13875–13880. doi:10.1074/jbc.M300458200
- Yamakawa, N., Sato, C., Miyata, S., Maehashi, E., Toriyama, M., Sato, N., et al. (2007). Development of sensitive chemical and immunochemical methods for detecting sulfated sialic acids and their application to glycoconjugates from sea urchin sperm and eggs. *Biochimie* 89, 1396–1408. doi:10.1016/j.biochi.2007.04.010
- Yamakawa, N., Vanbeselaere, J., Chang, L. Y., Yu, S. Y., Ducrocq, L., Harduin-Lepers, A., et al. (2018). Systems glycomics of adult zebrafish identifies organ-specific sialylation and glycosylation patterns. *Nat. Commun.* 9, 4647. doi:10.1038/s41467-018-06950-3
- Yanagimachi, R., Harumi, T., Matsubara, H., Yan, W., Yuan, S., Hirohashi, N., et al. (2017). Chemical and physical guidance of fish spermatozoa into the egg through the micropyle. *Biol. Reprod.* 96, 780–799. doi:10.1093/biolre/iox015
- Yanagimachi, R. (2011). Mammalian sperm acrosome reaction: Where does it begin before fertilization. *Biol. Reprod.* 85, 4–5. doi:10.1095/biolreprod.111.092601
- Yang, P., Major, D., and Rutishauser, U. (1994). Role of charge and hydration in effects of polysialic acid on molecular interactions on and between cell membranes. *J. Biol. Chem.* 269, 23039–23044. doi:10.1016/s0021-9258(17)31616-2
- Yeşilyurt, B., Şahar, U., and Devci, R. (2015). Determination of the type and quantity of sialic acid in the egg jelly coat of the sea urchin *Paracentrotus lividus* using capillary LC-ESI-MS/MS. *Mol. Reprod. Dev.* 82, 115–122. doi:10.1002/mrd.22448
- Yonezawa, N., Fukui, N., Kuno, M., Shinoda, M., Goko, S., Mitsui, S., et al. (2001). Molecular cloning of bovine zona pellucida glycoproteins ZPA and ZPB and analysis for sperm-binding component of the zona. *Eur. J. Biochem.* 268, 3587–3594. doi:10.1046/j.1432-1327.2001.02269.x
- Yonezawa, N., Mitsui, S., Kudo, K., and Nakano, M. (1997). Identification of an N-glycosylated region of pig zona pellucida glycoprotein ZPB that is involved in sperm binding. *Eur. J. Biochem.* 248, 86–92. doi:10.1111/j.1432-1033.1997.00086.x
- Yoshida, Y., Kojima, N., and Tsuji, S. (1995). Molecular cloning and characterization of a third type of N-glycan alpha 2, 8-sialyltransferase from mouse lung. *J. Biochem.* 118, 658–664. doi:10.1093/oxfordjournals.jbchem.a124960
- Yudin, A. I., Generao, S. E., Tollner, T. L., Treece, C. A., Overstreet, J. W., and Cherr, G. N. (2005). Beta-defensin 126 on the cell surface protects sperm from immunorecognition and binding of anti-sperm antibodies. *Biol. Reprod.* 73, 1243–1252. doi:10.1095/biolreprod.105.042432
- Yurewicz, E. C., Pack, B. A., and Sacco, A. G. (1991). Isolation, composition, and biological activity of sugar chains of porcine oocyte zona pellucida 55K glycoproteins. *Mol. Reprod. Dev.* 30, 126–134. doi:10.1002/mrd.1080300209
- Zhou, C. X., Zhang, Y. L., Xiao, L., Zheng, M., Leung, K. M., Chan, M. Y., et al. (2004). An epididymis-specific β -defensin is important for the initiation of sperm maturation. *Nat. Cell Biol.* 6, 458–464. doi:10.1038/ncb1127
- Zlatina, K., Lütteke, T., and Galuska, S. P. (2017). Individual impact of distinct polysialic acid chain lengths on the cytotoxicity of histone H1, H2A, H2B, H3 and H4. *Polym. (Basel)* 12, E720. doi:10.3390/polym9120720