



HAL
open science

Natural Organic Matter Membrane Fractionation: A new Approach

Karlien Dejaeger

► **To cite this version:**

Karlien Dejaeger. Natural Organic Matter Membrane Fractionation: A new Approach. Euromembrane Conference 2022, Nov 2022, Sorrento, Italy. hal-03920040

HAL Id: hal-03920040

<https://hal.univ-lille.fr/hal-03920040>

Submitted on 16 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Natural organic matter membrane fractionation: a new approach

Dejaeger Karlien

24th November 2022

Marjolein Vanoppen, Gabriel Billon, Justine Criquet, Cécile Vignal, Emile R. Cornelissen

Why membrane fractionation?


1. To split the **complex organic matter** mixture into fractions with **similar physical properties**
2. To **correlate** these properties to certain **water-related issues** such as disinfection by-product formation

Regulated ($\mu\text{g/L}$)

Trihalomethanes (THM)	Haloacetic acids (HAA)
$\begin{array}{c} \text{H} \\ \\ \text{X}-\text{C}-\text{X} \\ \\ \text{X} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{X}-\text{C}-\text{C}-\text{OH} \\ \\ \text{X} \end{array}$

Unregulated (ng/L)



Haloacetonitriles (HAN)	Haloacetamides (HAcAm)
$\begin{array}{c} \text{X} \\ \\ \text{X}-\text{C}-\text{C}\equiv\text{N} \\ \\ \text{X} \end{array}$	$\begin{array}{c} \text{O} \\ \\ \text{X}-\text{C}-\text{C}-\text{N}-\text{H} \\ \\ \text{X} \\ \\ \text{H} \end{array}$





Current approach

- Fractionation based on the MWCO of a series of ultrafiltration membranes






Identification of disinfection by-product precursors by natural organic matter fractionation: a review

Karlien Dejaeger^{1,2,3} · Justine Criquet² · Marjolein Vanoppen^{1,4} · Cécile Vignal³ · Gabriel Billon² · Emile R. Cornelissen^{1,4,5}



The new approach

- Fractionation based on HPSEC-TOC





STEP 1: Membrane selection


- Ultrafiltration membrane screening



Separation efficiency factor:

$$\alpha = 1 - \frac{1 - Retention_{BP}}{1 - Retention_{HS}}$$

	1 BAR	2 BAR
MT MEMBRANE	0.72	0.85
XT MEMBRANE	0.78	0.91



- Nanofiltration membrane screening



Separation efficiency factor:







$$\alpha = 1 - \frac{1 - Retention_{HS}}{1 - Retention_{LMW}}$$

	3 BAR	5 BAR
NFW MEMBRANE	0.93	0.92
NFX MEMBRANE	0.90	0.91



STEP 2: Fractionation of surface water

Start composition:







STEP 3: Massbalance model (black box) to predict fractionation

- Input variables:
 - Flux
 - Start concentration
 - Retention
- Validated during nanofiltration with a synthetic ionmatrix

Concentration





Diafiltration





Conclusions and outlook

- Fractionation based on HPSEC-TOC yields better separated fractions compared to the fractionation based on MWCO



- Improvements needed in terms of recovery

Identification of disinfection by-product precursors by natural organic matter fractionation: a review

Karlien Dejaeger^{1,2,3} · Justine Criquet² · Marjolein Vanoppen^{1,4} · Cécile Vignat³ · Gabriel Billon² · Emile R. Cornelissen^{1,4,5}





karlien.dejaeger@ugent.be
karlien.dejaeger@univ-lille.fr




[https://www.linkedin.com/in/
karlien-dejaeger/](https://www.linkedin.com/in/karlien-dejaeger/)






Supporting info: MWCO determination

$$R(MW^*) = \int_0^{MW^*} \frac{1}{s_{MW} * \sqrt{2\pi}} * \frac{1}{MW} * e^{-\frac{(\ln(MW) - \ln(MWCO) + 0.56 * s_{MW})^2}{2 * s_{MW}^2}} dMW$$



NFX

150-300 Da membrane → MWCO: 130 g/mol



NFW

300-500 Da membrane → MWCO: 139 g/mol



Supporting info: Flux during fractionation

