



**HAL**  
open science

## Openprot 2021: deeper functional annotation of the coding potential of eukaryotic genomes

Marie A. Brunet, Jean-Francois Lucier, Maxime Levesque, Sebastien Leblanc, Jean-Francois Jacques, Hassan R. H. Al-Saedi, Mariano Avino, Isabelle Fournier, Michel Salzet, Aida Ouangraoua, et al.

### ► To cite this version:

Marie A. Brunet, Jean-Francois Lucier, Maxime Levesque, Sebastien Leblanc, Jean-Francois Jacques, et al.. Openprot 2021: deeper functional annotation of the coding potential of eukaryotic genomes. Nucleic Acids Research, 2020, Nucleic Acids Research, 49 (D1), pp.D380-D388. 10.1093/nar/gkaa1036 . hal-04028636

**HAL Id: hal-04028636**

<https://hal.univ-lille.fr/hal-04028636v1>

Submitted on 14 Mar 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# OpenProt 2021: deeper functional annotation of the coding potential of eukaryotic genomes

Marie A. Brunet<sup>1,2,\*</sup>, Jean-François Lucier<sup>3,4</sup>, Maxime Levesque<sup>3,4</sup>, Sébastien Leblanc<sup>1,2</sup>, Jean-François Jacques<sup>1,2</sup>, Hassan R.H. Al-Saedi<sup>1</sup>, Noé Guillois<sup>1,2</sup>, Frederic Grenier<sup>3,4</sup>, Mariano Avino<sup>1</sup>, Isabelle Fournier<sup>5</sup>, Michel Salzet<sup>5</sup>, Aïda Ouangraoua<sup>6</sup>, Michelle S. Scott<sup>1</sup>, François-Michel Boisvert<sup>7</sup> and Xavier Roucou<sup>1,2,\*</sup>

<sup>1</sup>Department of Biochemistry and Functional Genomics, Université de Sherbrooke, 3201 Jean Mignault, Sherbrooke, QC J1E 4K8, Canada, <sup>2</sup>PROTEO, Quebec Network for Research on Protein Function, Structure, and Engineering, Université Laval, Quebec City, QC G1V0A6, Canada, <sup>3</sup>Center for Computational Science, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada, <sup>4</sup>Biology Department, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada, <sup>5</sup>INSERM U1192, Laboratoire Protéomique, Réponse Inflammatoire & Spectrométrie de Masse (PRISM), Université de Lille, F-59000 Lille, France, <sup>6</sup>Informatics Department, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada and <sup>7</sup>Department of Immunology and Cellular Biology, Université de Sherbrooke, Sherbrooke, QC J1E 4K8, Canada

Received September 14, 2020; Revised October 15, 2020; Editorial Decision October 16, 2020; Accepted October 16, 2020

## ABSTRACT

OpenProt ([www.openprot.org](http://www.openprot.org)) is the first proteogenomic resource supporting a polycistronic annotation model for eukaryotic genomes. It provides a deeper annotation of open reading frames (ORFs) while mining experimental data for supporting evidence using cutting-edge algorithms. This update presents the major improvements since the initial release of OpenProt. All species support recent NCBI RefSeq and Ensembl annotations, with changes in annotations being reported in OpenProt. Using the 131 ribosome profiling datasets re-analysed by OpenProt to date, non-AUG initiation starts are reported alongside a confidence score of the initiating codon. From the 177 mass spectrometry datasets re-analysed by OpenProt to date, the unicity of the detected peptides is controlled at each implementation. Furthermore, to guide the users, detectability statistics and protein relationships (isoforms) are now reported for each protein. Finally, to foster access to deeper ORF annotation independently of one's bioinformatics skills or computational resources, OpenProt now offers a data analysis platform. Users can submit their dataset for analysis and receive the results from the analysis by OpenProt. All data on OpenProt are freely available and downloadable for each species, the release-based format ensuring a

continuous access to the data. Thus, OpenProt enables a more comprehensive annotation of eukaryotic genomes and fosters functional proteomic discoveries.

## INTRODUCTION

Genome annotations are the cornerstones of all research endeavours as they define the proteomic landscape. In the face of such a crucial role, they enforce data-driven and arbitrary criteria to reduce spurious annotations to a minimum (1–3). For example, annotations are limited to open reading frames (ORFs) longer than 100 codons, a single coding sequence per transcript and an ATG start codon, excepted for previously characterized examples. These criteria substantially shape and limit the exploration of the proteome (1,4–6) and an ever-increasing number of studies report the need for deeper ORF annotation to better explore and understand cellular mechanisms (1,7–9). With the development of ribosome profiling (10), a technique to detect translation events throughout the genome, inaccuracies from current annotations have been experimentally proven (8,11–15). Building on ribosome profiling data, several small ORF repositories have been published to foster the functional characterization of these previously overlooked ORFs (16–18). These repositories have greatly participated in the incorporation of small ORFs in genome annotations (1,16). Yet, because they rely sometimes exclusively on ribosome profiling data analyses, these repositories suffer from the technical biases inherent to the experiment. This may hin-

\*To whom correspondence should be addressed. Tel: +1 819 821 8000 (Ext 72240); Email: [xavier.roucou@usherbrooke.ca](mailto:xavier.roucou@usherbrooke.ca)  
Correspondence may also be addressed to Marie A. Brunet. Email: [marie.brunet@usherbrooke.ca](mailto:marie.brunet@usherbrooke.ca)

der the detection of overlapping ORFs or ORFs in low-abundance transcripts or in repetitive regions (1,10,19,20). Furthermore, the accuracy of the detection of isoforms may vary depending on the algorithm used to mine the data (21–23).


Using a different approach, OpenProt (24) first predicts all possible ORFs within the transcriptome retrieved from two annotations [NCBI RefSeq (25) and Ensembl (26)] and then gathers supporting evidence. The prediction pipeline does not enforce a maximal length threshold, although it does filter for a minimal length of 30 codons and an AUG initiating codon. The predicted encoded proteins are then categorized as follows: RefProt or reference proteins are known proteins annotated in NCBI RefSeq (25), Ensembl (26) and/or UniProt (27); novel isoforms are unannotated proteins with a significant sequence identity to a RefProt from the same gene; and AltProts are unannotated proteins with no significant identity to a RefProt from the same gene. Finally, to assert confidence of the predicted proteins, OpenProt retrieves evidence for each annotated protein. These are *in silico* evidence (conservation and prediction of functional domains) and experimental evidence (translation evidence from ribosome profiling data and expression evidence from mass spectrometry (MS) data). OpenProt thus offers a deep annotation of the genome of 10 species, identifying novel isoforms and novel proteins in an unbiased yet data-driven manner.

The initial release of OpenProt provided a much needed resource for functional proteomic discoveries (7,28–32) and was the first proteogenomic resource supporting a polycistronic model of annotation for eukaryotic genomes (24). Here, we present an update of the OpenProt database incorporating the re-analysis of an additional 63 MS datasets and 44 ribosome profiling datasets, for a total of 177 and 131 datasets, respectively. This increase in the number of analysed datasets demonstrates the constant growth of the OpenProt resource. However, it does not constitute the core of this update. In this release, OpenProt implemented the transition to the latest annotations from NCBI RefSeq and Ensembl, the annotation of non-AUG initiation sites from ribosome profiling datasets, the incorporation of a time-independent quality control of the unicity of detected peptides and the annotation of protein identities within a gene (isoform prediction). Additionally, OpenProt website was upgraded to a more user-friendly interface and it now provides a platform for users to submit their datasets for re-analysis with the OpenProt pipeline. This platform aims to grant access to a deeper ORF annotation for data analysis to the wider community, independently of bioinformatics skills or computational resources. With a user-friendly and interactive interface, OpenProt ([www.openprot.org](http://www.openprot.org)) aims to foster discoveries of functional yet currently non-annotated proteins (Figure 1; Supplementary Material S1).

## DATABASE CONTENT UPDATES

### Genome annotations for ORF prediction

Genome annotations are constantly evolving, inserting novel transcripts, discarding others and updating annotations of coding sequence. Although most changes are minor from one release to the next, they do accumulate over time



**Figure 1.** The OpenProt (v1.6) proteogenomic resource. OpenProt pipeline (dark blue arrows) contains two main features: prediction (on the left, blue) and evidence retrieval (middle, green). OpenProt enforces a polycistronic model of eukaryotic genes contrary to the actual dogma of one CDS per transcript. It retrieves all possible ORFs from transcripts annotated in NCBI RefSeq and/or Ensembl (ORF prediction, blue). The ORF-encoded proteins are then categorized as follows: RefProt if already annotated in NCBI RefSeq, Ensembl and/or UniProt; novel isoforms of known CDS (IL\_ accessions); or novel alternative proteins (IP\_ accessions). All predicted proteins are available on the OpenProt website ([www.openprot.org](http://www.openprot.org)). Furthermore, OpenProt retrieves supporting evidence for all proteins from MS, ribosome profiling, protein conservation, functional domain prediction and related proteins. All data are inserted in the OpenProt resource and freely available to the community (yellow arrow). OpenProt also allows data submission by users for analysis using the OpenProt pipeline (purple dashed arrows). The results are then returned to the users and inserted in OpenProt at the next release (purple dashed arrows). With a symbiotic behaviour between the scientific community, linking experimental data, bioinformatics resources and deep ORF annotation, OpenProt participates in the implementation of novel, experimentally supported proteins in genome annotations and protein databases (red arrows).

(3,33). The initial release of OpenProt was based on genome annotations from December 2015 for each species (24). These were updated to the latest release for each species as of January 2019. This major update necessitated an entire re-run of the OpenProt pipeline (Supplementary Material S1). For example, in humans alone 9873 messenger RNAs (mRNAs) and 2520 non-coding RNAs (ncRNAs) from the NCBI RefSeq annotation, and 1087 mRNAs and 371 ncRNAs from the Ensembl annotation were discarded or had their sequence changed. In parallel, 13 833 novel mRNAs and 2146 novel ncRNAs were included in the latest release of NCBI RefSeq, and 5970 novel mRNAs and 4884 novel ncRNAs in that of Ensembl. Consequently, 4256 RefProts from the NCBI RefSeq annotation and 2785 for Ensembl were discarded or had their amino acid sequence changed, while 4428 RefProts were added from NCBI RefSeq and 7182 from Ensembl. Genome assemblies, annotation releases and the associated number of predicted coding sequences for the OpenProt update are listed in Table 1.

Table 1. OpenProt (v1.6) prediction pipeline output

Species	Genome assembly		Annotations		ORFeome (both annotations)		
	NCBI RefSeq	Ensembl	Total #	Ref #	IL_#	IP_#	
<i>Homo sapiens</i>	GRCh38.p12	GRCh38.95	692 045	134 477	68 612	488 956	
<i>Pan troglodytes</i>	Pan_tro_3.0	Pan_tro_3.0.95	331 247	79 070	14 308	237 869	
<i>Mus musculus</i>	GRCm38.p6	GRCm38.95	558 632	87 339	40 870	430 423	
<i>Rattus norvegicus</i>	Rnor_6.0	Rnor_6.0.95	294 727	51 662	7872	235 193	
<i>Bos taurus</i>	ARS-UCDI.2	ARS-UCDI.2.95	285 565	67 753	11 382	206 430	
<i>Ovis aries</i>	Oar_v3.1	Oar_v3.1.95	162 972	30 283	6339	126 350	
<i>Danio rerio</i>	GRCz11	GRCz11.95	287 990	68 272	11 896	207 822	
<i>Drosophila melanogaster</i>	Release 6 plus ISO1 MT	BDGP6.95	97 834	22 058	2125	73 651	
<i>Caenorhabditis elegans</i>	WBcel235	WBcel235.95	94 890	28 516	3034	63 340	
<i>Saccharomyces cerevisiae</i> S288c	R64-1-1	R64-1-1.95	16 873	6615	28	10 230	

Ref = currently annotated protein (RefProt); IL\_ = novel isoforms of known protein; IP\_ = novel protein from alternative ORF (AltProt).

Table 2. OpenProt fosters annotation of novel isoforms and novel proteins

Species	Predicted AltProt (OP v1.3), now RefProt (OP v1.6) (#)	Predicted novel isoform (OP v1.3), now RefProt (OP v1.6) (#)	Total changes (#)
<i>H. sapiens</i>	69	736	805
<i>P. troglodytes</i>	268	1702	1970
<i>M. musculus</i>	28	150	178
<i>R. norvegicus</i>	0	0	0
<i>B. taurus</i>	98	269	367
<i>O. aries</i>	0	0	0
<i>D. rerio</i>	155	661	816
<i>D. melanogaster</i>	1	1	2
<i>C. elegans</i>	8	51	59
<i>S. cerevisiae</i> S288c	0	0	0

OP = OpenProt; v1.3 = OpenProt version 1.3 [annotations from December 2015 (24)]; v1.6 = current OpenProt version (annotations from January 2019); RefProt = currently annotated protein; novel isoform = novel isoform of known protein; AltProt = novel protein from an alternative ORF.

Interestingly, out of the newly included RefProts, 90 in NCBI RefSeq and 1835 in Ensembl had a length of  $\leq 100$  codons. These numbers reflect the efforts from the small ORF and alternative ORF community to foster annotation of these functional yet overlooked proteins (16,17,24). Thus, in humans 736 novel isoforms and 69 AltProts annotated in the OpenProt initial release are now RefProts since they have been included in annotations (NCBI RefSeq, Ensembl and/or UniProt). For example, the protein previously annotated IP\_211724 in the initial release of OpenProt (<https://openprot.org/p/savedSearch/jCa>) was detected and described (renamed altDDIT3) in a publication building on OpenProt data (7). Consequently, the protein was included in the Ensembl annotation (ENSP00000494177) and in UniProt (A0A2R8YD15) in 2018, and became a RefProt in the latest OpenProt release (<https://openprot.org/p/savedSearch/kCa>). The numbers of AltProts and novel isoforms that have been included in NCBI RefSeq, Ensembl and/or UniProt since the initial release of OpenProt, across all supported species, are listed in Table 2.

### ORF calling from ribosome profiling datasets

In its initial release, OpenProt only reported evidence of translation for ORFs starting with an AUG. Yet pervasive translation with alternative start codons is increasingly reported by the community (14–17), with up to 83% of the sORF repository consisting of non-AUG small ORFs (28). The latest release of OpenProt now integrates data-driven annotation of non-AUG initiating codons. The pipeline of analysis for ribosome profiling datasets (RIBO-seq) uses the PRICE algorithm (21), which can accurately predict non-AUG start ORFs (Supplementary Material S1). Briefly, PRICE uses a logistic regression model to accurately predict initiating codons in RIBO-seq data, whether the dataset is enriched or not for initiating ribosomes. PRICE thus reconstitutes a set of codons most likely to yield the observed reads. We observed that when including all codons as possible initiation sites, the *P*-value of the reported ORF would increase as the overlap with the ORF predicted by OpenProt decreases (Supplementary Material S2). In PRICE, the *P*-value is the result of a generalized binomial test; thus, it in-

icates the confidence of that ORF not being attributable to noise. As the length of the ORF decreases, it is expected to be more difficult to distinguish true translation event from noise. In order to only annotate the most confident alternative initiation starts, we set the following filters based on our observation (Supplementary Material S2): any codon as initiating codon, and an overlap of the ORF candidate with the ORF predicted by OpenProt above 70%. Using these new filters, almost half a million (488 231) translation events could be retrieved in human datasets, corresponding to 33 836 ORFs across OpenProt.

### Peptide unicity in MS datasets

Unique and recurrent peptide detection in datasets of MS-based proteomics is considered a gold standard evidence of novel protein expression (1,34). However, the characterization of a unique peptide is often dependent on the database used at the time of the analysis. In classical MS analyses, a peptide is considered unique if it matches to one and only one protein. However, if a protein is not included in the database, it will not be taken into account in the evaluation of the unicity of the peptide. In the initial release of OpenProt, we ensured unicity by using an exhaustive proteome derived from both NCBI RefSeq and Ensembl annotations, as well as UniProt (24). Yet, as mentioned earlier, annotations evolve and with that the proteomic landscape. In the latest OpenProt release, we ensured a control of peptide unicity from MS datasets through time and releases. At each OpenProt release, the set of peptides mapped from all MS datasets to proteins is rescanned for protein assignment enforcing the OpenProt assignment rules. (i) When a peptide matches to multiple protein sequences from different genes, it is discarded. (ii) When a peptide matches to multiple protein sequences from the same gene, with at least one of the proteins being a RefProt, the peptide is given to the RefProt(s) only. (iii) When a peptide matches multiple AltProts or novel isoforms but not RefProts, the peptide is assigned to all AltProts and novel isoforms. Using such rules, OpenProt reports MS-based evidence of expression of novel proteins (AltProts and novel isoforms) only if the detected peptide does not match any known protein.

Furthermore, during this control of peptide unicity, the isobaric residues leucine and isoleucine are treated as undistinguishable. For example, the peptide EIGNLISDAMK was called by our SearchGUI/PeptideShaker MS pipeline to support the identification of IP\_658154 (HSPD1P7). However, this peptide only differs by its leucine from the canonical protein of the parental gene HSPD1, which contains the peptide EIGNIISDAMK. Thus, this peptide was re-assigned as EIGNIISDAMK to the RefProt (P10809).

### MS statistics to guide data interpretation

Based on the OpenProt pipeline and stringent peptide assignment criteria regarding novel proteins, one must realize that not all predicted AltProts or novel isoforms are detectable by MS (35). To help the OpenProt users with the interpretation of an MS score of 0, we added MS coverage statistics to the MS tab of the details page for each protein. The coverage statistics tab contains the number of independent datasets in which the protein was detected, the number

of unique peptides detected across all datasets and the number of peptide spectrum matches across all datasets. Alongside these metrics, the theoretical and current coverages by MS for each given protein are indicated (named possible sequence coverage and detected sequence coverage, respectively). The theoretical coverage is calculated from all possible tryptic peptides that would fit the OpenProt criteria to be assigned: a minimal length of seven amino acids, a maximal mass of 4600 Da and peptide unicity given the protein type (RefProt, novel isoform or AltProt). Thus, in *H. sapiens* 44 350 of OpenProt annotated proteins would not be detectable by MS, 13 595 in *P. troglodytes*, 31 275 in *M. musculus*, 11 311 in *R. norvegicus*, 11 454 in *B. taurus*, 8592 in *O. aries*, 15 294 in *D. rerio*, 1575 in *D. melanogaster*, 2267 in *C. elegans* and 245 in *S. cerevisiae* S288c.

A summary of experimental detections across all species supported by OpenProt is provided in Table 3 and general statistics are now provided in the About page of the OpenProt website (<https://openprot.org/p/about>; Supplementary Material S3) alongside a link to a stand-alone, freely accessible version of the script for calculating MS statistics.

### Identification of similarities of coding sequences within genes

In this OpenProt update, the relationships between proteins from a same gene are now reported. In order to predict proteins as isoforms, OpenProt evaluates the protein sequence identity between proteins from a same gene using an all-versus-all BLAST (Basic Local Alignment Search Tool) (36). OpenProt reports proteins as isoforms if the result of their BLAST search yields a bit score over 40 for an overlap over 50% of the queried sequence, as previously published (37,38). The bit score was chosen as it is dependent on the size of the alignment, but not on the size of the database used. Thus, the higher the bit score, the better the sequence similarity. This isoform feature allows the user to grasp the complexity brought by alternative splicing and by the polycistronic nature of a transcript. For example, the AltProt IP\_191523 from the *PIDD1* gene is related to six other AltProts from different transcripts of the *PIDD1* gene. These range from 284 amino acids (IP\_191530) to 109 (IP\_784984) (Figure 2A). In parallel, the canonical protein of the same transcript (ENSP00000416801) has eight isoforms ranging from 934 amino acids (XP\_005253063) to 526 (Q9HB75-5).

Furthermore, OpenProt now also displays for novel isoforms (accessions starting with IL\_) the RefProt(s) responsible for the annotation as isoform instead of AltProt. This information is displayed at the top of the details page for each novel isoform and in detail in the isoforms tab. For example, the novel isoform IL\_794710 is annotated isoform because of four RefProts sharing identical genomic coordinates (start and/or end) and a sequence identity over 20% of the protein length. These are NP\_001310470, NP\_071322 and XP\_005270110 from NCBI RefSeq, and ENSP00000359149 from Ensembl (Figure 2B).

### Database development

All data are generated using in-house Perl (version 5.18.2) and Python (version 3.6.9) scripts and stored in a PostgreSQL database (version 9.6). All re-analysed MS and ri-

Table 3. OpenProt (v1.6) evidence collection output

Species	Conservation evidence			Translation evidence (RIBO-seq)			Protein evidence (MS)					
	Sp #	Ref #	IL #	IP #	St #	Ref #	IL #	IP #	St #	Ref #	IL #	IP #
<i>H. sapiens</i>	9	116 295	45 938	217 897	69	31 453	6912	9738	95	120 925	2446	40 883
<i>P. troglodytes</i>	9	73 137	13 315	131 321	0	N/A	N/A	N/A	2	25 821	15	360
<i>M. musculus</i>	9	78 193	25 867	98 047	40	23 740	3309	5908	40	66 875	236	4458
<i>R. norvegicus</i>	9	49 994	6012	77 768	3	16 205	996	1731	8	21 210	16	398
<i>B. taurus</i>	9	64 866	7148	83 993	0	N/A	N/A	N/A	4	20 120	6	80
<i>O. aries</i>	9	30 091	6223	73 034	0	N/A	N/A	N/A	4	8618	22	104
<i>D. rerio</i>	9	58 386	6826	26 669	5	11 186	490	299	8	32 286	27	220
<i>D. melanogaster</i>	9	13 567	757	477	3	5370	147	243	4	11 560	24	168
<i>C. elegans</i>	9	13 182	1064	450	5	12 796	396	196	6	13 332	38	96
<i>S. cerevisiae</i> S288c	9	2895	5	34	6	5429	4	269	6	4191	0	44

Sp = number of species evaluated for orthology relationships (not counting the queried species); St = number of studies re-analysed by OpenProt; Ref = currently annotated CDS (RefORF); IL = novel isoforms of known CDS; IP = novel CDS from alternative ORF (AltORF); N/A = when no dataset has been re-analysed for this species yet (OpenProt release 1.2). Conservation evidence = all proteins with at least one ortholog in at least one species. Translation evidence = all ORFs detected in at least one detection by PRICE analysis of RIBO-seq data. Protein evidence = all proteins with at least one unique peptide in at least one study.

bosome profiling studies are accessible from the Help page (<https://openprot.org/p/help>; Supplementary Material S4).

## USER INTERFACE UPDATES

### Summary of the features presented in the initial release

At the time of the initial release, the OpenProt website provided three interfaces: a genome browser, a search page and a platform for downloads. The genome browser enables rapid lookup of all transcripts, proteins and detected peptides for a specific genomic region, while the search interface is designed for exploration of specific genes, transcripts and/or proteins. The platform for downloads allows custom retrieval of any data on the OpenProt resource as tsv, fasta or bed files (24,35). Each predicted ORF and encoded protein annotated in OpenProt has its own page that contains a genome browser centred on the dedicated ORF alongside all supporting information. All data from the initial release are still present on the website (under the release number 1.3); however, as genome annotations evolve, so does OpenProt, and we strongly encourage using the latest OpenProt release.

### Novel OpenProt web interface

The OpenProt resource is hosted at [www.openprot.org](http://www.openprot.org) since its initial release. The graphics of the website have, however, considerably changed for a more user-friendly and graphical interface. The home page (<https://openprot.org/>) now hosts a presentation of the project with tutorials and visual explanations. The query interface has been updated with an additional advanced search filter. The users can now filter results for specific MS or ribosome profiling datasets (Figure 1). Furthermore, a novel interface has been added for the submission of MS or ribosome profiling datasets to be analysed by OpenProt. This novel submission platform allows users to query their datasets for unannotated proteins, independently of their bioinformatics skills or computational resources. To ensure the quality of the data inserted into the OpenProt database, users first have to deposit their data into public repositories, the PRIDE archive (39) or ProteomeXchange (40) for MS datasets and Gene Omnibus (41) for ribosome profiling ones. After submission, the parameters are validated by the members of the OpenProt team prior to the analysis with the OpenProt pipeline (24) (Supplementary Material S1). Upon completion of the analysis, the results are sent to the users and the data are stored to be inserted in the next OpenProt release. For computational reasons, releases will be launched when 10 datasets are ready for implementation, or if it is 60 days since a dataset has been analysed. The Help page and contents have been updated to support this new interface (<https://openprot.org/p/help>; Supplementary Material S5).

### Website development

The OpenProt web platform was built using the Flask framework (version 1.1.2) and developed using HTML, SQL and JavaScript. The OpenProt website uses an HTTPS protocol to ensure protection of personal information.

A

OpenProt Home Browse Search Submit Study Downloads About Help

back to main table

IP\_191523

Info Mass spectrometry (48) Translation (2) Domains (1) Conservation (1) Isoforms (6)

Identity (%) 30 40 50 60 70 80 90 100

Isoforms

- IP\_191524
- IP\_191527
- IP\_784984
- IP\_2314968
- IP\_191530
- IP\_784982

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_191524

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
100	171	0	0	1	171	55	225	1e-109	318	100	171	225

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_191527 IP\_191527

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
100	171	0	0	1	171	94	264	2e-109	319	100	171	264

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_784984 IP\_784984

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
100	109	0	0	1	109	1	109	8e-65	199	64	171	109

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_2314968 IP\_2314968

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
100	171	0	0	1	171	36	206	1e-109	317	100	171	206

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_191530 IP\_191530

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
89.53	191	0	1	1	171	94	284	1e-104	308	100	171	284

**Blast**  
Query: IP\_191523 (Homo sapiens)  
Subject: IP\_784982 IP\_784982 IP\_784982 IP\_784982 IP\_784982

pident	length	mismatch	gapopen	qstart	qend	sstart	send	eval	bitscore	qcovs	qlen	slen
89.53	191	0	1	1	171	1	191	7e-106	307	100	171	191

B

OpenProt Home Browse Search Submit Study Downloads About Help

back to main table

IL\_794710

This protein was annotated as a novel isoform of NP\_001310470.1NP\_071322.1XP\_005270110.2ENSP00000359149.2 Please refer to the isoform tab for details.

Info Mass spectrometry (45) Translation (6) Domains (16) Conservation (8) Isoforms (3)

Reasons for the annotation as novel isoform

Protein accession	Reason for isoform annotation
NP_001310470.1	identical genomic coordinates of start or end codon with a sequence identity over 20% of the protein length
NP_071322.1	identical genomic coordinates of start or end codon with a sequence identity over 20% of the protein length
XP_005270110.2	identical genomic coordinates of start or end codon with a sequence identity over 20% of the protein length
ENSP00000359149.2	identical genomic coordinates of start or end codon with a sequence identity over 20% of the protein length

**Figure 2.** Annotation of novel protein relationships in OpenProt. (A) Content of the isoforms tab from the details page for IP\_191523 on OpenProt (v1.6—available here). The protein accession always sits at the top left corner of the page. The isoform tab displayed here is the last tab of the page. The number indicated at the top corresponds to the number of identified isoforms. The tab contains a protein tree where nodes correspond to unique protein sequences present in OpenProt and are coloured based on their identity to the looked-up protein (here, IP\_191523). Each node is accompanied by the corresponding protein accession. By clicking on the protein accession, a pop-up window (represented in dotted boxes) appears with the results of the BLAST search. (B) Content of the top of the isoforms tab from the details page for IL\_794710 on OpenProt (v1.6—available here). The protein accession always sits at the top left corner of the page. In the case of novel isoforms (accession starting with IL\_), a phrase under the accession indicates the RefProts responsible for the novel isoform annotation (circled in a red dotted line). The top of the isoform tab, displayed here, will thus contain an additional table. This table contains the accession of the RefProts responsible for the isoform annotation of the looked-up protein (here, IL\_794710), and the associated reason. Please note that because the filter for a novel protein to be annotated as novel isoform is more inclusive than that of the prediction of related proteins from the same gene, the RefProts listed in the table may not appear in the predicted isoform tree.

OpenProt is accessible via all major web browsers supporting JavaScript, such as Safari, Firefox, Chrome or Internet Explorer. All pages can be viewed on mobiles, but the interfaces have been optimized for display on computers or tablets.

## DISCUSSION AND COMPARISON TO EXISTING RESOURCES

The inaccuracies of current genome annotations have been increasingly reported in the last decades (1,4,14,28,29,42,43), and the OpenProt resource has offered a systematic approach to mend the gap between annotations and experimental data (24,35). OpenProt annotates thousands of novel predicted proteins supported by experimental evidence and functional predictions. As our pipeline evolves and more RIBO-seq and MS datasets are constantly added to the database, evidence for functional yet unannotated proteins is growing. At the time of this manuscript, the OpenProt resource lists 1477 novel proteins (AltProts or novel isoforms) in humans that were detected by both ribosome profiling and MS. Out of these, 935 are located on at least one mRNA that contains another annotated coding sequence. These numbers highlight not only the need for resources like the OpenProt database, but also the polycistronic nature of eukaryotic transcripts. Indeed, in each species for which OpenProt re-analysed at least one MS dataset and one ribosome profiling dataset, novel proteins detected by both methods and encoded in at least one mRNA containing another annotated sequence were reported. In particular, 98 were reported in *M. musculus*, 35 in *R. norvegicus*, 23 in *C. elegans*, 16 in *D. melanogaster*, 4 in *S. cerevisiae* and 2 in *D. rerio*. As the number of available datasets for these species increases, we expect these numbers of detection to rise.

To the best of our knowledge, OpenProt was the first database to fully endorse a polycistronic model of eukaryotic genome annotation. OpenProt differs from other small ORF databases (16,17) in that it does not uphold a maximum length threshold (below 100 codons for smORFs); it allows for multiple ORFs per transcript, supports two transcriptome annotations and allows for the identification and detection of novel isoforms of annotated proteins. Thus, OpenProt reaches a deeper ORF annotation throughout the genome. Furthermore, OpenProt now supports data submission to be analysed using the pipeline of OpenProt and with the results returned to the users. To the best of our knowledge, other small ORF databases do not offer this service although they may accept data suggestions (16).

In addition, OpenProt differentiates itself from UniProt (27) or NextProt (44) as it provides a genome browser and implements a polycistronic annotation model. Notwithstanding, as UniProt and NextProt are well-established proteomic resources, when a novel protein regroups sufficient evidence to meet UniProt and NextProt annotation requirements, it becomes a RefProt in the OpenProt database with the UniProt accession for ease of translation between databases. This symbiotic behaviour is exemplified by hundreds of novel proteins that have been annotated in NCBI RefSeq, Ensembl and/or UniProt since the first release of OpenProt (Table 2).

With the necessity of careful curation for the annotation of novel proteins comes the need for thorough and meticulous groundwork that annotation consortia are not designed to conduct (1,3,4). OpenProt and other small ORF resources are thus of utmost importance to foster less serendipitous discoveries of novel proteins and further our understanding of biological questions.

## FUTURE DIRECTIONS

Since its initial release (24), the OpenProt website counted over 36 000 visits and about 3000 downloads. The resource was used in 17 publications to discover novel proteins (5 of these were from our lab or collaborators) (30–32,45–57). This shows the OpenProt resource answered a need of the scientific community. The OpenProt development is heavily driven by suggestions from the community and recommendations of new features, species or experimental evidence are always welcome. These can be submitted via the OpenProt discussion forum.

The OpenProt pipeline is automated so that iGenomes updates and releases of NCBI RefSeq, Ensembl and/or UniProt are taken into account. The data are updated at the beginning of every odd year to ease management of computational resources access. OpenProt is a release-based platform, developed in accordance to the FAIR guiding principles for scientific data management and stewardship (58). This ensures an up-to-date, continuous availability of all OpenProt data through time. Furthermore, the shareable links created from the ‘Share’ feature at the top of the results table on the query interface are also release-based and persistent in time. This allows the inclusion of such links in publications for an exact snapshot of the OpenProt annotation at the time. All of the scripts behind OpenProt are available upon reasonable requests to the authors. Stand-alone scripts related to the mining of OpenProt data are available in the About page, under the Related Scripts section (<https://openprot.org/p/about>).

In future releases, OpenProt aims to tackle remaining challenges. For example, OpenProt is currently developing cutting-edge machine learning algorithms (e.g. convolutional neural networks) to more accurately predict coding sequences throughout genomes. Using such algorithms, OpenProt aims to insert ORFs shorter than 30 codons while avoiding spurious annotations. Furthermore, new tools and features will be added, such as a mass spectra viewer for evidence of protein detection, visualization of ribosomal coverage on the genome browser, networks of detected protein–protein interactions from affinity purification MS experiments and integration of genomic variants.

The quantity and quality of data provided by OpenProt along with its ease of use and transparent data availability hold potential to make it a long-lasting and popular proteogenomic resource.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

## ACKNOWLEDGEMENTS

We thank our collaborators for providing additional MS datasets. We thank Florian Erhard for his help updating the PRICE pipeline on RIBO-seq data. We also thank Mylène Brunelle, Alan A. Cohen and Darel Hunting for helpful discussions and contributions. X.R. is a member of the Fonds de Recherche du Québec Santé (FRQS)-supported Centre de Recherche du Centre Hospitalier Universitaire de Sherbrooke. We thank the team at Calcul Québec and Compute Canada for their support with the use of the supercomputer mp2 from Université de Sherbrooke. S.L. was supported by a CIHR scholarship for graduate studies. Operation of the mp2 supercomputer was funded by the Canada Foundation of Innovation (CFI), le ministère de l'Économie, de la science et de l'innovation du Québec (MESI) and les Fonds de Recherche du Québec—Nature et technologies (FRQ-NT).

## FUNDING

Canada Research Chairs [to X.R.]; Canadian Institutes of Health Research (CIHR) [MOP-137056 and MOP-136962]. Funding for open access charge: Canada Research Chairs.

*Conflict of interest statement.* None declared.

## REFERENCES

- Brunet, M.A., Levesque, S.A., Hunting, D.J., Cohen, A.A. and Roucou, X. (2018) Recognition of the polycistronic nature of human genes is critical to understanding the genotype–phenotype relationship. *Genome Res.*, **28**, 609–624.
- Cheng, H., Chan, W.S., Li, Z., Wang, D., Liu, S. and Zhou, Y. (2011) Small open reading frames: current prediction techniques and future prospect. *Curr. Protein Pept. Sci.*, **12**, 503–507.
- Mudge, J.M. and Harrow, J. (2016) The state of play in higher eukaryote gene annotation. *Nat. Rev. Genet.*, **17**, 758–772.
- Orr, M.W., Mao, Y., Storz, G. and Qian, S.-B. (2019) Alternative ORFs and small ORFs: shedding light on the dark proteome. *Nucleic Acids Res.*, **48**, 1029–1042.
- Olexiouk, V. and Menschaert, G. (2016) Identification of small novel coding sequences, a proteogenomics endeavor. *Adv. Exp. Med. Biol.*, **926**, 49–64.
- Hellens, R.P., Brown, C.M., Chisnall, M.A.W., Waterhouse, P.M. and Macknight, R.C. (2016) The emerging world of small ORFs. *Trends Plant Sci.*, **21**, 317–328.
- Samandi, S., Roy, A.V., Delcourt, V., Lucier, J.-F., Gagnon, J., Beaudoin, M.C., Vanderperre, B., Breton, M.-A., Motard, J., Jacques, J.-F. *et al.* (2017) Deep transcriptome annotation enables the discovery and functional characterization of cryptic small proteins. *eLife*, **6**, e27860.
- Menschaert, G., Crieckinge, W.V., Notelaers, T., Koch, A., Crappé, J., Gevaert, K. and Damme, P.V. (2013) Deep proteome coverage based on ribosome profiling aids mass spectrometry-based protein and peptide discovery and provides evidence of alternative translation products and near-cognate translation initiation events. *Mol. Cell. Proteomics*, **12**, 1780–1790.
- Ma, J., Ward, C.C., Jungreis, I., Slavoff, S.A., Schwaid, A.G., Neveu, J., Budnik, B.A., Kellis, M. and Saghatelian, A. (2014) Discovery of human sORF-encoded polypeptides (SEPs) in cell lines and tissue. *J. Proteome Res.*, **13**, 1757–1765.
- Ingolia, N.T. (2014) Ribosome profiling: new views of translation, from single codons to genome scale. *Nat. Rev. Genet.*, **15**, 205–213.
- Andreev, D.E., O'Connor, P.B.F., Zhdanov, A.V., Dmitriev, R.I., Shatsky, I.N., Papkovsky, D.B. and Baranov, P.V. (2015) Oxygen and glucose deprivation induces widespread alterations in mRNA translation within 20 minutes. *Genome Biol.*, **16**, 90.
- Andreev, D.E., O'Connor, P.B.F., Fahey, C., Kenny, E.M., Terenin, I.M., Dmitriev, S.E., Cormican, P., Morris, D.W., Shatsky, I.N. and Baranov, P.V. (2015) Translation of 5' leaders is pervasive in genes resistant to eIF2 repression. *eLife*, **4**, e03971.
- Bazzini, A.A., Johnstone, T.G., Christiano, R., Mackowiak, S.D., Obermayer, B., Fleming, E.S., Vejnár, C.E., Lee, M.T., Rajewsky, N., Walther, T.C. *et al.* (2014) Identification of small ORFs in vertebrates using ribosome footprinting and evolutionary conservation. *EMBO J.*, **33**, 981–993.
- Chen, J., Brunner, A.-D., Cogan, J.Z., Nuñez, J.K., Fields, A.P., Adamson, B., Itzhak, D.N., Li, J.Y., Mann, M., Leonetti, M.D. *et al.* (2020) Pervasive functional translation of noncanonical human open reading frames. *Science*, **367**, 1140–1146.
- Ingolia, N.T. (2016) Ribosome footprint profiling of translation throughout the genome. *Cell*, **165**, 22.
- Olexiouk, V., Van Crieckinge, W. and Menschaert, G. (2018) An update on sORFs.org: a repository of small ORFs identified by ribosome profiling. *Nucleic Acids Res.*, **46**, D497–D502.
- Hao, Y., Zhang, L., Niu, Y., Cai, T., Luo, J., He, S., Zhang, B., Zhang, D., Qin, Y., Yang, F. *et al.* (2017) SmProt: a database of small proteins encoded by annotated coding and non-coding RNA loci. *Brief. Bioinform.*, **19**, 636–643.
- Xie, S.-Q., Nie, P., Wang, Y., Wang, H., Li, H., Yang, Z., Liu, Y., Ren, J. and Xie, Z. (2016) RPFdb: a database for genome wide information of translated mRNA generated from ribosome profiling. *Nucleic Acids Res.*, **44**, D254–D258.
- Ingolia, N.T., Ghaemmaghami, S., Newman, J.R.S. and Weissman, J.S. (2009) Genome-wide analysis *in vivo* of translation with nucleotide resolution using ribosome profiling. *Science*, **324**, 218–223.
- Raj, A., Wang, S.H., Shim, H., Harpak, A., Li, Y.I., Engelmann, B., Stephens, M., Gilad, Y. and Pritchard, J.K. (2016) Thousands of novel translated open reading frames in humans inferred by ribosome footprint profiling. *eLife*, **5**, e13328.
- Erhard, F., Halenius, A., Zimmermann, C., L'Hernault, A., Kowalewski, D.J., Weekes, M.P., Stevanovic, S., Zimmer, R. and Dölken, L. (2018) Improved Ribo-seq enables identification of cryptic translation events. *Nat. Methods*, **15**, 363–366.
- Reixachs-Solé, M., Ruiz-Orera, J., Albà, M.M. and Eyras, E. (2020) Ribosome profiling at isoform level reveals evolutionary conserved impacts of differential splicing on the proteome. *Nat. Commun.*, **11**, 1768.
- Cui, H., Hu, H., Zeng, J. and Chen, T. (2019) DeepShape: estimating isoform-level ribosome abundance and distribution with Ribo-seq data. *BMC Bioinformatics*, **20**, 678.
- Brunet, M.A., Brunelle, M., Lucier, J.-F., Delcourt, V., Levesque, M., Grenier, F., Samandi, S., Leblanc, S., Aguilar, J.-D., Dufour, P. *et al.* (2019) OpenProt: a more comprehensive guide to explore eukaryotic coding potential and proteomes. *Nucleic Acids Res.*, **47**, D403–D410.
- O'Leary, N.A., Wright, M.W., Brister, J.R., Ciuffo, S., Haddad, D., McVeigh, R., Rajput, B., Robbertse, B., Smith-White, B., Ako-Adjei, D. *et al.* (2016) Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.*, **44**, D733–D745.
- Yates, A.D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M.R., Armean, I.M., Azov, A.G., Bennett, R. *et al.* (2020) Ensembl 2020. *Nucleic Acids Res.*, **48**, D682–D688.
- Consortium, T.U. (2019) UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.*, **47**, D506–D515.
- Brunet, M.A., Leblanc, S. and Roucou, X. (2020) Reconsidering proteomic diversity with functional investigation of small ORFs and alternative ORFs. *Exp. Cell Res.*, **393**, 112057.
- Peeters, M. and Menschaert, G. (2020) The hunt for sORFs: a multidisciplinary strategy. *Exp. Cell Res.*, **391**, 111923.
- Dubois, M.-L., Meller, A., Samandi, S., Brunelle, M., Frion, J., Brunet, M.A., Toupin, A., Beaudoin, M.C., Jacques, J.-F., Lévesque, D. *et al.* (2020) UBB pseudogene 4 encodes functional ubiquitin variants. *Nat. Commun.*, **11**, 1306.
- Cardon, T., Franck, J., Coyaud, E., Laurent, E.M.N., Damato, M., Maffia, M., Vergara, D., Fournier, I. and Salzet, M. (2020) Alternative proteins are functional regulators in cell reprogramming by PKA activation. *Nucleic Acids Res.*, **48**, 7864–7882.
- Cao, X., Khitun, A., Na, Z., Dumitrescu, D.G., Kubica, M., Olatunji, E. and Slavoff, S.A. (2020) Comparative proteomic profiling of unannotated microproteins and alternative proteins in human cell lines. *J. Proteome Res.*, **19**, 3418–3426.

33. Salzberg, S.L. (2019) Next-generation genome annotation: we still struggle to get it right. *Genome Biol.*, **20**, 92.
34. Mann, M. (2016) Origins of mass spectrometry-based proteomics. *Nat. Rev. Mol. Cell Biol.*, **17**, 678.
35. Brunet, M.A., Lekehal, A.M. and Roucou, X. (2020) How to illuminate the dark proteome using the multi-omic OpenProt resource. *Curr. Protoc. Bioinformatics*, **71**, e103.
36. Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment search tool. *J. Mol. Biol.*, **215**, 403–410.
37. Sonnhammer, E.L.L. and Östlund, G. (2015) InParanoid 8: orthology analysis between 273 proteomes, mostly eukaryotic. *Nucleic Acids Res.*, **43**, D234–D239.
38. Chen, H., Shaw, D., Zeng, J., Bu, D. and Jiang, T. (2019) DIFFUSE: predicting isoform functions from sequences and expression profiles via deep learning. *Bioinformatics*, **35**, i284–i294.
39. Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J., Inuganti, A., Griss, J., Mayer, G., Eisenacher, M. *et al.* (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res.*, **47**, D442–D450.
40. Deutsch, E.W., Bandeira, N., Sharma, V., Perez-Riverol, Y., Carver, J.J., Kundu, D.J., García-Seisdedos, D., Jarnuczak, A.F., Hewapathirana, S., Pullman, B.S. *et al.* (2020) The ProteomeXchange consortium in 2020: enabling 'big data' approaches in proteomics. *Nucleic Acids Res.*, **48**, D1145–D1152.
41. Barrett, T., Wilhite, S.E., Ledoux, P., Evangelista, C., Kim, I.F., Tomashevsky, M., Marshall, K.A., Phillippy, K.H., Sherman, P.M., Holko, M. *et al.* (2013) NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Res.*, **41**, D991–D995.
42. Merino-Valverde, I., Greco, E. and Abad, M. (2020) The microproteome of cancer: from invisibility to relevance. *Exp. Cell Res.*, **392**, 111997.
43. Landry, C.R., Zhong, X., Nielly-Thibault, L. and Roucou, X. (2015) Found in translation: functions and evolution of a recently discovered alternative proteome. *Curr. Opin. Struct. Biol.*, **32**, 74–80.
44. Zahn-Zabal, M., Michel, P.-A., Gateau, A., Nikitin, F., Schaeffer, M., Audot, E., Gaudet, P., Duek, P.D., Teixeira, D., Rech de Laval, V. *et al.* (2020) The neXtProt knowledgebase in 2020: data, tools and usability improvements. *Nucleic Acids Res.*, **48**, D328–D334.
45. Cardon, T., Salzet, M., Franck, J. and Fournier, I. (2019) Nuclei of HeLa cells interactomes unravel a network of ghost proteins involved in proteins translation. *Biochim. Biophys. Acta: Gen. Subj.*, **1863**, 1458–1470.
46. Fesenko, I., Kirov, I., Kniazev, A., Khazigaleeva, R., Lazarev, V., Kharlampieva, D., Grafkaia, E., Zgoda, V., Butenko, I., Arapidi, G. *et al.* (2019) Distinct types of short open reading frames are translated in plant cells. *Genome Res.*, **29**, 1464–1477.
47. Brunet, M.A. and Roucou, X. (2019) Mass spectrometry-based proteomics analyses using the OpenProt database to unveil novel proteins translated from non-canonical open reading frames. *J. Vis. Exp.*, doi:10.3791/59589.
48. Kinyry, S.J., Michel, A.M. and Baranov, P.V. (2020) Computational methods for ribosome profiling data analysis. *WIREs RNA*, **11**, e1577.
49. Cardon, T., Hervé, F., Delcourt, V., Roucou, X., Salzet, M., Franck, J. and Fournier, I. (2020) Optimized sample preparation workflow for improved identification of ghost proteins. *Anal. Chem.*, **92**, 1122–1129.
50. Vergara, D., Verri, T., Damato, M., Trerotola, M., Simeone, P., Franck, J., Fournier, I., Salzet, M. and Maffia, M. (2020) A hidden human proteome signature characterizes the epithelial mesenchymal transition program. *Curr. Pharm. Des.*, **26**, 372–375.
51. Wang, B., Hao, J., Pan, N., Wang, Z., Chen, Y. and Wan, C. (2020) Identification and analysis of small proteins and short open reading frame encoded peptides in Hep3B cell. *J. Proteomics*, **230**, 103965.
52. Simoneau, J., Gosselin, R. and Scott, M.S. (2020) Factorial study of the RNA-seq computational workflow identifies biases as technical gene signatures. *NAR Genomics Bioinform.*, **2**, lqaa043.
53. Erady, C., Chong, D., Meena, N., Puntambekar, S., Chauhan, R., Umrانيا, Y., Andreani, A., Nel, J., Wayland, M.T., Pina, C. *et al.* (2019) Translational products encoded by novel ORFs may form protein-like structures and have biological functions. bioRxiv doi: <https://doi.org/10.1101/567800>, 05 March 2019, preprint: not peer reviewed.
54. Erady, C., Puntambekar, S. and Prabakaran, S. (2020) Use of short-read RNA-seq data to identify transcripts that can translate novel ORFs. bioRxiv doi: <https://doi.org/10.1101/2020.03.21.001883>, 23 March 2020, preprint: not peer reviewed.
55. Brunet, M.A., Jacques, J.-F., Nassari, S., Tyzack, G.E., McGoldrick, P., Zinman, L., Jean, S., Robertson, J., Patani, R. and Roucou, X. (2020) FUS gene is dual-coding with both proteins united in FUS-mediated toxicity. bioRxiv doi: <https://doi.org/10.1101/848580>, 14 April 2020, preprint: not peer reviewed.
56. Jagannathan, N.S., Meena, N., Bhayankaram, K.P. and Prabakaran, S. (2019) Proteins encoded by novel ORFs have increased disorder but can be biochemically regulated and harbour deleterious mutations. bioRxiv doi: <https://doi.org/10.1101/562835>, 29 March 2019, preprint: not peer reviewed.
57. Murgoci, A.-N., Cardon, T., Aboulouard, S., Duhamel, M., Fournier, I., Cizkova, D. and Salzet, M. (2020) Reference and ghost proteins identification in rat C6 glioma extracellular vesicles. *iScience*, **23**, 101045.
58. Wilkinson, M.D., Dumontier, M., Aalbersberg, I.J.J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L.B., Bourne, P.E. *et al.* (2016) The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data*, **3**, 160018.