



**HAL**  
open science

## The role of active touch: differential mechanism in blindness

Maria Casado-Palacios, Claudio Campus, Alessia Tonelli, Monica Gori

### ► To cite this version:

Maria Casado-Palacios, Claudio Campus, Alessia Tonelli, Monica Gori. The role of active touch: differential mechanism in blindness. Cognitive Neuroscience Society Annual Meeting, Apr 2022, San Francisco, United States. hal-04192656

**HAL Id: hal-04192656**

**<https://hal.univ-lille.fr/hal-04192656>**

Submitted on 31 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# The role of active touch: differential mechanism in blindness

1. Unit for Visually Impaired people (U-VIP), Istituto Italiano di Tecnologia, Genova Italy  
2. University of Genoa, Italy



Casado M.<sup>1,2</sup>,



Campus C.<sup>1</sup>,



Tonelli A.<sup>1</sup>,



Gori M.<sup>1</sup>



This project has received funding from the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No 860114

## Introduction

Research focus on active touch lead to the conclusion that its advantage on the perception might depend on the experimental task [2]. In fact, passive and active touch are two different process, with some studies showing a suppression of afferent information to the somatosensory cortex during active touch. This is called *movement-related sensory gating* and could lead to a worst encoding [3]. Another open discussion refers to the possible enhanced tactile sensitivity in blind individuals, with some authors reporting better performance in this group compared to their sighted counterparts [4], while others found no differences between the groups [5], highlighting the important role of familiarity and experience in their performance [6].

With this work, we wanted to shed some light into the role of active touch in sighted and blind individuals using dynamic stimuli discrimination.

## Method

### Participants:

- 18 sighted
- 18 blind

### Task:


2AFC - The participant has to perceive a sequence of two movement with different speeds and to discriminate which was faster between the two.

Standard velocity:

3 cm/s

Comparison speed:

QUEST [6]




### Conditions:




- Passive touch
- Active touch

### Stimulation:

- Tactile stimulus: 10 cycles/cm
- Tactile area of stimulation: fingertip of index



## Results



Group\*Action  $p=.035^*$

Post Hoc: significant differences between Blind-Active and Blind-Passive  $p<.001^{***}$

## Conclusions

### Sighted individuals:

No differences between the Active and Passive conditions

The similar performance might be caused by an enhanced transmission due to central influences, such as attention and motor set [1]. Also, as active touch involves kinaesthesia and proprioception combined with the cutaneous perception [7] the integration of this information might also contribute to maintain the threshold.

### Blind individuals:

Significant worse performance in the active condition

Blind individuals might be more sensitive to movement-related sensory gating. This might be due to the weakened proprioceptive spatial representations [8] and their difficulty to optimally integrate multisensory information [9].

## References

1. Elaine Chapman, C., Tremblay, F., & Ageranioti-Bélanger, S. A. (1996). Role of Primary Somatosensory Cortex in Active and Passive Touch. *Hand and Brain*, 18(1), 329–347.
2. Güçlü, B., & Murat, A. (2007). Active touch does not improve sequential processing in a counting task. *Acta Neurobiologiae Experimentalis*, 67(2), 165–169.
3. (Chapman, C. E. (1994). Active versus passive touch: Factors influencing the transmission of somatosensory signals to primary somatosensory cortex. *Canadian Journal of Physiology and Pharmacology*, 72(5), 558–570.
4. Postma A, Zuidhoek S, Noordzij ML, Kappers AML (2007) Differences between early-blind, late-blind, and blindfoldedsighted people in haptic spatial-configuration learning and resulting memory traces. *Perception* 36(8):1253–1265
5. Abramowicz A, Klatzky RL, Lederman SJ (2010) Learning and generalization in haptic classification of 2-D raised-line drawings of facial expressions of emotion by sighted and adventitiously blind observers. *Perception* 39(9):1261–1275.
6. Fernandes, A. M., & Albuquerque, P. B. (2012). Tactile perception: A review of experimental variables and procedures. *Cognitive Processing*, 13(4), 285–301.
7. Saal, H. P., & Bensmaia, S. J. (2014). Touch is a team effort: Interplay of submodalities in cutaneous sensibility. *Trends in Neurosciences*, 37(12), 689–697.
8. Cappagli, G., Cocchi, E., & Gori, M. (2017). Auditory and proprioceptive spatial impairments in blind children and adults. *Developmental Science*, 20(3).
9. Ocellli, V., Bruns, P., Zampini, M., & Röder, B. (2012). Audiotactile integration is reduced in congenital blindness in a spatial ventriloquism task. *Neuropsychologia*, 50(1), 36–43.

## Acknowledgments

We thank Elisa Freddi for assistance in collecting the data. This project has been funded by the European Union's Horizon 2020 Research and Innovation Programme under Grant Agreement No 860114