



## Identification of two distinct OH infrared signatures in pyroxenes from pyroxenite xenoliths

Jannick Ingrin (1), Samantha Azevedo-Vannson (2), and Lydéric France (2)

(1) Univ.Lille, CNRS, INRA, ENSCL, UMR 8207, UMET, Unité Matériaux et Transformations, F 59 000 Lille, France. jannick.ingrin@univ-lille.fr, (2) CRPG, UMR 7358, CNRS, Université de Lorraine, Vandoeuvre-lès-Nancy, France

Water concentration in pyroxenes from mantle xenoliths is generally used to trace the water content of lithospheric mantle. However, the infrared OH signatures recorded by pyroxenes provides also important information on their previous equilibrium. We present results of infrared analyses of garnet pyroxenites xenoliths, two from the French Massif Central (Devès, FMC) and two from Cameroon (Youkou Maar, Adamaou, Africa). The FMC pyroxenites have low water content and present dominant OH bands at 3595, 3517  $\text{cm}^{-1}$  and 3571, 3517  $\text{cm}^{-1}$  for cpx and opx, respectively. The pyroxenites from Cameroon have high water content and have dominant bands at 3635, 3445  $\text{cm}^{-1}$  and 3602, 3417  $\text{cm}^{-1}$  for cpx and opx. Like for OH spectra from peridotite xenoliths from Nógrád-Gömör (Hungary), it was possible to isolate two distinct end-member spectra in pyroxenes (Patkó et al. 2019). Spectra signatures of cpx and opx are correlated, showing that they have both recorded the same events. The two end-members spectra clearly point out for two distinct spectral signatures related to distinct OH defects configurations.

The pyroxenites from FMC have crystallized at higher temperature than those of Cameroon (1050°C versus 950°C; France et al. 2015). These results suggest that the FMC pyroxenites have also crystallized in drier conditions than the ones from Cameroon or have experienced dry metasomatism after their crystallization.

### References

- France, L. et al. (2015) Mantle refertilization and magmatism in old orogenic regions: The role of late-orogenic pyroxenites. *Lithos*, 232, 49-75.
- Patkó L. et al. (2019) Extremely low structural hydroxyl contents in upper mantle xenoliths from the Nógrád-Gömör volcanic field (northern Pannonian Basin): Geodynamic implications and the role of post-eruptive re-equilibration. *Chemical Geology*, <https://doi.org/10.1016/j.chemgeo.2018.12.017>.