



HAL
open science

Power Is More Relevant Than Ascensional Speed to Determine Metabolic Demand at Different Gradient Slopes During Running

Corentin Hingrand, Nicolas Olivier, Adrien Combes, Samir Bensaid, Frédéric Daussin

► **To cite this version:**

Corentin Hingrand, Nicolas Olivier, Adrien Combes, Samir Bensaid, Frédéric Daussin. Power Is More Relevant Than Ascensional Speed to Determine Metabolic Demand at Different Gradient Slopes During Running. *Journal of Strength and Conditioning Research*, 2023, *Journal of Strength and Conditioning Research*, 37 (11), pp.2298-2301. 10.1519/jsc.0000000000004598 . hal-04274690

HAL Id: hal-04274690

<https://hal.univ-lille.fr/hal-04274690>

Submitted on 8 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Power is more relevant than ascensional speed to determine metabolic demand at different gradient slopes during running

Corentin Hingrand¹, Nicolas Olivier², Adrien Combes², Samir Bensaid², Frédéric N. Daussin²

Running head: Trail training parameter

Affiliations:

¹ U1075, COMETE, Université de Caen

² Univ. Lille, Univ. Artois, Univ. Littoral Côte d'Opale, ULR 7369 - URePSSS - Unité de Recherche Pluridisciplinaire Sport Santé Société, F-59000 Lille, France

Corresponding author: Frédéric N. Daussin, Ph.D., Faculté des Sciences du Sport, 9 rue de l'Université, F-59790 Ronchin, France; E-mail: frederic.daussin@univ-lille.fr; Phone: +33 (0)3.20.88.73.69; Fax: +33 (0)3.20.88.73.63

Funding: No funding to disclose

ABSTRACT

Trail running is characterised by successive uphill and downhill running sessions. In order to prescribe training intensity, an assessment of maximal running capacity is required. This study compared two uphill incremental tests using the same ascensional speed increment to identify the influence of the slope gradient on performance. Ten subjects (eight men and two women) performed three incremental exercises on various slope (1%: IT01, 10%: IT10, and 25%: IT25), the ascensional speed increment was similar between IT10 and IT25 (100 m/h every minute). Gas exchanges, heart rate, and power were monitored continuously during the tests. Similar VO_{2max} levels were observed in the three conditions: 68.7 ± 6.2 for IT01, 70.1 ± 7.3 for IT10, and 67.6 ± 7.0 for IT25. A greater maximal ascensional speed was reached in the IT25 (1760 ± 190 vs 1330 ± 106 for IT25 and IT 10 respectively, $p < 0.01$). A significant relationship was observed between relative VO_2 levels and relative power without any effect of slope. Power should be the parameter used for prescribing training intensity compared to ascensional speed in trail.

Keywords: VO_{2max} , slope, running, performance

INTRODUCTION

Trail running (TR) is defined by the International Trail Running Association as “a pedestrian and off-road race conducted in a natural environment (e.g. mountain) with minimal possible paved or asphalt road (<20% of the total duration race)”. TR is challenging, as races are characterised by successive uphill and downhill running sections that are associated with major physiological changes (14). Using the classic endurance running model, maximal oxygen consumption (VO_{2max}) has been shown to account for 90% of TR performance (11).

In order to prescribe adequate exercise intensity, coaches use a test protocol to establish performance levels, and plan training accordingly. The majority of exercise tests are performed by running on a horizontal grade treadmill. Recently, the comparison of an outdoor 16% uphill exercise test has been compared to 1% laboratory test. This suggested that metabolic parameter values differ despite similar VO_{2max} values (22). Moreover, new tests have been designed to evaluate TR athletes. Scheer et al. (21) proposed a trail test with a continuous increase of both speed (+0.5 km/h per minute) and inclination (1% per minute). The athletes reached higher VO_{2max} values during the trail test compared to the horizontal test, suggesting that the use of graded inclinations during testing should be employed to match to the specificity of TR. However, this test limits the ability to use the results to prescribe training. Doucende et al. (10) developed a specific incremental trail test with increment expressed in metres per hour. The test was performed on a 25% slope, and the speed increment was designed to increase ascending speed (AS) by 100 m/h every minute. In others tests, different constant slopes (from 10 to 30%) have been used, and allow the determination of sub-maximal and maximal aptitude in TR racers

(4,9,11). Whether ascending speed may be used by coaches to prescribe training has not yet been determined.

The aim of the present study was to compare incremental tests using an identical ascensional speed increment (100 m/h) at different slopes to determine whether AS or power better reflects the metabolic cost of running at different slope gradients. We hypothesised that the maximal AS would vary with the slope and would be a reference parameter to prescribe training intensity.

METHODS

Experimental Approach to the Problem

Subjects completed three laboratory tests in this randomised, cross-over study. All visits were separated by at least 48 hours, over a 10-day period. Subjects performed three maximal graded tests on a treadmill to determine VO_{2max} . Power and O_2 consumption were monitored to assess the relationship between AS and power, or O_2 consumption.

Subjects

For this study, 10 subjects were recruited (female $n=2$ and male $n=8$), who were trail runners, (mean \pm SD: age 27.9 ± 8.74 years; height 175 ± 9.3 cm; body mass 65.3 ± 12.54 kg; and fat mass $10.9 \pm 3.99\%$) and considered to be strong to advanced athletes by the ITRA performance index (index: 542-781; (26)). All subjects were healthy and free of any history of metabolic, cardiovascular, hormonal, and respiratory diseases, or musculoskeletal disorders. None of the subjects were taking any medications or supplements, and they were advised to maintain their

normal dietary intake throughout the study. All subjects provided written informed consent after ethical approval was obtained for the Ethics Committee of the University of Lille (2019-381-S77).

Procedures

All subjects completed the three different exercise test protocols on three different occasions. All subjects were familiar with treadmill running, and performed the tests with their usual training shoes. Before all testing, subjects refrained from severe physical activity for at least 48 hours.

A standardised warm-up programme of 15 minutes preceded each test. The three tests were performed on a motorised treadmill (Intelligent Suspension 3, Cybex International Inc, MA, USA) as follows: (1) incremental test (IT01) starting at 8km/h with a progressive increase of 1km/h every minute and an inclination of 1%; (2) incremental test (IT10) starting at an AS of 500 m/h with a progressive increase of 100 m/h every minute with an inclination of 10%; and (3) incremental test (IT25) starting at an AS of 500 m/h with a progressive increase of 100 m/h every minute with an inclination of 25%.

During all tests, gas exchanges were collected breath by breath, together with heart rate, by a Metamax 3B-R2 system (Biophysik GmbH, Leipzig, Germany). VO_{2max} was defined as the highest 30-s average VO_2 value. The mean value over the last 20-s of each step was used to determine the relationship between AS and VO_2 .

Power was monitored using a power meter (Stryd Powermeter; Stryd Inc., Boulder, CO, USA), which is a foot-mounted inertia sensor of 9.1 g reinforced with carbon fibre that estimates power in watts. Based on 6-axis inertial motion sensor with a 1 Hz sampling frequency, the device provides several metrics to characterize performance: pace, elevation, distance, cadence, ground contact time, vertical oscillation, leg stiffness, running power, form power. The device has been validated previously at different running velocities (13,16). The power meter was firmly attached to the shoe according to the manufacturer recommendations. Subjects provided their body mass prior to its use; this was a requisite for the power estimation. The data extracted from the Stryd power meter were timed averages of 10-s, and the mean value over the last 20-s of each step was used to determine the relationship between AS and power.

At the end of each test, participants were asked to provide perceived exertion on the 6–20 point Borg scale (5).

Statistical Analyses

Descriptive statistics are represented as mean \pm SD. Tests of normal distribution and homogeneity (Shapiro–Wilk and Levene’s test, respectively) were performed on all data before analysis. Pairwise comparisons of mean (t-test) were conducted between the conditions at the different AS over the incremental test. A One-way ANOVA for repeated measure was performed to compare variables between the three conditions and a Tukey post-hoc test was used to identify significant difference between protocols. Relationships between variables were analysed using a Pearson’s product–moment correlation. Correlation coefficients were evaluated using the

following criteria: small = 0.10–0.29, moderate = 0.30–0.49, large = 0.50–0.69, very large = 0.70–0.89, nearly perfect = 0.90–0.99 (15). Data analysis was performed using Prism 9.1 (GraphPad Software, Inc., San Diego, CA). The magnitude of the differences between values was also interpreted using the Cohen's *d* effect size (ES)(7). Effect sizes of <0.20, 0.20–0.50, 0.50–0.80, and > 0.80 were considered as trivial, small, moderate, and large, respectively. The level of significance used was set a $p < 0.05$.

RESULTS

Table 1 summarises the mean maximal values of the three exercise test protocols. No difference was observed between the protocols for $\text{VO}_{2\text{max}}$, heart rate, power, and ratings of perceived exertion. A higher maximal velocity was observed in IT01 vs. IT10 and IT25 (ES = 3.96 and 8.76, respectively, $p < 0.001$), and maximal velocity was higher in IT10 compared to IT25 (ES = 6.55, $p < 0.001$). A greater AS was reached in the IT25 vs IT10 (ES = 2.79, $p < 0.001$).

--- INSERT TABLE 1 HERE ---

A significant relationship between AS and VO_2 was observed in IT10 and IT25 ($r^2=0.991$, $p < 0.001$, and $r^2=0.997$, $p < 0.001$, respectively, Figure 1A), and a significant difference between the slopes of the linear regression was observed ($P < 0.001$). Similarly, a significant relationship between AS and power was observed in IT10 and IT25 ($r^2=0.992$, $p < 0.001$, and $r^2=0.965$, $p < 0.001$, respectively,

Figure 1B), and a significant difference between the slopes of the linear regression was observed ($P<0.001$). A significant relationship between relative VO_2 and relative power was observed in IT10 and IT25 ($r^2=0.895$, $p<0.001$, and $r^2=0.852$, $p<0.001$, respectively, Figure 1C), and no difference was observed between the slopes ($P=0.38$).

--- INSERT FIGURE 1 HERE ---

DISCUSSION

The goal of this study was to determine whether the slope gradient influenced the peak AS in order to determine the utility of peak AS to prescribe training programs. Our results demonstrate that the slope gradient influence the peak AS but not maximal cardiometabolic values. The relationships between AS and VO_2 and AS and power were different across the slopes. The power is related to VO_2 independently of the slope gradient.

The peak AS reached by athletes was different upon the slope gradient. However, they reached similar $\text{VO}_{2\text{max}}$ in the three conditions. In the literature, the results of the influence of the slope on $\text{VO}_{2\text{max}}$ are contradictory: some studies observed an influence of slope (21,25) while others did not (3,8,17). The level of the population may explain these discrepancies as differences in $\text{VO}_{2\text{max}}$ became smaller when investigation trained subject (8,22). In our study, the subjects were considered to be trained, which may mitigate $\text{VO}_{2\text{max}}$ differences across exercise protocols. Therefore, these results suggest that all the protocols are appropriate for $\text{VO}_{2\text{max}}$ assessment for

trained trail runners. The protocol of test that was used to determine VO_{2max} may explain the discrepancies across the studies (23). In our study, we used a constant increment (100m/h) while Scheer et al.'s study (21) used incremental increments (107 m/h to 217 m/h). Therefore, the difference of VO_{2max} might be explained by the different increments in workload over time. An underestimation of VO_{2max} would be expected due to steeper increases of metabolic requirement (4).

We observed a linear relationship between VO_2 and AS, and between power and AS, and greater metabolic and power values were observed on a 25% slope vs a 10% slope at a same absolute AS. These results are interesting for training purposes while considering AS as an intensity index during training sessions. In line, previous studies reported a linear increase in energy cost with each increment (3,19,24). The greater energy cost at a same absolute AS may result from the higher muscular mass involved while increasing the sloped gradient (22, 50, 52). The uphill running locomotion is associated with stronger muscular contraction and mechanical work (52). Moreover, for a same AS, the horizontal speed will be higher when the slope increment is lower suggesting that horizontal speed may be a potent explanation for the greater metabolic cost observed at low gradient slope. All together these data indicate that for a same absolute AS the increase of the slope gradient modifies running speed and running patterns which alter running economy and induce an increase of net energy generation.

The linear relationship between oxygen uptake and work rate has been widely described (12,20). Previous studies assessed the relationship between power and oxygen uptake, and all found a

linear and positive relationship between the two parameters (1,2,16,18). Our results are in line with the literature where there is a strong and positive relationship between power and oxygen uptake. As previously suggested, we pay attention to the methodology employed to analyse VO_2 and Stryd power data (16). We used the last 20 seconds of the step to avoid the time delay for VO_2 to increase while power increases instantaneously. Our results also suggest that the power assessed using the Stryd was unaffected by the gradient of the slope, and should be an interesting tool to monitor exercise intensity during trail running.

PRACTICAL APPLICATIONS

The present study demonstrated that the use of peak AS should not be used by coaches to prescribe or individualise training programmes as the cardiometabolic responses were slope-dependent. However, AS would be a relevant parameter when coaches want to individualise uphill training sessions as long as the slope gradient remained similar. In that situation and considering that the slope gradient influences the cardiometabolic responses, the coaches should choose the AS according to the cardiometabolic target of the training session.

The use of a power meter could be an interesting tool to prescribe and individualise training intensity. In fact, the relationship between O_2 consumption and power were correlated and similar between the two-slope gradient explored in this study. These results suggest that power could be valid for different terrains as it was the case during trail running. Despite the growing offer of power meters, specific attention should be paid to the validity of the devices used.

ACKNOWLEDGEMENTS

The authors thank all the trail runners for their enthusiastic participation in the study. The authors have no conflict of interest to declare. The study was financed by the authors' own institutional resources. The results of the present study do not constitute endorsements of the products by the authors or the NSCA.

REFERENCES

1. Aubry, RL, Power, GA, and Burr, JF. An Assessment of Running Power as a Training Metric for Elite and Recreational Runners. *J Strength Cond Res* 32: 2258–2264, 2018.
2. Austin, CL, Hokanson, JF, McGinnis, PM, and Patrick, S. The Relationship between Running Power and Running Economy in Well-Trained Distance Runners. *Sports (Basel)* 6: 142, 2018.
3. Balducci, P, Cléménçon, M, Morel, B, Quiniou, G, Saboul, D, and Hautier, CA. Comparison of Level and Graded Treadmill Tests to Evaluate Endurance Mountain Runners. *J Sports Sci Med* 15: 239–246, 2016.
4. Beltz, NM, Gibson, AL, Janot, JM, Kravitz, L, Mermier, CM, and Dalleck, LC. Graded Exercise Testing Protocols for the Determination of VO₂max: Historical Perspectives, Progress, and Future Considerations. *J Sports Med (Hindawi Publ Corp)* 2016, 2016. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5221270/>
5. Borg, G. Perceived exertion as an indicator of somatic stress. *Scandinavian Journal of Rehabilitation Medicine* 2: 92–98, 1970.
6. Cassirame, J, Godin, A, Chamoux, M, Doucende, G, and Mourot, L. Physiological Implication of Slope Gradient during Incremental Running Test. *International Journal of Environmental Research and Public Health* 19: 12210, 2022.
7. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. New York: Routledge, 1988.
8. Davies, CTM, Sargeant, AJ, and Smith, B. The physiological responses to running downhill. *Europ J Appl Physiol* 32: 187–194, 1974.
9. Doucende, G, Risetto, C, Defer, T, Mourot, L, and Cassirame, J. Field Adaptation of a Specific Trail Running Incremental Test: IncremenTrail. 2018.
10. Doucende, G, Risetto, C, Mourot, L, and Cassirame, J. Biomechanical adaptation of

preferred transition speed during an incremental test in a gradient slope. *Computer Methods in Biomechanics and Biomedical Engineering* 20: 69–70, 2017.

11. Ehrström, S, Tartaruga, MP, Easthope, CS, Brisswalter, J, Morin, J-B, and Vercruyssen, F. Short Trail Running Race: Beyond the Classic Model for Endurance Running Performance. *Med Sci Sports Exerc* 50: 580–588, 2018.
12. Gaesser, GA and Poole, DC. The slow component of oxygen uptake kinetics in humans. *Exerc Sport Sci Rev* 24: 35–71, 1996.
13. García-Pinillos, F, Roche-Seruendo, LE, Marcén-Cinca, N, Marco-Contreras, LA, and Latorre-Román, PA. Absolute Reliability and Concurrent Validity of the Stryd System for the Assessment of Running Stride Kinematics at Different Velocities. *J Strength Cond Res* 35: 78–84, 2021.
14. Giandolini, M. Fatigue associated with prolonged graded running. *Eur J Appl Physiol* 15, 2016.
15. Hoffman, J. Norms for Fitness, Performance, and Health. Human Kinetics, 2006.
16. Imbach, F, Candau, R, Chailan, R, and Perrey, S. Validity of the Stryd Power Meter in Measuring Running Parameters at Submaximal Speeds. *Sports (Basel)* 8, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7404478/>
17. Kasch, FW, Wallace, JP, Huhn, RR, Krogh, LA, and Hurl, PM. VO₂max during horizontal and inclined treadmill running. *J Appl Physiol* 40: 982–983, 1976.
18. Lara, F, Shearer, L, Coppi, M, Hayden, N, Ogden, J, Murr, S, et al. Reliability Of A Running Power Meter Between Trials Of Submaximal Running On Three Different Surfaces. *Medicine & Science in Sports & Exercise* 50: 436, 2018.
19. Padulo, J, Powell, D, Milia, R, and Ardigò, LP. A Paradigm of Uphill Running. *PLoS One* 8: e69006, 2013.
20. Poole, DC, Barstow, TJ, Gaesser, GA, Willis, WT, and Whipp, BJ. VO₂ slow component: physiological and functional significance. *Med Sci Sports Exerc* 26: 1354–1358, 1994.
21. Scheer, V, Ramme, K, Reinsberger, C, and Heitkamp, H-C. VO₂max Testing in Trail Runners: Is There a Specific Exercise Test Protocol? *Int J Sports Med* 39: 456–461, 2018.
22. Schöffl, I, Jasinski, D, Ehrlich, B, Dittrich, S, and Schöffl, V. Outdoor Uphill Exercise Testing for Trail Runners, a More Suitable Method? *J Hum Kinet* 79: 123–133, 2021.
23. Schöffl, V, Pöppelmeier, O, Emmler, J, Schöffl, I, Küpper, T, and Lutter, C. Ski Mountaineering - Evaluation of a Sports Specific Performance Diagnosis. *Sportverletz Sportschaden* 32: 233–242, 2018.
24. Snyder, KL and Farley, CT. Energetically optimal stride frequency in running: the effects of incline and decline. *J Exp Biol* 214: 2089–2095, 2011.
25. Taylor, HL, Buskirk, E, and Henschel, A. Maximal Oxygen Intake as an Objective Measure of Cardio-Respiratory Performance. *Journal of Applied Physiology* 8: 73–80, 1955.
26. All about the ITRA performance index - ITRA. Available from: <https://itra.run/content/indice-performance>

TABLE**Table 1: The mean maximal values of the three exercise test protocols**

Parameters	IT01	IT10	IT25
VO _{2max} (mL·min ⁻¹ ·kg ⁻¹)	68.7 ± 6.2 [64.3–73.1]	70.1 ± 7.3 [64.9–75.3]	67.6 ± 7.0 [62.6–72.6]
Maximal velocity (km·h ⁻¹)	19.2 ± 1.8 [18.0–20.5]	13.3 ± 1.1 [12.5–14.1] ^a	7.0 ± 0.8 [6.5–7.6] ^{a,b}
Ascensional speed (m·h ⁻¹)		1330 ± 106 [1254–1406]	1760 ± 190 [1624–1896] ^b
Heart rate (bpm)	185 ± 11 [177–193]	182 ± 11 [174–190]	183 ± 14 [173–193]
Power (W·kg ⁻¹)	4.9 ± 1.1 [4.1–5.7]	5.4 ± 0.7 [4.9–6.0]	4.9 ± 1.4 [3.9–6.0]
Ratings of perceived exertion	16.7 ± 1.1 [15.9–17.5]	16.8 ± 1.7 [15.6–18.0]	17.3 ± 1.2 [16.5–18.1]

Mean values ± SD [95% CI], ^a*p*<0.001 vs. IT01, ^b*p*<0.001 vs. IT10

FIGURE

Figure 1: Relationships between A. oxygen consumption and ascensional velocity, B. power and ascensional velocity, and C. oxygen consumption and power. Oxygen consumption was expressed in $\text{mL}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$, and power was expressed in $\text{W}\cdot\text{kg}^{-1}$. This represents pooled data from 10 subjects.

