



HAL
open science

Original bar fixation technique in minimally invasive repair of pectus excavatum in adolescents: A 36-case series.

Adrien Fournier, Damien Fron, Michel Bonnevalle, Bernard Herbaux, Aurelie Mezel, Eric Nectoux, A Fournier

► To cite this version:

Adrien Fournier, Damien Fron, Michel Bonnevalle, Bernard Herbaux, Aurelie Mezel, et al.. Original bar fixation technique in minimally invasive repair of pectus excavatum in adolescents: A 36-case series.. Orthopaedics & Traumatology: Surgery & Research, 2020, Orthopaedics & Traumatology: Surgery & Research, 106 (1), pp.155-157. 10.1016/j.otsr.2019.10.007 . hal-04356220

HAL Id: hal-04356220

<https://hal.univ-lille.fr/hal-04356220v1>

Submitted on 20 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Technical note

Original bar fixation technique in minimally invasive repair of pectus excavatum in adolescents: a 36-case series.

Adrien **Fournier**^a, Damien **Fron**^a, Michel **Bonnevalle**^b,

Bernard **Herbaux**^a, Aurélie **Mezel**^a, Eric **Nectoux**^{a,*}

a Equipe de Chirurgie Orthopédique Infantile, Service de Chirurgie et Orthopédie de l'Enfant, Hôpital Jeanne de Flandre, Centre Hospitalier Régional Universitaire de Lille, 2 Avenue Oscar Lambret, 59000 Lille, France

b Equipe de Chirurgie Viscérale Pédiatrique, Service de Chirurgie et Orthopédie de l'Enfant, Hôpital Jeanne de Flandre, Centre Hospitalier Régional Universitaire de Lille, 2 Avenue Oscar Lambret, 59000 Lille, France

* Corresponding author: Eric Nectoux, Service de Chirurgie et Orthopédie de l'Enfant, Hôpital Jeanne de Flandre, Centre Hospitalier Régional Universitaire de Lille, 2 Avenue Oscar Lambret, 59000 Lille, France

E-mail: eric.nectoux@chru-lille.fr (E. Nectoux).

Cell: +33 (0)6 66 22 90 58 (E. Nectoux).

Abstract

In pectus excavatum, minimally invasive endoscopic repair is the current gold standard in adolescents. The main postoperative pitfall is bar displacement, making fixation a major issue. We report an original transcostal technique using absorbable USP-2 suture in 36 patients.

There was a single case of bar displacement (2.78%; 95% CI, 0.07-14.53).

Bar stability was comparable to literature reports; but the present technique avoided the complications potentially incurred by use of metal wire, pericostal fixation points and stabilizers.

Key-words: Pectus excavatum, Nuss procedure, minimally invasive surgery, bar fixation.

1. Introduction

Pectus excavatum mainly affects males [1] and is usually present from birth. The deformity persists or worsens during rapid thorax growth at puberty. Impact is mainly psychological, or more rarely cardiorespiratory. The Nuss procedure, also known as minimally invasive repair of pectus excavatum (MIRPE), is the gold standard in adolescence, providing lasting results [2,3]. It has become an attractive alternative to sternochondroplasty, with smaller lateral scars and conserved costal cartilage [4]. There are, however, complications: secondary bar displacement, with risk of recurrence and of mediastinal injury, and residual pain at the fixation points [2-5]. Kabbaj et al. [6] described costal bar fixation by pericostal wire, with a single case of traumatic displacement in a series of 70 patients. Bond et al. [7] described use of pericostal and subxyphoid absorbable suture, with only 2 cases of displacement in 73 patients. And Yoo et al. [8] reported zero displacement in 44 patients, using “sutureless” fixation with 2 bars connected by metal devices. The present technique extends the range of bar fixation options, using absorbable transcostal suture. It seems technically easier than pericostal suture and less expensive than use of stabilizers, and facilitates hardware removal.

2. Surgical technique

Bar placement follows the princeps technique described by Nuss et al. [3].

The configuration between the ribs and the emergence of the bar is analyzed, and an absorbable USP (United States Pharmacopeia) size 2 polyglactin 910 transcostal suture is performed using a large needle under the positioned bar, facing the notches (Medxpert, Eschbach, Germany) intended for stabilizer fixation (**Figures 1 and 2**). If possible, a second suture is made contralaterally. No drainage is used.

Movement and sport are restricted:

- no twist or lateral flexion of the trunk for 3 weeks;
- no individual sports activity for 6 weeks;
- no team sports for 3 months.

The bar is removed at 2 years.

3. Study population and results

Thirty-seven pectus excavatum patients operated on by this technique between January 2010 and December 2016 were included, 1 of whom was subsequently excluded for early bar ablation due to infection.

Table 1 shows results, and Table 2 complications. There was 1 case of neurologic deficit, in the left upper-limb C5 and C6 root territories due to bad sleep posture, in a patient receiving morphine derivatives. There was 1 case of bar displacement: 1/36, 2.78%; 95% CI, 0.07-14.53.

4. Discussion

Incidence of bar displacement was comparable to the 2.1% reported by Nuss and Kelly [5] for stabilizers associated to pericostal suture, within the present 95% confidence interval, suggesting no real difference.

The present technique has several advantages:

- Absorbable suture avoids the complications of metal wire breakage (subcutaneous irritation, pulmonary parenchyma lesion) and facilitates subsequent bar removal.
- Transcostal, unlike pericostal, suture avoids intercostal needle passage with

risk of neurovascular bundle lesion (neuropathic pain, dysesthesia, bleeding with hemopneumothorax) or pulmonary parenchyma lesion.

- The absence of stabilizer reduces local discomfort and scar disorder, reduces procedural cost and shortens operative time.

5. Conclusion

The present study demonstrated the efficacy and safety of this bar fixation technique over follow-up until removal: only 1 displacement was observed, out of 36 cases.

Acknowledgments

Thanks to Pr Jean-Luc Jouve and Pr Henri Migaud for kindly re-editing the manuscript.

Thanks to Mr Martinet, medical illustrator, for his contribution to the illustrations.

Funding

The University of Lille 2 – Law and Health medical faculty provided funding for the illustrations.

Author contributions

Eric Nectoux: study concept and design, re-editing.

Adrien Fournier: data collection, analysis and interpretation, article writing.

Damien Fron, Michel Bonnevalle, Bernard Herbauts, Aurélie Mézel: re-editing and advice.

Conflicts of interest

The authors have no conflicts of interest to disclose.

References

- [1] Faglin P, Nectoux É, Belkhou A, Guerreschi P, Duquennoy-Martinot V. The unsightly chest: Analysis and anomaly. Curative or palliative approach? *Ann Chir Plast Esthet* 2016;61:680–93.
- [2] Felts E, Jouve JL, Blondel B, Launay F, Lacroix F, Bollini G. Child pectus excavatum: Correction by minimally invasive surgery. *Orthop Traumatol Surg Res* 2009;95:190-5.
- [3] Nuss D, Kelly RE, Croitoru DP, Katz ME. A 10-year review of a minimally invasive technique for the correction of pectus excavatum. *J Pediatr Surg* 1998;33:545-52.
- [4] Wurtz A, Rouse N, Benhamed L, Conti M, Hysi I, Pinçon C, et al. Simplified open repair for anterior chest wall deformities. Analysis of results in 205 patients. *Orthop Traumatol Surg Res* 2012;98:319-26.
- [5] Nuss D, Kelly RE. Indications and technique of Nuss procedure for pectus excavatum. *Thor Surg Clin* 2010;20:583-97.
- [6] Kabbaj R, Burnier M, Kohler R, Loucheur N, Dubois R, Jouve J-L. Minimally invasive repair of pectus excavatum using the Nuss technique in children and adolescents: indications, outcomes, and limitations. *Orthop Traumatol Surg Res* 2014;100:625–30.
- [7] Bond SJ, Rapstine E, Bond JM. Results of pectus excavatum correction using a minimally invasive approach with subxyphoid incision and three-point fixation. *Pediatr Surg Int* 2018;34:75-78.
- [8] Yoo G, Rha EY, Jeong JY, Lee J, Sim SB, Jo KH. Emerging Fixation Technique to Prevent Pectus Bar Displacement: Needlescope-Assisted 3-Point Fixation. *Thorac Cardiovasc Surg* 2016;64:78–82.

Figure legends

Figure 1: Bar fixation by absorbable transcostal suture. Left: cross-sectional view; bar in front, rib behind.

Figure 2: Intraoperative photographs. 2A: Transcostal passage of suture, positioning the two ends in the notches on either side of the bar. 2B: Tying to bar.

Table 1: Patient data (n=36).



Male n(%)	31 (86.1)
Age at first consultation (years)	14.7 ± 2.2 (8.0 18.2)
Reason for consultation n(%)	
<i>Esthetic</i>	27 (75.0)
<i>Intolerance of effort</i>	7 (19.4)
<i>Pain</i>	3 (8.3)
<i>Feeding issues</i>	1 (2.8)
Symmetric pectus excavatum n(%)	28 (77.8)
Associated spinal deformity n(%)	8 (22.2)
Depth of excavatum (cm)	3.2 ± 0.9 (1.0 5.0)
Age at surgery (years)	15.7 ± 1.3 (13.0 18.7)
Preoperative Haller index	3.7 ± 0.8 (2.8 5.7)
Hospital stay (days)	7.4 ± 1.2 (5.0 10.0)
Operative time (min)	61.1 ± 14.7 (39.0 117.0)
Bar in place (years)	2.1 ± 0.2 (1.6 2.9)

Qualitative variables reported as number (percentage), and quantitative variables as mean ± standard deviation (range).

Table 2: Postoperative complications (n = 36)

Bar displacement	1 (2.78)
Thoracic drainage	2 (5.56)
Neuropathic pain	4 (11.1)
Hardware discomfort	0 (0.0)
Pneumopathy	1 (2.78)
Atelectasis	2 (5.56)
Metal allergy	1 (2.78)
Neurologic deficit	1 (2.78)

Data reported as number (percentage).



MARTINET
pour A. Fournier

