



HAL
open science

The Complexity of Aqueous Alteration Veins in Nakhlites

Lisa Krämer Ruggiu, Lydie Bonal, Laurette Piani, Hugues Leroux, Olivier
Grauby, J. Gattacceca, Bertrand Devouard

► **To cite this version:**

Lisa Krämer Ruggiu, Lydie Bonal, Laurette Piani, Hugues Leroux, Olivier Grauby, et al.. The Complexity of Aqueous Alteration Veins in Nakhlites. 85th Annual Meeting of The Meteoritical Society, Aug 2022, Glasgow, Scotland, United Kingdom. hal-04447461

HAL Id: hal-04447461

<https://hal.univ-lille.fr/hal-04447461v1>

Submitted on 8 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE COMPLEXITY OF AQUEOUS ALTERATION VEINS IN NAKHLITES

L. Krämer Ruggiu^{1,2}, B. Devouard², J. Gattacceca², L. Bonal³, L. Piani⁴, H. Leroux⁵, O. Grauby⁶

¹Analytical-, Environmental- and Geo-Chemistry, Vrije Universiteit Brussel, Brussels, Belgium (lisa.kramer.ruggiu@vub.be), ²Aix Marseille Univ, CNRS, IRD, INRAE, CEREGE, Aix-en-Provence, France, ³Univ. Grenoble Alpes, CNRS, IPAG, Grenoble, France 11, ⁴CRPG, CNRS, Université de Lorraine, UMR 7358, Vandoeuvre-les-Nancy, France, ⁵Univ. Lille, CNRS, INRAE, Centrale Lille, UMR 8207 - UMET, F-59000, Lille, France, ⁶CINaM, Aix-Marseille Université, Campus de Luminy, 13288 Marseille, France

Introduction: Nakhilites are a group of basaltic meteorites originating from the mid-Amazonian igneous crust of Mars [1]. They represent an interesting set of samples to study the aqueous alteration at the Martian subsurface, as they contain aqueous alteration phases in olivine called “iddingsite” (e.g. [2]). Studying iddingsite in nakhilites can reveal the water-crust interactions and the hydrothermal physico-chemical conditions that prevailed at the Martian subsurface, with implications for the past habitability of Mars, and in particular for microbial life during the Amazonian period. We studied iddingsite in Caleta el Cobre (CeC) 022, a nakhilite that contains a high amount of alteration products, and compared this meteorite to 7 other nakhilites.

Methods: A serie of analyses were conducted on eight nakhilites polished sections: CeC 022, Governador Valadares, Lafayette, Miller Range (MIL) 03346, Nakhla, North-West Africa (NWA) 817, NWA 998 and Yamato (Y) 000953 to assess the mineralogy, the chemical and the isotopic composition of the alteration products in nakhilites: electronic microscopy (FEG-SEM and TEM), EDX microanalyses and chemical maps, LA-ICPMS major elements analyses, Raman spectra, and finally, hydrogen isotopic measurements by SIMS.

Results: In all nakhilites, the alteration veins show at least two types of iddingsite (Fig. 1A): (i) a coarse iddingsite in contact with olivine at the border of the alteration veins, showing 50-200 nm crystals, composed of a mixture of 1:1 phyllosilicates of greenalite-cronstedtite composition, and Fe-oxyhydroxides ; (ii) a fine iddingsite, in the inner part of the alteration veins, with <10 nm crystals, with a composition close to saponite (2:1 phyllosilicate). Complex chemical zoning of Mg, Ca, Mn, S, P and Al, has been observed on EDS-TEM mappings in NWA 10153. In most nakhilites, olivine grains also display planes of secondary inclusions, composed of pyroxene, magnetite and a void potentially filled by a fluid (Fig. 1B). In most nakhilites, magnetite-pyroxene symplectites are found in olivine grains, often at the border of the secondary inclusion planes. Those secondary inclusions and symplectites can also be observed at the center of the iddingsite veins (Fig. 1A). Also, sulfide-magnetite veinlets are observed at the center of the iddingsite veins, and also crosscutting olivine and pyroxene grains and mesostasis. Finally, organic matter is observed on Raman spectra in the iddingsite of many nakhilites, and is located in coarse iddingsite, as shown by TEM observations of NWA 10153.

Discussion and conclusion: Our favored scenario is that the secondary inclusions and the symplectites are formed by a first fluid alteration event triggered by late magmatic fluids, prior to the iddingsite formation event,. These secondary inclusions represent weakness planes that facilitate the circulation of the alteration fluid forming the iddingsite inside the olivine grains. The composition and texture of both types of iddingsite is suggestive of a crystallization by filling of existing fractures, with an alteration fluid enriched in elements from basaltic glass and host olivine dissolution, and changes in the alteration conditions or fluid composition. The sulfide-magnetite veinlets represent either the late stage of the same alteration event as the iddingsite formation, or a different later fluid injection. With the two types of iddingsite, the complex centers of veins and the chemical zoning in the fine iddingsite, we suggest that fluid alteration in the nakhilites has a complex multistage history of fluid injections.

References: [1] Nyquist, L. E., et al. (2001) *Chronology and evolution of Mars*. 105-164, [2] Changela, H. G., and J. C. Bridges. (2010) *Meteoritics & Planetary Science* 45 .12:1847-1867.




Figure 1: BSE images in olivine of CeC 022. A: iddingsite veins with magnetite-pyroxene symplectites as center of vein. B: symplectites of magnetite and pyroxene, and secondary inclusions composed of magnetite, pyroxene and a void. Idd c: coarse iddingsite; Idd f: fine iddingsite; Mt: magnetite; Ol: olivine; Px: pyroxene; Symp: symplectite.