



Make me more comfortable: effects of a hypnosis session on pain perception in chronic pain patients

David Ogez, Mathieu Landry, Rémi Caron-Trahan, Anne-Eva Jusseaux, Maryse Aubin, Jade Véronneau, Valentyn Fournier, Nadia Godin, Moulay Idrissi, Pierre Rainville, et al.

► To cite this version:

David Ogez, Mathieu Landry, Rémi Caron-Trahan, Anne-Eva Jusseaux, Maryse Aubin, et al.. Make me more comfortable: effects of a hypnosis session on pain perception in chronic pain patients. Frontiers in Psychology, 2024, Frontiers in Psychology, 15, 10.3389/fpsyg.2024.1362208 . hal-04549207

HAL Id: hal-04549207

<https://hal.univ-lille.fr/hal-04549207>

Submitted on 17 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



OPEN ACCESS

EDITED BY

Ernil Hansen,
University Medical Center Regensburg,
Germany

REVIEWED BY

Burkhard Peter,
University of Bern, Switzerland
Sonia Lorente,
Autonomous University of Barcelona, Spain

*CORRESPONDENCE

David Ogez
✉ david.ogez@umontreal.ca

RECEIVED 27 December 2023

ACCEPTED 12 February 2024

PUBLISHED 28 February 2024

CITATION

Ogez D, Landry M, Caron-Trahan R,
Jusseaux A-E, Aubin M, Véronneau J,
Fournier V, Godin N, Idrissi M, Rainville P and
Richebé P (2024) Make me more
comfortable: effects of a hypnosis session on
pain perception in chronic pain patients.
Front. Psychol. 15:1362208.
doi: 10.3389/fpsyg.2024.1362208

COPYRIGHT

© 2024 Ogez, Landry, Caron-Trahan,
Jusseaux, Aubin, Véronneau, Fournier, Godin,
Idrissi, Rainville and Richebé. This is an open-
access article distributed under the terms of
the [Creative Commons Attribution License](#)
(CC BY). The use, distribution or reproduction
in other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Make me more comfortable: effects of a hypnosis session on pain perception in chronic pain patients

David Ogez^{1,2*}, Mathieu Landry³, Rémi Caron-Trahan³,
Anne-Eva Jusseaux¹, Maryse Aubin¹, Jade Véronneau¹,
Valentyn Fournier⁴, Nadia Godin¹, Moulay Idrissi¹,
Pierre Rainville^{5,6} and Philippe Richebé^{1,2}

¹Research Centre, Hôpital Maisonneuve-Rosemont, Montréal, QC, Canada, ²Department of Anesthesiology and Pain Medicine, Université de Montréal, Montréal, QC, Canada, ³Département de Psychologie, Université de Montréal, Montréal, QC, Canada, ⁴SCALab – Sciences Cognitives et Sciences Affectives, Université de Lille, Lille, France, ⁵Faculty of Dental Medicine, Université de Montréal, Montréal, QC, Canada, ⁶Research Centre, Institut Universitaire de Gériatrie de Montréal, Montréal, QC, Canada

Introduction: Approximately one-quarter of Canadians experience chronic pain, a debilitating condition often necessitating opioid use, which raises concerns regarding dependency and overdose risks. As an alternative, we developed the HYLADO program (Hypnose de la Douleur, hypnosis of Pain in French), a novel self-hypnosis approach for chronic pain management. The development of this program followed the ORBIT model, a comprehensive framework for designing interventions encompassing several phases ranging from design to efficacy assessment.

Methods: In the present work, we conducted a preliminary evaluation of the HYLADO program with 21 participants (18 of the 21 patients were included in the analysis). The primary objective was to determine one session of the program's effectiveness in altering pain, anxiety and relaxation via pre-post analysis. The secondary goal was to examine the long-term effects across the same measures, in addition to the overall quality of life.

Results: The results highlight the benefits of our approach, while participants reported short-term significant pain reduction, decreased anxiety, and increased relaxation. Additionally, preliminary trends suggest improvements in physical activity and quality of life metrics.

Discussion: These positive outcomes highlight HYLADO's potential as an alternative to opioid therapy for chronic pain. Encouraged by these results, we aim to extend our research to a broader and more diverse cohort, paving the way for comprehensive randomized controlled trials. This expansion will further validate HYLADO's efficacy and its role in transforming chronic pain management.

KEYWORDS

self-hypnosis, chronic pain, anxiety, quality of life, intervention program, pre-post study

Introduction

In Canada, chronic pain is a prevalent issue, affecting more than 7 million individuals, or about one in four, over their lifetimes (Canadian pain task force, 2019). This condition not only impacts personal health but also imposes a considerable economic strain. The total cost, including both direct expenses such as medical services and indirect costs like lost productivity, is approximately \$40 billion (Canadian pain task force, 2019). At the level of individual, chronic pain significantly affects mental health, quality of life, and social inclusion among Canadians. Most notably, between 35 and 60% of those suffering from chronic pain is at an elevated risk of developing anxiety disorders (Choinière et al., 2020). Social consequences are also evident, with chronic pain contributing to early disability and impairing both personal and professional lives (Mills et al., 2019). In Canada, these social and health deficits are exacerbated by the lack of readily available services and delays in accessing specialized care (Canadian pain task force, 2019). The average wait time for pain management clinics is between 8 and 10 months, allowing ample time for pain to become chronic. These prolonged waits increase levels of pain and distress, while also reducing the likelihood of successful therapy. In parallel the lack of services, effective treatments, and lengthy wait times yield increased usage of potent analgesics, such as opioids. This heightened usage results in tolerance and dependence, limiting their availability in pain clinics in a timely manner (Mills et al., 2019). In sum, the current situation regarding chronic pain in Canada is difficult and is likely to get worse due aging population.

Following this adverse context, the Canadian government mandated a group of specialists, the Pain Task Force, to guide decision-makers in enhancing chronic pain prevention and management strategies (Canadian pain task force, 2020). A key recommendation from this panel was to increase the adoption of non-pharmacologic interventions, such as hypnosis, mindfulness, and acceptance therapy. This recommendation followed from strong empirical supporting their effectiveness in pain management. Furthermore, focused on cultivating pain self-management skills, these interventions present innovative solutions to address issues like healthcare accessibility, substance misuse, and the escalation of severe pain-related outcomes. Research consistently demonstrates that these strategies are effective in mitigating the risk of chronic pain persistence and its associated comorbidities. They integrate well within multimodal and biopsychosocial treatment frameworks, significantly benefiting patients' mental health and overall quality of life (Langlois et al., 2022).

Among these approaches, research indicates that clinical hypnosis represents an efficient non-pharmacological intervention for pain management in various clinical populations suffering from chronic pain (Langlois et al., 2022). Hypnotic interventions stand out for their ability to maintain its effects over an extended period of time based on a procedure that can simply reinstate suggestions for analgesia (Houzé et al., 2021). Moreover, in addition to pain reduction, evidence shows that clinical hypnosis can also reduce anxiety, improve sleep, and enhance quality of life of patients (Thompson et al., 2019). In summary, hypnosis represents a viable non-pharmacological treatment option for chronic pain (Jensen et al., 2006, 2020).

In contrast to hetero-hypnosis, which involves the guidance of a clinician, self-hypnosis is characterized by the patient's performance of hypnotic induction and suggestion procedures (Hammond, 2001). This approach promotes self-management of chronic pain outside therapeutic sessions with a healthcare professional (Langlois et al., 2022). Self-hypnosis relies on two key elements: instructions for

practicing self-hypnosis and audio recordings (Bicego et al., 2021; Eaton et al., 2021). Despite its apparent effectiveness, self-hypnosis training remains largely unexplored, with limited information in the literature about the optimal method for providing self-hypnosis instructions (Milling et al., 2021; Samami et al., 2021). The present work aims to address this lacuna by further developing a new standardized program specifically tailored for self-hypnosis training in chronic pain management.


Objectives

This research introduces a self-hypnosis training program, developed based on insights from prior chronic pain studies. Our approach to designing this program followed the ORBIT framework (Figure 1; Czajkowski et al., 2015). The ORBIT model represents a flexible overarching framework to guide and optimize the development of behavioral treatment across four distinct phases. During phase I, the program is conceptualized and refined, incorporating feedback and suggestions from potential users to ensure its relevance and efficacy. Phase II involves conducting preliminary studies to assess the program's initial impact and to set the stage for more extensive research. The subsequent phases, III and IV, are dedicated to rigorous efficacy and effectiveness studies, respectively. These phases are crucial for establishing the program's validity and determining its practical applicability in real-world settings. Based on this framework, the current study present work that was done during phase II. In this regard, the aim of the present work is to evaluate the effects of this program's hypnosis techniques in the context of chronic pain. Our evaluation is twofold: first, we aim to assess the immediate impact of a single hypnosis session on participants' levels of pain intensity, anxiety, and relaxation; second, our goal is also to examine the long-term benefits of ongoing self-hypnosis practice on the same measure and overall quality of life. We hypothesize that regular self-hypnosis will significantly improve the quality of life for these individuals, alongside marked reductions in anxiety and pain. This hypothesis is predicated on the notion that self-hypnosis can effectively modulate pain perception and bolster coping strategies, thus positively influencing both mental and physical health outcomes.

Materials and methods

HYlaDO intervention program

HYlaDO ("HYpnose pour la DOuleur"; hypnosis for pain in French) is a self-hypnosis program designed for improving the quality of life of chronic pain patients. As we mentioned previously, the design of our program tracks the stages of the ORBIT model. Our previous research documents the outcome of Phases Ia and Ib (Caron-Trahan et al., 2023a,b). The initial version of the HYlaDO program comprised eight sessions: a session introducing the participants, a session introducing hypnosis, 5 sessions offering 5 heterohypnosis exercises and recommendations for self-hypnosis practice, and a session for conclusion and feedback. The five exercises aimed at emotional release, acceptance, pain modification, pain reducing with magic glove, and confidence building. Each exercise followed a structured pattern, including a hypnotic induction procedure, a deepening phase, specific hypnotic work depending on the session objective,






TABLE 1 Descriptive data.

	N = 21	%
Sex		
Female	15	71
Male	6	29
Age (years)		
Mean (SD)	52 (11)	
Marital status		
Single	6	29
Divorced	4	19
Relationship	11	52
Having children		
Yes	17	81
No	4	19
Education		
Secondary	7	33
College	2	10
University	12	57
Professional status		
Unemployed	2	10
Temporary	3	14
Retired	3	14
Invalidity	5	24
Part-time work	1	5
Full-time work	7	33

pre- and post- hypnosis sessions at both T1 and T2, and between T1 and T2 for pre- and post-hypnosis time points, respectively. The difference in pain, anxiety, and relaxation scores between T1 and T2

were also compared. Also, total quality of life scores and sub-scores were compared between T1 and T2. Two-tailed Wilcoxon signed-rank test was used given the repeated measures design and the small sample size. Statistical significance threshold were settled at $\alpha=0.05$. All analyses were run using SPSS 28.0.1. software.


Results

As illustrated in the flowchart (Figure 3), 18 of the 32 participants successfully completed the measurements in our study. Dropouts were due to participants' unavailability, logistical challenges such as residing too far from the laboratory, and the worsening of physical health conditions.

Perceived pain, anxiety, and relaxation

For perceived pain and anxiety scores, non-parametric paired comparisons were conducted with Wilcoxon tests to compare the scores before and after the hypnosis sessions at T1 and T2. Results show a significant difference between the scores on the variables of interest before and after hypnosis session at T1 with lower scores of perceived pain [$W(18)=-3.366$; $p<0.001$] and anxiety [$W(18)=-2.955$; $p=0.003$], and higher score of relaxation [$W(18)=2.996$; $p=0.003$]. The same pattern is observed for measures before and after hypnosis at T2 with lower score of perceived pain [$W(17)=-3.415$; $p<0.001$] and anxiety [$W(17)=-3.18$; $p=0.001$], and higher scores of relaxation [$W(17)=3.638$; $p<0.001$].

No significant difference is observed between the scores of perceived pain before hypnosis session at T1 and T2 [$W(17)=0.095$; $p=0.925$], neither is for anxiety scores [$W(17)=-1.177$; $p=0.239$]. The comparison of scores after hypnosis sessions between T1 and T2 does not show significant results for perceived pain [$W(17)=-1.166$; $p=0.243$] or anxiety [$W(17)=-0.820$; $p=0.412$] (Figure 4).



Quality of life

Total scores for quality of life and sub-scores on the SF-36 were compared between T1 and T2. Analyses reveal no significant difference between the total scores of quality of life at T1 and T2 [$W(10) = 1.125$; $p = 0.260$], neither for different sub-scores such as physical limitation [$W(10) = 1.350$; $p = 0.177$], emotional limitation [$W(10) = 1.382$; $p = 0.167$], emotional well-being [$W(16) = 0.739$; $p = 0.460$], pain [$W(16) = 0.09$; $p = 0.929$], and global health [$W(16) = -0.751$; $p = 0.452$]. When considered as a whole, mental quality of life (i.e., sum of limitations in social activities and usual activities because emotional problem, general mental health and vitality sub-scores) did not differ significantly between T1 and T2 [$W(10) = 0.969$; $p = 0.333$]. However, statistical tendency is observed for several sub-scores such as physical functioning [$W(16) = 1.728$; $p = 0.084$], social functioning [$W(16) = 1.667$; $p = 0.095$], and energy

[$W(16) = 1.657$; $p = 0.097$]. When computed as such (i.e., sum of limitations in social activities and usual activities because physical problem, body pain and general perception of health sub-scores), the dimension of physical quality of life tends to be different between T1 and T2 [$W(10) = 1.682$; $p = 0.093$].

Discussion

This study aimed to evaluate the short- and long-term benefits of HYlaDO, a self-hypnosis approach, on reducing pain and anxiety levels, increasing relaxation level, and improving quality of life in patients with chronic pain. This work was done in the context of the preliminary test phase (Phase II) of the ORBIT model. The results confirmed that the HYlaDO program improves the perception of pain, anxiety and relaxation. Furthermore, we also observed




FIGURE 4

Significant changes in pain, anxiety and relaxation levels for pre- and post-hypnosis across T1 and T2. (A) Pain level (T1), (B) Anxiety level (T1), (C) Relaxation level (T1), (D) Pain level (T2), (E) Anxiety level (T2), (F) Relaxation level (T2).

improvements at 6-month follow-up on quality-of-life sub-scores for some individuals ($N=11$). Our findings regarding the effects of a hypnosis session from the HYLaDO program in pre-post-intervention represents a critical step in the development of a non-pharmacological approach in pain clinical practice. Already, previous research highlights the efficacy of hypnosis in pain management, as demonstrated in numerous fundamental and clinical studies (Lang et al., 2000; Mills et al., 2019; Thompson et al., 2019). Consistent with this body work, our program led to a reduction in perceived pain and anxiety, two central targets among patients suffering from chronic pain, would indicate that these patients can experience a sense of physical and emotional comfort without pharmacological interventions. It is further complemented by the patients' ability to achieve relaxation during the hypnosis session, demonstrating their capacity to regain control over their bodies, often perceived as beyond their control.

Our research corroborates previous findings in chronic pain management, which underline the crucial role of self-care skills in patient treatment (Jensen et al., 2020; Langlois et al., 2022). Our study builds on this work by harnessing patients' endogenous abilities to enhance their well-being, thereby promoting their autonomy. However, our analysis did not show significant differences in the assessments conducted between T1 and T2, with a six-month interval between these measurements. Several reasons may account for the absence of this effect. First, we noted pronounced improvements in self-reported pain perception, anxiety, and relaxation at the session-level. The substantial benefits observed within a single session suggest that improvements across sessions may only be marginal, indicative

of a ceiling effect for these measures. Still, the consistency in results between T1 and T2 suggests that participants were at least able to maintain these improvements over time, underscoring the sustained impact of the intervention.

Secondly, as a preliminary study, our limited sample size constrained our capacity to detect anything but large effect sizes. Despite this, we noted improvements in quality of life for some individuals ($N=11$) between T1 and T2. Quality of life encompasses various factors, divided into emotional and physical sub-scales (Lins and Carvalho, 2016). This possible improvement is specific to the physical sub-scale of the SF-36, which relates to physical functioning, social functioning, and energy. A more extensive sample size in future research would enable a more accurate estimation of effect sizes and provide the statistical power necessary to evaluate these potential benefits more thoroughly. Conversely, the emotional sub-scale of the SF-36, which includes distress and the general perception of one's health. It seems unlikely that a brief intervention, like the one we are proposing here, can swiftly address the complexity of these patients' mental fragility in such a short timeframe. For example, chronic pain conditions lead to significant socio-professional and financial consequences, such as a loss of time at work and reduced financial contributions. Therefore, the absence of an effect across T1 and T2 for this subscale is hardly surprising.

Third, the participants in this study were treated at a pain clinic, and we did not document the proposed treatments in a detailed manner. We could not isolate these treatments either due to our small sample size. For future studies, we will document and introduce them as variables in our analyses.

Lastly, inter-individual variability in hypnotic responding represent another aspects, a fundamental yet frequently neglected aspect of hypnotic phenomena in clinical settings (Houzé et al., 2021). Such variability can stem from diverse factors including psychological background, individual susceptibility to hypnosis, and previous experiences with hypnotic techniques (REFs). This variability implies that individuals react distinctly to identical suggestions (Houzé et al., 2021). Unfortunately, in the context of our study, we did not collect specific information regarding this variability in hypnotic susceptibility among participants. Consequently, our analysis lacks an exploration of how these individual differences in response to hypnosis might have played a role in the outcomes observed. This limitation is significant, as understanding the extent to which hypnotic susceptibility influences therapeutic outcomes could provide valuable insights for tailoring hypnotic interventions more effectively. Future research in this domain should aim to incorporate measures of hypnotic susceptibility to better assess its impact on clinical results. This approach could potentially lead to more personalized and effective therapeutic strategies in the application of hypnotherapy.

In sum, this preliminary study confirmed that a single session from the HYlaDO program benefits chronic pain patients along several dimensions. However, we could not confirm the benefits of the program between T1 and T2. Considering the milestones outlined in the ORBIT model, this justifies advancing to the next stage, a pilot randomized controlled study, aimed at testing our protocol and gathering data. This step will enable us to estimate effect sizes and calculate the sample size required for a future clinical trial (ORBIT III). Ultimately, if this project proves effective, it could be widely offered in pain clinics as a non-pharmacological approach based on hypnosis.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The studies involving humans were approved by Maisonneuve-Rosemont Hospital Ethics (CER No: 2022-2741). The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

References

- Bicego, A., Monseur, J., Collinet, A., Donneau, A. F., Fontaine, R., Libbrecht, D., et al. (2021). Complementary treatment comparison for chronic pain management: a randomized longitudinal study. *PLoS One* 16:e0256001. doi: 10.1371/journal.pone.0256001
- Canadian pain task force (2019) *Chronic pain in Canada: laying a foundation for action*. Available at: <https://www.canada.ca/content/dam/hc-sc/documents/corporate/about-health-canada/public-engagement/external-advisory-bodies/canadian-pain-task-force/report-2019/canadian-pain-task-force-june-2019-report-en.pdf>
- Canadian pain task force (2020). Working together to better understand, prevent and manage chronic pain: What We Heard. Available at: <https://www.canada.ca/en/health-canada/corporate/about-health-canada/public-engagement/external-advisory-bodies/canadian-pain-task-force/report-2020.html>
- Caron-Trahan, R., Jusseaux, A.-E., Aubin, M., Cardinal, É., Aramideh, J., Idrissi, M., et al. (2023a). Practicing self-hypnosis to reduce chronic pain: a qualitative exploratory study of HYlaDO. *Br. J. Pain* 18, 28–41. doi: 10.1177/20494637231200324
- Caron-Trahan, R., Jusseaux, A. E., Aubin, M., Urbanowicz, R., Richebé, P., and Ogez, D. (2023b). Definition and refinement of HYlaDO, a self-hypnosis training program for chronic pain management: a qualitative exploratory study. *Explore* 19, 417–425. doi: 10.1016/j.explore.2022.08.008
- Choinière, M., Peng, P., Gilron, I., Buckley, N., Williamson, O., Janelle-Montcalm, A., et al. (2020). Accessing care in multidisciplinary pain treatment facilities continues to be a challenge in Canada. *Reg. Anesth. Pain Med.* 45, 943–948. doi: 10.1136/rapm-2020-101935
- Czajkowski, S. M., Powell, L. H., Adler, N., Naar-King, S., Reynolds, K. D., Hunter, C. M., et al. (2015). From ideas to efficacy: the ORBIT model for developing

Author contributions

DO: Conceptualization, Formal analysis, Funding acquisition, Methodology, Project administration, Supervision, Validation, Writing – original draft, Writing – review & editing. ML: Formal analysis, Validation, Writing – review & editing. RC-T: Formal analysis, Investigation, Writing – review & editing. A-EJ: Investigation, Writing – review & editing. MA: Conceptualization, Funding acquisition, Writing – review & editing. JV: Writing - review & editing. VF: Formal analysis, Validation, Writing – review & editing. NG: Writing – review & editing. MI: Formal analysis, Validation, Writing – review & editing. PRA: Writing – review & editing. PRI: Resources, Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. DO and MA are recipients of funding from the Fonds de Recherche du Québec (Programme Engagement Citoyen). DO is also a recipient of a Fonds de recherche du Québec salary grant (clinician-researcher J1).

Acknowledgments

We would like to thank the Pain Clinic patients who participated in this study, as well as our partner patient. We also acknowledge the support of the Fonds de Recherche du Québec (Citizen Engagement program) for the funding of this project.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

behavioral treatments for chronic diseases. *Health Psychol.* 34, 971–982. doi: 10.1037/hea0000161

Eaton, L. H., Beck, S. L., and Jensen, M. P. (2021). An audio-recorded hypnosis intervention for chronic pain Management in Cancer Survivors: a randomized controlled pilot study. *Int. J. Clin. Exp. Hypn.* 69, 422–440. doi: 10.1080/00207144.2021.1951119

Hammond, D. C. (2001). Treatment of chronic fatigue with neurofeedback and self-hypnosis. *NeuroRehabilitation* 16, 295–300. doi: 10.3233/NRE-2001-16415

Houzé, B., Streff, A., Piché, M., and Rainville, P. (2021). Spinal and supraspinal modulation of pain responses by hypnosis, suggestions, and distraction. *Am. J. Clin. Hypn.* 63, 329–354. doi: 10.1080/00029157.2020.1863184

Jensen, M. P., McArthur, K. D., Barber, J., Hanley, M. A., Engel, J. M., Romano, J. M., et al. (2006). Satisfaction with, and the beneficial side effects of, hypnotic analgesia. *Int. J. Clin. Exp. Hypn.* 54, 432–447. doi: 10.1080/00207140600856798

Jensen, M. P., Mendoza, M. E., Ehde, D. M., Patterson, D. R., Molton, I. R., Dillworth, T. M., et al. (2020). Effects of hypnosis, cognitive therapy, hypnotic cognitive therapy, and pain education in adults with chronic pain: a randomized clinical trial. *Pain* 161, 2284–2298. doi: 10.1097/j.pain.0000000000001943

Lang, E. V., Benotsch, E. G., Fick, L. J., Lutgendorf, S., Berbaum, M. L., Berbaum, K. S., et al. (2000). Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomised trial. *Lancet* 355, 1486–1490. doi: 10.1016/S0140-6736(00)02162-0

Langlois, P., Perrochon, A., David, R., Rainville, P., Wood, C., Vanhaudenhuyse, A., et al. (2022). Hypnosis to manage musculoskeletal and neuropathic chronic pain: a systematic review and meta-analysis. *Neurosci. Biobehav. Rev.* 135:104591. doi: 10.1016/j.neubiorev.2022.104591

Lins, L., and Carvalho, F. M. (2016). SF-36 total score as a single measure of health-related quality of life: scoping review. *SAGE Open Med* 4:2050312116671725. doi: 10.1177/2050312116671725

Milling, L. S., Valentine, K. E., LoStimolo, L. M., Nett, A. M., McCarley, H. S. (2021). Hypnosis and the alleviation of clinical pain: a comprehensive meta-analysis. *Int. J. Clin. Exp. Hypn.* 69, 297–322. doi: 10.1080/00207144.2021.1920330

Mills, S. E. E., Nicolson, K. P., and Smith, B. H. (2019). Chronic pain: a review of its epidemiology and associated factors in population-based studies. *Br. J. Anaesth.* 123, e273–e283. doi: 10.1016/j.bja.2019.03.023

Samami, E., Shahhosseini, Z., and Elyasi, F. (2021). The effect of psychological interventions on the quality of life in women with fibromyalgia: a systematic review. *J. Clin. Psychol. Med. Settings* 28, 503–517. doi: 10.1007/s10880-021-09794-0

Thompson, T., Terhune, D. B., Oram, C., Sharangparni, J., Rouf, R., Solmi, M., et al. (2019). The effectiveness of hypnosis for pain relief: a systematic review and meta-analysis of 85 controlled experimental trials. *Neurosci. Biobehav. Rev.* 99, 298–310. doi: 10.1016/j.neubiorev.2019.02.013

Thong, I. S. K., Jensen, M. P., Miro, J., and Tan, G. (2018). The validity of pain intensity measures: what do the NRS, VAS, VRS, and FPS-R measure? *Scand J Pain* 18, 99–107. doi: 10.1515/sjpain-2018-0012