



HAL
open science

Nitrous oxide abuse in the emergency practice, and review of toxicity mechanisms and potential markers.

Marie Joncquel, Guillaume Grzych, Celine Tard, Julien Lannoy, Sylvie Deheul, Riyad Hanafi, Claire Douillard, Joseph Vamecq

► To cite this version:

Marie Joncquel, Guillaume Grzych, Celine Tard, Julien Lannoy, Sylvie Deheul, et al.. Nitrous oxide abuse in the emergency practice, and review of toxicity mechanisms and potential markers.. Food and Chemical Toxicology, 2022, Food and Chemical Toxicology, 162, pp.112894. 10.1016/j.fct.2022.112894 . hal-04610412

HAL Id: hal-04610412

<https://hal.univ-lille.fr/hal-04610412v1>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Nitrous oxide abuse in the emergency practice, and Review of toxicity mechanisms and potential markers

Marie Joncquel ChevalierCurt¹, Guillaume Grzych^{1,2}, Céline Tard³, Julien Lannoy⁴, Sylvie Deheul⁵, Riyad Hanafi⁶, Claire Douillard⁷ and Joseph Vamecq^{8,*}

¹ CHU Lille, Service d'Endocrinologie, Métabolisme, Nutrition, Oncologie, F-59000 Lille, France

² Univ. Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U1011- EGID, F-59000 Lille, France

³NeurologyDepartment, CHU Lille, Lille, France

⁴ CHU Lille, Service de Neurologie, Lille, France

⁵ CHU Lille, Addictovigilance, Lille, France

⁶ CHU Lille, Medical Imagery, Lille France,

⁷ Endocrinology-Diabetology-Metabolism Department and Medical Reference Center for Inherited Metabolic Diseases Jeanne de Flandre Hospital, CHU Lille, Lille, France,

⁸ Inserm, Univ. Lille EA 7364 RADEME, CHU Lille, Centre de Biologie Pathologie Génétique, UF Métabolisme Général et Maladies Rares, F-59000 Lille, France (ORCID ID 0000-0002-4089-7663).

M. Joncquel Chevalier Curt (First Author)
MARIE.JONCQUEL@chru-lille.fr

G. Grzych
Guillaume.GRZYCH@chu-lille.fr

C. Tard
CELINE.TARD@chru-lille.fr

J. Lannoy
Julien.LANNOY@chru-lille.fr

S. Deheul
Sylvie.DEHEUL@chru-lille.fr

R. Hanafi
Riyad.HANAFI@chru-lille.fr

C. Douillard
Claire.DOUILLARD@chru-lille.fr

J. Vamecq (✉)
joseph.vamecq@inserm.fr

* Corresponding author. Tel.: +33 320 44 57 02. e-mail: joseph.vamecq@inserm.fr.

Full postal address: Dr Joseph Vamecq, Biochimie et Biologie Moléculaire, HMNO, CBP, CHRU Lille, 2, Avenue Prof Jules Leclercq, 59037 Lille Cedex, France

ABSTRACT

Nitrous oxide (N₂O) toxicity is a concern common to several medical fields. Here, retrospective study of four N₂O abuses with neurological signs in the emergency practice provides a preliminary basis for a metabolic Discussion/Review. This latter highlights N₂O abuse as pathology of DNA/RNA/protein methylations, for instance consistent with impairments of protein arginine methyltransferases involved in myelinogenesis and myelopathy in patients. Basically, pathogenesis starts with oxidation by N₂O of coordinated cobalamine cobalt ions at enzyme sites with impairments of vitamin-B12-dependent pathways. Methionine synthase (methylcobalamine) and methylmalonyl-CoA mutase (adenosylcobalamine) are inactivated and cofactor-depleted, respectively. The number of impacted pathways (folate cycle, methylation cycle, S-adenosylmethionine-dependent methyltransferases, transulfuration pathway, Krebs cycle fueling by methylmalonyl-CoA, glutathione synthesis) explains the variety of potential research/laboratory markers, and may provide new clues and future angles to explore N₂O toxicity. Overall, homocysteine measurements obviously help diagnosis of N₂O abuses. Additional markers may include vitamin-B12, methionine, methylmalonate, dimethylglycine, sarcosine, S-adenosylmethionine to S-adenosylhomocysteine ratio, various S-adenosylamino acids, S-adenosylmethionine-dependent cellular methylations, and additional analytes (propionylcarnitine, propionylglycine, cystathionine and derived metabolites, methylated amino acids [eg arginine], betaine).

Keywords

Nitrous oxide abuse; cobalamins; vitamin B12; methionine synthase; methylmalonyl-CoA mutase; methylation cycle; folate cycle; homocysteine; biological markers

1. Introduction

Nitrous oxide (N₂O) was discovered in 1772. This 'laughing gas' has medical indications which include analgesia, anxiolysis and anaesthesia, and date back to 1844 with analgesia in dental surgery, then anaesthesia (Kongara et al., 2021; Oussalah et al., 2019). Recreational use also dates from this period. Facilitated by its easy availability as a propellant gas in aerosol cans for food use, it has recently re-emerged as a cause of hallucinations, excitement and psychological dependence in the young population (van Amsterdam et al., 2015). N₂O poured into "whippits" is inhaled by users to meet euphoria. However, prolonged high consumption of N₂O causes side effects among which vitamin B12 deficiency, severe neurological deterioration such as the degeneration of the spinal cord (Dong et al., 2019; Lan et al., 2019; McArdle and Gaillard, 2020; Seed and Jogia, 2020) and, *via* hyperhomocysteinemia, pulmonary embolisms, venous and arterial thromboses, psychiatric disorders and ischaemic strokes (Bajaj et al., 2018; Chien et al., 2020; den Uil et al., 2018; Pratt et al., 2020).

Though well documented for methylcobalamin at methionine synthase enzyme site, the inactivating effects of N₂O on non-enzyme bound vitamin B12 and adenosylcobalamin encased in methymalonyl-CoA mutase still need to be clarified (see Discussion). Vitamin B12 deficiency is double-edged being consequence and worsening factor of N₂O toxicity. The present retrospective study of data concerns four young (18-23 years old) patients who developed N₂O abuse before the Covid-19 outbreak. The discussion stresses the clinical and biological signs that should alert on N₂O intoxication, and the panoply of biomarkers potentially available to diagnosing this addiction. A brief account of patient 3 has been published previously (Grzych et al., 2020).

2. N₂O abuse in the emergency practice

The present retrospective study of cases of N₂O abuse in the emergency practice serves as a preliminary basis to review underlying toxicity mechanisms and potential markers (Discussion). Plasma amino acids concentrations (μmol/L) were determined as described by Boemer et al., 2015. Briefly, plasma amino acids were amine-modified using the aTRAQ™ labeling reagent, which provides a specific mass tag for tandem mass spectrometry (MS). The tag is identified by MS/MS fragmentation of analytes and standards under the multiple reaction monitoring mode. An internal standard (IS) set of aTRAQ™ labeled amino acids was used for both detection and quantification of analytes by HPLC (Shimadzu C18 column, Kyoto, Japan) separation coupled with MS/MS (Sciex 3200 Qtrap, Framingham, MA, US). Quantification was performed by dividing peak area of the analyte with

peak area of the respective IS, the calculated quotient being multiplied by the respective IS concentration. Vitamins B1 and B6 were assayed using ChromSystemS protocols (Grärfeling, Germany) on whole blood and plasma samples, respectively. Sample clean-up was achieved by protein precipitation. Derivatization of supernatants was performed to obtain fluorescent analytes. Isocratic HPLC separation was followed by fluorimetric detection through a procedure adapted from the method of [Reynold and Brain \(1992\)](#). Seric vitamins B9 (folates) and B12 were measured by immunoassays (Access Folates and Access Vitamin B12 cobalamine protocols, respectively, Beckman Coulter, CA).

Patient 1 - A 19-year-old man was admitted for gait disturbance and paresthesia affecting left leg and both hands. These signs had developed gradually over the preceding week. One week ago, patient had also developed balance problems with a fall episode. The history indicated no personal and family history for a neurological disease. Alcohol consumption and smoking (6 packs per year) were occasional. By contrast, cannabis was consumed daily (2 to 3 times per day), and N₂O at 200 to 300 whippits per day. Clinically, the ataxia was major with the need for an assistive device to walk. Generalized areflexia and apallesthesia of the lower limbs were also observed. Vitamin B12 assessed on admission was at the lower limit of reference values (0.2 nmol/L [control range: 0.2-1.0 nmol/L]) (Table 1). The MRI scan showed posterior cervical spinal cord lesion spanning from the C2 to C6 vertebral levels and suggestive of subacute combined degeneration (Fig. 1). This myelopathy associated with vitamin B12 deficiency prompted biological investigations to identify any underlying nutritional or metabolic deficiency. Low plasma vitamin B6 (8 nmol/L [control values: 15-73 nmol/L]), and significantly increased total plasma homocysteine (144 µmol/L [control values < 14 µmol/L]) were observed (Table 1). There was no anaemia and the GMV was normal. IV vitamin B12 supplementation was followed by oral relay (1 mg/day for 1 month), as well as completed with vitamin B9 (5 mg/day for 1 month) and vitamin B6 (500 mg/day for 1 month) (Table 1) therapies. Discontinuation of N₂O was also undertaken during upon hospitalization. Signs gradually improved and patient was discharged after a few days of the multi-vitamin therapy.

Patient 2 - A 19-year-old woman was admitted with paresthesia of the lower limbs. The patient followed a meat-free diet. For 1 year and at repeated time intervals, she consumed 200 whippits of N₂O per day (Table 1). Four months ago, paresthesia developed with hypoesthesia extending from the feet to the knees and from the hands to the elbows without motor deficit. The symptoms regressed with the withdrawal of N₂O, and sensory disturbances in the lower limbs reappeared as

soon as patient resumed N₂O consumption for a few days. Unstable walking caused a fall with knee trauma. On examination, patient was conscious without signs of disorientation. MRI of the spine was normal. Vitamin B12 was collapsed below 0.2 nmol/L in parallel with a significant increase in total plasma homocysteine (142 µmol/L [control values < 14 µmol/L]) (Table 1). IM vitamin B12 supplementation followed by oral relay (1 mg/day for 1 month) was combined with oral folic acid (5 mg x2/day for 1 month) (Table 1).

Patient 3 - An 18-year-old man was referred for subacute walking difficulties. For six weeks, he suffered from heavy leg signs. For one year, patient consumed 100 whippits of N₂O every weekend (Table 1); he spontaneously halved consumption when affected with walking difficulties, and stopped intake when convinced of a link between consumption and signs. The sensation of heavy leg was accentuated on exertion, when climbing stairs and walking for long periods. Three weeks before present admission, the patient referred to another hospital and diagnosed with subacute combined spinal cord degeneration caused by N₂O, omitted to take the prescribed vitamin B12 when out of the hospital. Walking difficulties with neuropathic pain were the cause of the present consultation. The electroneuromyogram showed a severe symmetrical sensitivomotor polyneuropathy of axonal origin. Bilateral carpal tunnel syndrome was also diagnosed. Plasma vitamin B12 was at the lower limit of reference values, total serum homocysteine (123 µmol/L [control values < 14 µmol/L]) was increased (Table 1). IM vitamin B12 supplementation was administered at a dose of 1 mg/day. After 7 days of hospitalisation, the patient was transferred to a rehabilitation unit and 10 days later, without medical agreement, escaped from this unit, resulting in a loss of medical follow-up (Table 1).

Patient 4 - A 23-year-old man was admitted for delusion in the context of a paranoid disorientation for 7 days. On arrival he was agitated with hallucinations. He was treated with Tercian and Rivotril. Heteroanamnesis revealed a similar episode a few months ago, and consumption of 100 whippits of N₂O per day for 2 years. Patient had no personal psychiatric history. The cerebral CT scan failed to detect haemorrhagic lesion. Plasma vitamin B12 concentration was below baseline and total serum homocysteine was increased (164 µmol/L [control values < 14 µmol/L]) (Table 1).

3. Discussion, Review of mechanisms and potential markers of N₂O toxicity

3.1. Recapitulative clinics, biology and therapeutic measures in the four reported cases - Alert signs for nitrous oxide abuse

Table 1 provides a brief summary of the clinical history, laboratory data and medical treatments of the four patients. Diagnosing N₂O abuse has been aided with the anamnesis of patients, and,

heteroanamnesis conducted in their immediate social entourage. Recreational uptakes of 100 or more whippits per day caused signs which led patients to consult and be hospitalized. Patients had deficient or low levels of vitamin B12 (Table 1). Gait disturbances were present in all but one patient. In this patient, clinical signs were dominated by a delirium. This change in mental status has been previously reported in cases of N₂O abuse (Sterman and Coyle, 1983). In the other three patients, gait disturbances were associated with paresthesia, heaviness and pain of legs. Hands were also affected as reported spontaneously by patients or diagnosed after medical examination including EMG. Paraesthesia, gait instability and weakness are common in patients with N₂O abuse (Garakani et al., 2016; Xiang et al., 2021). These signs should thus lead to include N₂O abuse in the differential diagnosis (Fernández et al., 2017). One of the two patients with subacute combined degeneration of spinal cord had also vitamin B6 deficiency, raising a possible protection of vitamin B6 against myelopathy caused by vitamin B12 deficiency (see below).

3.2. Effects of nitrous oxide on cobalamin-dependent proteins, and general consequences on metabolism and pathogenesis

Vitamin B12 encompasses a group of molecules usually referred to as cobalamins and which share in common a molecular structure which coordinates a cobalt atom within a corrinoid ring (Froese et al., 2019). Essential forms of cobalamins involved in human metabolism include non-exhaustively methylcobalamin and adenosylcobalamin which are cofactors of cytoplasmic methionine synthase and mitochondrial methylmalonyl-CoA-mutase, respectively (Froese et al., 2019). Vitamin B12 is also involved in erythropoiesis and CNS homeostasis. N₂O impairs cobalamin (vitamin B12)-dependent metabolic pathways through the oxidation by N₂O of cobalt ions from the vitamin corrinoid ring leading to inactivation and deficiency of vitamin B12 (Sharma et al., 2003) along with its biological manifestations including increased homocysteine and methylmalonic acid (Xiang et al., 2021).

Fig.2 illustrates the metabolic pathways dependent on cobalamin cofactors, stressing their targeting by N₂O as well as some reciprocal regulations taking place between methylation and folate cycles. Subcellular sites for the main metabolic reactions are illustrated as well as interactions between pathways including folate cycle, methylation cycle, transsulfuration pathway, mitochondrial succinyl-CoA metabolism, cytosolic glutathione synthesis, and cell/tissue S-adenosylmethionine-dependent methyltransferases (for a more complete consideration including the nuclear component of folate metabolism, see Lan et al, 2018; Tibbetts and Appling 2010).

Fig.2 points a potential protective role of vitamin B6 (evoked above for Patient 1) against vitamin B12 deficiency by linking homocysteine to glutathione synthesis *via* cysteine formation. Other mechanisms for neuroprotective properties of vitamin B6 rely on PMK2-NRF2 signaling and up-regulation of glutathione biosynthesis genes (Wei et al., 2020). Reduced availability of vitamin B6

could thus compromise antioxidant defenses and repair mechanisms that might protect against subacute combined spinal cord degeneration. Functional deficiency of S-adenosylmethionine-dependent methyltransferases including protein arginine methyltransferases (PRMT) 1 and 5 which are critical for myelin formation (Hashimoto et al., 2021) might also promote this myelopathy. The inactivation by N₂O of methionine synthase [vitB12-dependent enzyme] might impair PRMT1 and 5 *via* either reduced supply in S-adenosylmethionine or increased product inhibition by S-adenosylhomocysteine. The latter vs former mechanism might prevail based on normal levels of methionine observed in patients.

The analysis of N₂O-impacted pathways suggests two mechanisms contributing to the rise of methylmalonic acid : its decreased handling by mitochondrial methylmalonyl-CoA mutase and its enhanced synthesis from the α -ketobutyrate derived, *via* cystathionine, from homocysteine accumulating secondarily to impaired methionine synthase (Fig.2).

3.3. Biological markers for N₂O abuse

3.3.1. Distinct inactivation patterns of methionine synthase and methylmalonyl-CoA mutase

Altered levels in blood vitamin B12, homocysteine and methionine, and urinary methylmalonic acid have been largely described in N₂O abuse. Their genesis, and kinetics, may be understood by taking into account the recoveries of methylmalonyl-CoA mutase and methionine synthase activities after exposure to N₂O. Riedel and coworkers (Riedel et al., 1999) provided evidence that cobalamin oxidation by N₂O at methionine synthase enzyme site involves hydroxyl radical and subsequent irreversible oxidative enzyme inactivation. As a result, *de novo* enzyme synthesis is needed to recovering enzyme function. By contrast, the apoenzyme of methylmalonyl-CoA mutase is preserved after N₂O, and recovery of the mutase activity requires only the replenishment of the corrinoid ring with the cobalamin enzyme cofactor (Riedel et al., 1999). As a conclusion, methionine synthase is oxidatively and irreversibly inactivated by N₂O whereas methylmalonyl-CoA mutase is essentially cofactor depleted. In addition, a rapid recovery of the mutase activity, which relies on release of the altered cobalamin cofactor and its replacement by an intact one, is promoted by MMAA (Takahashi-Iñiguez et al., 2017). These two distinct patterns of enzyme alterations by N₂O may explain the persistence of elevated homocysteine but not methylmalonic acid levels when patients both receive vitamin B12 therapy and maintain N₂O consumption (Nunn, 1984). Other considerations for distinct impactation of the two enzymes by N₂O might lie for the mutase (*vs* synthase), in the mitochondrial (*vs* cytosolic) location and cobalamin flanking adenosyl (*vs* methyl) substitution. Whether these enzyme specificities (mitochondrial membrane barriers, steric hindrance in the catalytic site) might protect the mutase enzyme site from N₂O poisoning remains to be clarified.

3.3.2. Mechanisms of vitamin B12 deficiency induced by N₂O oxidation of cobalamin cobalt ions

As mentioned just above, N₂O-driven irreversible oxidation of cobalamin cobalt ions at enzyme sites impacts irreversibly methionine synthase and reversibly methylmalonyl-CoA mutase. The recovery of methionine synthase and methylmalonyl-CoA mutase from N₂O-induced inactive states is inevitably made at the expense of available vitamin B12 stores *via* incorporation of cobalamins in newly synthesized and recycled apoenzymes, respectively. This may contribute to vitamin B12 deficiency which in turn may worsen back the impairment of the two vitamin B12-dependent enzymes.

3.3.3. Potential markers of N₂O abuse

Vitamin B12 is a potential reference biological marker of N₂O abuse. For the reasons just mentioned above, N₂O abuse may induce a drop in blood vitamin B12. The absence of status of vitamin measurements before N₂O abuse does not allow to assess the drop but only the possible dropped value. When vitamin B12 value is in normal range, vitamin B12 deficiency or drop may not be stated in N₂O abuse. In two of the four patients, serum vitamin B12 values were at the lowest normal range values. Such values might be considered to be pathological since reference values are based on a large population inevitably including some deficient patients. Finally, the social food habits of N₂O consumers do not rule out a poor nutritional status responsible for vitamin B12 deficiency. As mentioned above, such a pre-existing deficiency in vitamin B12 would exacerbate the gas toxicity.

Methylmalonic acid accumulation, and inhibition of methylmalonyl-CoA mutase may occur under N₂O abuse. For the reasons mentioned above, methylmalonic acid might be a less reliable marker than homocysteine as more as, in contrast to methionine synthase, inactivation of methylmalonyl-CoA mutase still remains unclear (Frasca et al., 1986; Waclawik et al., 2003). Nevertheless, methylmalonic aciduria is regularly observed in patients with N₂O abuse; it should lead to check vitamin B12 status and plasma homocysteine levels when drug abuse is suspected. Increased levels of plasma valine can result from methylmalonyl-CoA mutase deficiency (Riedel et al., 1999). This is consistent with metabolic fuelling of methylmalonyl-CoA by valine (Fig. 2).

Increased plasma homocysteine is the direct result of N₂O-targeted methionine synthase. It was observed in the four patients with values 10-fold the reference highest value, suggesting it might be a highly sensitive marker of N₂O abuse. Despite alternative etiologies, differential diagnosis should immediately evoke N₂O abuse in a context of drug intake. Reciprocally, plasma homocysteine might be measured when N₂O abuse is suspected.

Methionine, the product of methionine synthase, is expected to be decreased under N₂O. However, as observed in our patients, plasma methionine concentrations may be in normal range values. So, the drop in plasma methionine should not be considered as a reliable marker of N₂O abuse. Compensatory mechanisms that might maintain normal levels of methionine are diet and body

(proteolysis) supplies, alternative synthesis from homocysteine (BHCMT), and decreased consumption by the methylation cycle.

Dimethylglycine and sarcosine are discussed because of increased plasma sarcosine in one of the four patients. In this light, sarcosine does not appear to be a reliable marker in N₂O abuse. How dimethylglycine levels behave under N₂O needs clarification.

S-adenosylmethionine/S-adenosylhomocysteine balance, and related individual S-adenosyl-amino acid levels are physiologically controlled by GNMT activity. This latter is negatively regulated by 5-methyl THF (**Fig. 2**) (Simile et al., 2018). In return, S-adenosylmethionine one of the substrates (the other being glycine) of GNMT inhibits methylenetetrahydrofolate reductase (MTHFR) (Simile et al., 2018) with as consequences (in the absence of N₂O) a rise in 5-methyl THF (THF, tetrahydrofolate) and increased conversion of homocysteine to methionine. It also inhibits the other methionine-synthesizing enzyme BHCMT (Simile et al., 2018). (**Fig.2**). This balance between, from the one hand, methylation cycle inhibition (at the level of GNMT) by a folate cycle intermediate (5-methyl THF) and, on the other hand, folate cycle inhibition (at the MTHFR step) by a methylation cycle intermediate (S-adenosylmethionine) represents a fine tuning to adapt the folate cycle activity to the cellular demand for S-adenosyl-dependent methylations. N₂O may impact this S-adenosylmethionine/S-adenosylhomocysteine balance and related individual S-adenosyl-amino acid levels essentially by inducing a huge rise in S-adenosylhomocysteine (Molloy et al., 1990), making this S-substituted amino acid a potential interesting biological marker for N₂O abuse.

Cellular S-adenosylmethionine-dependent methylations is expected to be affected by the N₂O-mediated inactivation of methionine synthase. The enzyme inactivation though not leading to decreased levels of methionine (see above) might alter the steady-state concentrations and availability of methionine for S-adenosylmethionine synthesis, and hence the availability of S-adenosylmethionine for cellular methylations. The S-adenosyladenosine-dependent methylation product to substrate ratios might clarify subsequent impact on the individual S-adenosyladenosine methyltransferases (see **Fig. 2** for a rapid non-exhaustive listing), and in turn be exploited as markers of the N₂O abuse. Serious limits in this approach arise when metabolites take place in several metabolic pathways (for instance glycine) or are supplied by the diet. For instance, creatine the product of guanidinoacetate methyltransferase is present in energetic drinks which when taken-up by N₂O users may affect creatine levels and creatine to guanidinoacetate ratio. In the same time, increased guanidinoacetate secondarily to deficient S-adenosyladenosine availability could witness for N₂O abuse. Substrates and products of S-adenosyl-dependent methylations and their ratios warrant future clarifications about their suitability as biological markers of N₂O abuse.

Other metabolites appearing on **Fig.2** might be checked for their suitability to be biological markers of N₂O abuse. Briefly, these might include propionylcarnitine (reduced consumption and increased

production of propionyl-CoA secondarily to inactivated methylmalonyl-CoA mutase and methionine synthase, respectively), propionylglycine (for the same reasons as propionylcarnitine, and hypothetic, increased glycine availability), cystathionine and derived metabolites (secondarily to methionine synthase deficiency) such as 2-oxobutyrate (mentioned above) and intermediates of the homocysteine transulfuration pathway (taurine, for instance). Methylated aminoacids resulting from hydrolysis of proteins which have been subjected to S-adenosylmethionine-dependent methylations are also of interest, for instance symmetric and asymmetric dimethylarginine (SDMA and ADMA). Though, in contrast to homocysteine, ADMA was not impacted by the acute anesthetic use of N₂O (Myles et al., 2008), chronic recreational use of N₂O might affect dimethylarginine levels *via* chronic impairment of the methylation cycle resulting from methionine synthase inactivation. Betaine consumption (and hence reduced levels of betain and its precursor choline) which yields dimethylglycine can be decreased by the stimulated fuelling of BHCMT by high homocysteine levels (Imbard et al., 2015). So, ratios to betain of metabolites which are expected to increase in patients with N₂O abuse obviously also warrant future consideration.

4. Conclusions

Grosso modo, chronic N₂O toxicity exhibits most clinical and metabolic features of vitamin B12 deficiency. Clinically, recreational N₂O users may develop walking difficulties and pain, and more severe neurological signs ranging from sensory-motor polyneuropathy to subacute combined spinal cord degeneration and, delirium. Biologically, patients usually present with decreased serum vitamin B12 (near or below the lower limit of normal) and markers for deficiency of vitamin B12-dependent enzymes (cytosolic methionine synthase and mitochondrial methylmalonyl-CoA mutase) such as increased homocysteine and decreased methionine, and increased methylmalonate.. N₂O abuse should be primarily considered as a severe methionine synthase deficiency associated with partially to severely deficient methylmalonyl-CoA mutase. In this light, the pathogenesis of N₂O abuse should be viewed as a disorder of cellular DNA, RNA and protein methylations, consistent with impairment of S-adenosylmethionine-dependent methyltransferases such as PRMT1 and 5 involved in myelinogenesis and functional deficiencies of which are consistent with myelopathy seen in patients. Treatment relies on cessation of N₂O intake and supplementation of vitamin B12. Treatment with vitamin B6 is also recommended when blood levels of this vitamin are decreased. Moreover, vitamin B12 supplementation should be combined with vitamin B6 treatment in cases of N₂O overdose, even in the absence of vitamin B6 depletion. This dual measure might help the detoxifying of the methylation cycle by removing excess homocysteine *via* methionine [progressive recovery of methionine synthase activity, VitB12-dependent] and cysteine [stimulation of cystathionine synthase,

VitB6-dependent]. Limiting access to N₂O whippits and regular measurements of homocysteine or other potential relevant marker to monitor patient compliance can also be recommended.

Acknowledgments

The authors thank the CHU of Lille and the Inserm for the support given to the authors' work.

CRedit author statement

MJCC, GG, SD, JV: Investigation, Methodology ; MJCC, GG, JV: Biological Data curation ; CT, JL, SD, CD: Clinical Data curation ; RH: MRI Data curation ; MJCC, GG, JV: Writing-Original draft preparation ; MJCC, GG, CT, CD, JV: Writing - review & editing ; MJCC, GG, CT, SD, CD, JV: Supervision. All authors approve this submission for publication.

Declaration of Competing Interest

The authors declare that they have no conflict of interest.

REFERENCES

- Bajaj, D., Agrawal, A., Gupta, S., et al., 2018. Recreational Nitrous Oxide Abuse Causing Ischemic Stroke in a Young Patient: A Rare Case Report. *Cureus* 10, e3761. <https://dx.doi.org/10.7759/cureus.3761>.
- Boemer, F., Schoos, R., Deberg, M., 2015. Quantification des acides aminés physiologiques par le kit aTRAQ(®) : évaluation et implémentation de nouveaux paramètres [Quantification of physiological aminoacids using aTRAQ(®) kit: evaluation and implementation of new markers]. *Ann. Biol. Clin. (Paris)* 73, 427-442. <https://dx.doi.org/10.1684/abc.2015.1066>
- Chien, W.H., Huang, M.C., Chen, L.Y., 2020. Psychiatric and Other Medical Manifestations of Nitrous Oxide Abuse: Implications From Case Series. *J. Clin. Psychopharmacol.* 40, 80-83. <https://dx.doi.org/10.1097/JCP.0000000000001151>.
- den Uil, S.H., Vermeulen, E.G.J., Metz, R., et al., 2018. Aortic arch thrombus caused by nitrous oxide abuse. *J. Vasc. Surg. Cases Innov. Tech.* 4, 80-82. <https://dx.doi.org/10.1016/j.jvscit.2018.01.001>.
- Dong, X., Ba, F., Wang, R., et al., 2019. Imaging appearance of myelopathy secondary to nitrous oxide abuse: a case report and review of the literature. *Int. J. Neurosci.* 129, 225-229. <https://dx.doi.org/10.1080/00207454.2018.1526801>.
- Fernández, D., Fara, M.G., Biary, R., et al., 2017. Clinical Reasoning: A 27-year-old man with unsteady gait. *Neurology* 89, e120-e123. <https://dx.doi.org/10.1212/WNL.0000000000004327>.
- Frasca, V., Riazzi, B.S., Matthews, R.G, et al., 1986. In vitro inactivation of methionine synthase by nitrous oxide. *J. Biol. Chem.* 261, 15823-15826. [https://dx.doi.org/10.1016/S0021-9258\(18\)66636-0](https://dx.doi.org/10.1016/S0021-9258(18)66636-0).

- Froese, D.S., Fowler, B., Baumgartner, M.R., 2019. Vitamin B12 , folate, and the methionine remethylation cycle-biochemistry, pathways, and regulation. *J. Inherit. Metab. Dis.* 42, 673-685. <https://dx.doi.org/10.1002/jimd.12009>.
- Garakani, A., Jaffe, R.J., Savla, D., et al., 2016. Neurologic, psychiatric, and other medical manifestations of nitrous oxide abuse: A systematic review of the case literature. *Am. J. Addict.* 25, 358-369. <https://dx.doi.org/10.1111/ajad.12372>.
- Grzych, G., Douillard, C., Lannoy, J., et al., 2020. Very High Plasma Homocysteine without Malnutrition or Inherited Disorder. *Clin. Chem.* 66, 1468-1469. <https://dx.doi.org/10.1093/clinchem/hvaa070>.
- Hashimoto, M., Fukamizu, A., Nakagawa, T., et al., 2021. Roles of protein arginine methyltransferase 1 (PRMT1) in brain development and disease. *Biochim. Biophys. Acta GenO Subj.* 1865, 129776. <https://dx.doi.org/10.1016/j.bbagen.2020.129776>.
- Imbard, A., Benoist, J.F., Esse, R., et al., 2015. High homocysteine induces betaine depletion. *Biosci Rep.* 35, e00222. <https://dx.doi.org/10.1042/BSR20150094>.
- Kongara, G., Pula, R., Thakur, N., et al., 2021. Adverse Effects of Nitrous Oxide on Vitamin B12 Levels in Health Care Personnel of ESIC Tertiary Care Hospital. *J. Cell. Mol. Anesth.* 6, 9-14. <https://dx.doi.org/https://doi.org/10.22037/jcma.v6i1.32944>.
- Lan, S.Y., Kuo, C.Y., Chou, C.C., et al., 2019. Recreational nitrous oxide abuse related subacute combined degeneration of the spinal cord in adolescents - A case series and literature review. *Brain Dev.* 41, 428-435. [https://dx.doi.org/https://doi.org/10.1016/0165-6147\(84\)90426-7](https://dx.doi.org/https://doi.org/10.1016/0165-6147(84)90426-7).
- Lan, X., Field, M.S., Stover, P.J., 2018. Cell cycle regulation of folate-mediated one-carbon metabolism. *Wiley Interdiscip. Rev. Syst. Biol. Med.* 10, e1426. <https://doi.org/10.1002/wsbm.1426>.
- McArdle, D.J.T., Gaillard, F, 2020. Pernicious azotaemia? A case series of subacute combined degeneration of the cord secondary to nitrous oxide abuse. *J. Clin. Neurosci.* 72, 277-280. <https://dx.doi.org/10.1016/j.jocn.2019.11.003>.
- Molloy, A.M., Weir, D.G., Kennedy, G., et al., 1990. A new high performance liquid chromatographic method for the simultaneous measurement of S-adenosylmethionine and S-adenosylhomocysteine. Concentrations in pig tissues after inactivation of methionine synthase by nitrous oxide. *Biomed. Chromatogr.* 4, 257-260. <https://doi.org/10.1002/bmc.1130040611>.
- Myles, P.S., Chan, M.T., Kaye, D.M., et al., 2008. Effect of nitrous oxide anesthesia on plasma homocysteine and endothelial function. *Anesthesiology* 109, 657-663. <https://dx.doi.org/10.1097/ALN.0b013e31818629db>.

Nunn, J.F., 1984. Interaction of nitrous oxide and vitamin B12. *Trends Pharmacol. Sci.* 5, 225-227. [https://dx.doi.org/10.1016/0165-6147\(84\)90426-7](https://dx.doi.org/10.1016/0165-6147(84)90426-7).

Oussalah, A., Julien, M., Levy, J., et al., 2019. Global Burden Related to Nitrous Oxide Exposure in Medical and Recreational Settings: A Systematic Review and Individual Patient Data Meta-Analysis. *J. Clin. Med.* 8, 551. <https://dx.doi.org/10.3390/jcm8040551>.

Pratt, D.N., Patterson, K.C., Quin, K., 2020. Venous thrombosis after nitrous oxide abuse, a case report. *J. Thromb. Thrombolysis* 49, 501-503. <https://dx.doi.org/10.1007/s11239-019-02010-9>.

Reynolds, T. M., Brain, A., 1992. A simple internally-standardised isocratic HPLC assay for vitamin B6 in human serum. *J. Liq. Chromatogr. Rel. Technol.* 15, 897-914.

Riedel, B., Fiskerstrand, T., Refsum, H., et al., 1999. Co-ordinate variations in methylmalonyl-CoA mutase and methionine synthase, and the cobalamin cofactors in human glioma cells during nitrous oxide exposure and the subsequent recovery phase. *Biochem. J.* 341, 133-138. <https://dx.doi.org/10.1042/bj3410133>.

Seed, A., Jogia, M., 2020. Lessons of the month: Nitrous oxide-induced functional vitamin B12 deficiency causing subacute combined degeneration of the spinal cord. *Clin. Med. (Lond)* 20, e7-e9. <https://dx.doi.org/10.7861/clinmed.2020-0072>.

Sharma, V.S., Pilz, R.B., Boss, G.R., et al., 2003. Reactions of nitric oxide with vitamin B12 and its precursor, cobinamide. *Biochemistry* 42, 8900-8. <https://dx.doi.org/10.1021/bi034469t>.

Simile, M.M., Latte, G., Feo, C.F., et al., 2018. Alterations of methionine metabolism in hepatocarcinogenesis: the emergent role of glycine N-methyltransferase in liver injury. *Ann. Gastroenterol.* 31, 552-560. <https://dx.doi.org/10.20524/aog.2018.0288>.

Sterman, A.B., Coyle, P.K., 1983. Subacute toxic delirium following nitrous oxide abuse. *Arch. Neurol.* 40, 446-447. <https://dx.doi.org/10.1001/archneur.1983.04050070076021>.

Takahashi-Iñiguez, T., González-Noriega, A., Michalak, C., et al., 2017. Human MMAA induces the release of inactive cofactor and restores methylmalonyl-CoA mutase activity through their complex formation. *Biochimie* 142, 191-196. <https://dx.doi.org/10.1016/j.biochi.2017.09.012>.

Tibbetts, A.S., Appling, D.R., 2010. Compartmentalization of Mammalian folate-mediated one-carbon metabolism. *Annu. Rev. Nutr.* 30, 57-81. <https://dx.doi.org/10.1146/annurev.nutr.012809.104810>.

van Amsterdam, J., Nabben, T., van den Brink, W., 2015. Recreational nitrous oxide use: Prevalence and risks. *Regul. Toxicol. Pharmacol.* 73, 790-796. <https://dx.doi.org/10.1016/j.yrtph.2015.10.017>.

Waclawik, A.J., Luzzio, C.C., Juhasz-Pocsine, K., et al., 2003. Myeloneuropathy from nitrous oxide abuse: unusually high methylmalonic acid and homocysteine levels. *WMJ.* 102, 43-45 (Erratum in: *WMJ.* 2003;102(6):5). PMID: 12967021.

Wei, Y., Lu, M., Mei, M., et al., 2020. Pyridoxine induces glutathione synthesis via PKM2-mediated Nrf2 transactivation and confers neuroprotection. *Nat. Commun.*11:941. <https://dx.doi.org/10.1038/s41467-020-14788-x>.

Xiang, Y., Li, L., Ma, X., et al., 2021. Recreational Nitrous Oxide Abuse: Prevalence, Neurotoxicity, and Treatment. *Neurotox. Res.* 39, 975-985. <https://dx.doi.org/10.1007/s12640-021-00352-y>.

TABLE

Table 1

Clinics, biology, and medical treatments of four patients with N₂O abuse. Abnormal biological data appear in bold underlined characters. NA, not available. * Reference laboratory range values are given between round brackets.

	Patient 1	Patient 2	Patient 3	Patient 4
Age (years)	19	19	18	23
N₂O consumption	200 to 300 whippits/day	200 whippits/day	100 whippits/week-end	100 whippits/day
Clinical signs	gait disorder, paraesthesia of left leg and both hands	gait disorder, paresthesia of lower limbs	gait disorder, walking pain	delirium
Spinal MRI	extensive posterior C2 - C6 cervical cord injury suggestive of subacute combined degeneration	normal	NA	NA
EMG	NA	bilateral L5 root affection more marked on left side	severe symmetrical sensitivomotor axonal polyneuropathy, bilateral carpal tunnel syndrome	NA
<u>Blood/Plasma/Serum Biology*</u>				
Hemoglobin(13-17g/dL)	14.1	<u>11.4</u>	16.1	13.1
MGM (82-98 µm ³)	95.3	<u>88.6</u>	93.3	89.3
Vit B12 (0.2-1 ng/mL)	<u>0.2</u>	<u><0.2</u>	<u>0.2</u>	<u><0.2</u>
Vit B9 (3.1-19.9 ng/mL)	<u>7.5</u>	<u>11.9</u>	<u>4.4</u>	<u>21.3</u>
Vit B6 (15-73 nmol/L)	<u>8</u>	<u>59</u>	<u>47</u>	<u>NA</u>
Vit B1 (95-180 nmol/L)	<u>167</u>	<u>105</u>	<u>145</u>	<u>109</u>
Homocysteine(<14µmol/L)	<u>144</u>	<u>142 - 153</u>	<u>123</u>	<u>164</u>
Methionine (16-29 µmol/L)	<u>22</u>	<u>19.5</u>	<u>19</u>	<u>NA</u>
Sarcosine (0.5-2.7 µmol/L)	<u>3.5</u>	<u>2.1</u>	<u>7.7</u>	<u>NA</u>
Cystathionine (0.1-1.5µmol/L)	<u>1.7</u>	<u>2.4</u>	<u>1.7</u>	<u>NA</u>
<u>Therapeutic measures</u>				
Cessation of N ₂ O intakes	Yes	Yes	Yes	Yes
Vit B12	1 mg/day for 1 month	1 mg/day for 1 month	1 mg/day (for 2 weeks before loss of follow-up)	
Vit B9 (folate)	5 mg/day for 1 month	5 mg x2/day for 1 month	None	
Vit B6	500 mg/day for 1 month	None	None	

FIGURES




Fig. 1. MRI examination of the cervical spine in patient 1. MRI images show the swelling from C1 to C5 and the strong signal intensity in the dorsal cervical spinal cord on the sagittal (A) and axial (B) T2-weighted images. The contrast-enhanced T1-weighted axial image (C) further highlights the focal enhancement of the left posterior column of the spinal cord. These findings are consistent with subacute combined degeneration of the spinal cord, and possibly with a deficient function of S-adenosylmethionine-dependent methylases involved in myelogenesis (see Discussion).




Fig.2. Subcellular location and metabolic role of cellular pathways affected by N_2O . The vitamin B12-dependent methionine synthase (cytosol) and methylmalonyl-CoA mutase (mitochondria) are biological metabolic targets of N_2O . As explained in the main article text, methionine synthase is differently and more severely impacted than methylmalonyl-CoA mutase. The figure highlights the cellular roles of these enzymes in major cellular pathways which include folate cycle, methylation cycle, S-adenosylmethionine-dependent methyltransferases, transulfuration pathway, Krebs cycle fueling by methylmalonyl-CoA, and glutathione biosynthesis. A non-exhaustive description of proteins, metabolic steps, and intermediates involved in these pathways is given. Abbreviations are: BHCMT, betaine homocysteine methyltransferase; DMGDH, dimethylglycine dehydrogenase; GAMT, guanidinoacetate methyltransferase; GCS, glycine cleavage system; GNMT, glycine *N*-methyltransferase; MFT, mitochondrial folate transporter; MTHFR, methylenetetrahydrofolate reductase; PNMT, phenylethanolamine *N*-methyltransferase; SARDH, sarcosine dehydrogenase; SHMT, serine hydroxymethyltransferase; THF, tetrahydrofolate (yielded from folic acid, not shown). Note that serine produced by mitochondrial SHMT may fuel serine consumed by cytosolic SHMT, and glycine produced by cytosolic SHMT may fuel that consumed by mitochondrial SHMT. The purple broken lines stress some mutual inhibitions that can take place between the methylation and folate cycles.