



HAL
open science

Efficiency of four solutions in removing 23 conventional antineoplastic drugs from contaminated surfaces.

Nicolas Simon, Nicolas Guichard, Pascal Odou, Bertrand Décaudin, Pascal Bonnabry, Sandrine Fleury-Souverain

► To cite this version:

Nicolas Simon, Nicolas Guichard, Pascal Odou, Bertrand Décaudin, Pascal Bonnabry, et al.. Efficiency of four solutions in removing 23 conventional antineoplastic drugs from contaminated surfaces.. PLoS ONE, 2020, PLoS One, 15, pp.e0235131. 10.1371/journal.pone.0235131 . hal-04610861

HAL Id: hal-04610861

<https://hal.univ-lille.fr/hal-04610861v1>

Submitted on 13 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Efficiency of four solutions in removing 23 conventional antineoplastic drugs from contaminated surfaces

Nicolas Simon^{1,2*}, Nicolas Guichard¹, Pascal Odou², Bertrand Decaudin^{1,2}, Pascal Bonnabry¹, Sandrine Fleury-Souverain¹

1 Pharmacy, Geneva University Hospitals and Institute of Pharmaceutical Sciences of Western Switzerland, University of Geneva, Geneva, Switzerland, **2** Univ. Lille, CHU Lille, ULR 7365 –GRITA–Groupe de Recherche sur les Formes Injectables et les Technologies Associées, Lille, France

* nicolas.simon@univ-lille.fr



Abstract

Background

Residual contamination by intravenous conventional antineoplastic drugs (ICAD) is still a daily issue in hospital facilities. This study aimed to compare the efficiency (Eff_Q) of 4 different solutions to remove 23 widely used ICADs from surfaces.

Method and findings

A solution containing 23 ICADs (4 alkylating agents, 8 antimetabolites, 2 topo-I inhibitors, 6 topo-II inhibitors and 3 spindle poisons) was spread over 100 cm² stainless steel. After drying, decontamination was carried out using 10×10 cm wipes moistened with 300 μL of one of the following solutions: 70% isopropanol (S1); ethanol-hydrogen peroxide 91.6–50.0 mg/g (S2); 10⁻² M sodium dodecyl sulphate/isopropanol 80/20 (S3) or 0.5% sodium hypochlorite (S4). Six tests were performed for each decontamination solution. Two modalities were tested: a single wipe motion from top to bottom or vigorous wiping (n = 6 for each modality). Residual contamination was measured with a validated liquid chromatography with tandem mass spectrometry detection method. Solution efficiency (in %) was computed as follows: $Eff_Q = 1 - (\text{quantity after decontamination} / \text{quantity before decontamination})$, as median (min–max) for the 23 ICADs. The overall decontamination efficiency (Eff_Q) of the 4 solutions was compared by a Kruskal-Wallis test. Decontamination modalities were compared for each solution and per ICAD with a Mann-Whitney test (p<0.05).

Eff_Q were significantly different from one solution to the next for single wipe motion decontamination: 79.9% (69.3–100), 86.5% (13.0–100), 85.4% (56.5–100) and 100% (52.9–100) for S1, S2, S3 and S4 (p<0.0001), respectively. Differences were also significant for vigorous decontamination: Eff_Q of 84.3% (66.0–100), 92.3% (68.7–100), 99.6% (84.8–100) and 100% (82.9–100) for S1, S2, S3 and S4, respectively (p<0.0001). Generally, vigorous decontamination increased Eff_Q for all tested solutions and more significantly for the surfactant.

OPEN ACCESS

Citation: Simon N, Guichard N, Odou P, Decaudin B, Bonnabry P, Fleury-Souverain S (2020) Efficiency of four solutions in removing 23 conventional antineoplastic drugs from contaminated surfaces. PLoS ONE 15(6): e0235131. <https://doi.org/10.1371/journal.pone.0235131>

Editor: Girish Sailor, Bhagwan Mahvir College of Pharmacy, INDIA

Received: March 2, 2020

Accepted: June 8, 2020

Published: June 22, 2020

Copyright: © 2020 Simon et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: NS received a grant from French Society of Clinical Pharmacy, by la Fondation pour la Recherche en Pharmacie Hospitalière, AstraZeneca laboratories for his post-doctoral position in the University Hospital of Geneva. The specific roles of this author is articulated in the 'author contributions' section. No additional external funding was received for this study. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have read the journal's policy and the authors of this paper have the following competing interests: NS received a grant from AstraZeneca for his post-doctoral position in the University Hospital of Geneva. There are no patents, products in development or marketed products associated with this research to declare. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Conclusion

Decontamination efficiency depended on the solution used but also on the application modality. An SDS admixture seems to be a good alternative to sodium hypochlorite, notably after vigorous chemical decontamination with no hazard either to materials or workers.

Introduction

Occupational exposure to intravenous conventional antineoplastic drugs (ICADs) is a daily problem widely encountered in care settings. Traces of contamination have been found in many sectors of the drug supply chain, from industry [1] to hospitals [2,3] and to patients' homes [4]. More recently, veterinary clinics have also been found liable to occupational exposure [5].

Since the first report [6], numerous articles have highlighted different risks, varying according to the case. For example, sudden massive contamination (e.g. vial breaking) may cause acute symptomatology [7–9], which is an uncommon event. Chronic exposure could lead to clinical or biological disorders, notably reproductive risks or cytogenetic effects [10,11].

Rapidly, professional recommendations were published [12–14], followed by institutional recommendations [15–21]. All these recommendations insist on the need to combine several protective measures. Indeed, personal protective equipments (e.g. gloves, gowns, mask. . .) are recommended to avoid a direct contact between the operator and the chemical hazard. Such equipments have to be combined to collective equipments (e.g. compounding isolators or laminar air-flow hoods, specific sterile medical devices for preparation or administration. . .) which are devoted to decrease the widepreading of the chemical contamination to the surrounding [19,21,22,23]. Nonetheless, the respect for these recommendations is not absolute and appears to differ among healthcare professionals [24]. Moreover, the use of protective tools varies according to healthcare facilities. Consequently, healthcare workers are exposed to varying levels and types of contamination.

One safety measure which must be developed is the elimination of contaminants from workplaces. This was recently mentioned in the <USP800> monograph and in the latest recommendations of the American Society of Healthcare-system Pharmacists [23,25]. Removing contamination from surfaces has been studied for more than thirty years. A recent review of the literature discussed all the decontamination strategies studied and published [26] and in what conditions: contaminants used, contamination level, surface type and operating procedure [26]. According to the <USP800>, contaminants may be removed from surfaces either by deactivation, that is to say by chemical degradation, or by decontamination, involving a physical process (i.e. desorption). Desorption strategies were studied on different antineoplastic drugs after intentional surface contamination. Some studies focused on 1 or 2 drugs [27–32], testing solutions of different natures (quaternary ammonium, biguanide, alcohol-based solutions). Others were performed on 10 drugs with a wider range of solutions (organic solvent; anionic, non-ionic or cationic surfactants; oxidant or water) [33]. Considering the different physicochemical properties of ICADs, some strategies were found to be of little efficacy (e.g. 70% isopropanol). Nonetheless, certain approaches were promising such as diluted oxidant (e.g. sodium hypochlorite) or the combination of a surfactant agent and an alcoholic solvent (e.g. sodium dodecyl sulfate/isopropanol). Even if these studies were tested on 10 compounds and have yielded interesting results, they concern only a small number of the

antineoplastic drugs handled in hospital. Therefore, efficacy on numerous other drugs from different antineoplastic drug families requires evaluating.

The objective of this *in vitro* study was to assess the efficiency of four different solutions after intentional contamination on stainless steel surfaces. The originality of this study lies in the assessment of decontamination solutions and application modalities on 23 conventional antineoplastic drugs from all conventional antineoplastic drug families.

Materials and methods

Chemicals

The ICADs used for experiments were: 5-fluorouracil (batch# PE4SH-RM), methotrexate (batch# G07UG-01) and dacarbazine (batch# Z3J8O-SF), purchased from Tokyo Chemical Industry (Zwijndrecht, Belgium); gemcitabine (batch# A0375170), purchased from Acros Organic (Geel, Belgium); busulfan (batch# BCBN8120V), purchased from Sigma-Aldrich (Buchs, Switzerland); ganciclovir, purchased from Roche Pharma (Cymevene[®], batch# B4091B08, Reinach, Switzerland); cytarabine (batch# 3-YFD-59-1), epirubicin (batch# 11-CGS-118-1) and topotecan (batch# 8-MSW-162-1) purchased from Toronto Research Chemicals (North York, ON, Canada); raltitrexed (batch# MG10453-29102016), pemetrexed (batch# AGN2017-685), docetaxel (batch# CS13527-18112016), paclitaxel (batch# CS22539-20112016), vincristine sulfate (batch# MC-10452-0212016), doxorubicin hydrochloride (batch# MC10454-0612016), daunorubicin (batch# MC10456-1712016), idarubicin (batch# 5-CGS-96-1), etoposide (batch# MC10457-20102016), etoposide phosphate (batch# MC305547-21082017), irinotecan hydrochloride (batch# MC20783-21082017) and fludarabine phosphate (batch# MC444607-21082017) brought from Pharmaserv (Santsstad, Switzerland). Cyclophosphamide (batch# 7F124) and ifosfamide (batch# 7H025) were obtained from Baxter EG (Endoxan[®] and Holoxan[®], respectively, Opfikon, Switzerland). Internal standards (IS) for the analytical assay ($[^{13}\text{C}^{15}\text{N}_2]$ -fluorouracil, batch# JA-ALS-16-130P3; $[^{13}\text{C}^2\text{H}_3]$ -methotrexate, batch# TM-ALS-12-134-B1; $[^{13}\text{C}_6]$ -irinotecan, batch# TM-ALS-12-257-P1; $[^2\text{H}_8]$ -cyclophosphamide, batch# LSG-ALS-12-017-P1; and $[^2\text{H}_5]$ -paclitaxel, batch# TF-ALS-11-066-P1) were purchased from Alsachim (Strasbourg, France).

All solvents were mass spectrometry grade and all chemicals were obtained in the highest available analytical quality. Dimethylsulfoxide (DMSO) was purchased from Sigma-Aldrich (Buchs, Switzerland). Ammonium hydroxide, acetic acid and acetonitrile were obtained from Merck (Darmstadt, Germany). Ultrapure water (UPW) was obtained from a Milli-Q purification system from Millipore (Bedford, MA, USA).

Preparation of working solutions

Standard stock solutions were prepared by diluting standard compounds in DMSO or UPW for pemetrexed at a concentration of 1 mg/mL and were stored at -80°C . One stock solution was prepared with:

- Group A ($C_A = 10,000$ ng/mL for each compound) comprised of 5-fluorouracil, busulfan, cyclophosphamide, cytarabine, dacarbazine, docetaxel, etoposide, etoposide phosphate, ganciclovir, gemcitabine, idarubicin, ifosfamide, methotrexate, paclitaxel, pemetrexed and raltitrexed;
- Group B ($C_B = 50,000$ ng/ml for each compound) comprised of daunorubicin, doxorubicin, epirubicin, topotecan and vincristine;
- Group C comprised of only fludarabine at a concentration of $C_C = 100,000$ ng/ml.

Internal standard stock solutions were prepared by diluting individual radiolabelled IS in DMSO at 0.1 mg/mL and also stored at -80°C.

Contamination of surfaces

Contamination was carried out with 50 µL of a 1/50th dilution of the stock solution. Thus, the intentional contaminations were 200 ng, 1,000 ng and 2,000 ng for groups A, B and C, respectively.

The 50 µL was randomly dropped on a 10×10 cm stainless steel surface and allowed to dry for 1h under a ventilated biosafety cabinet protected from light.

Decontamination of surfaces

Decontamination followed 2 modalities: a single wipe motion from top to bottom or vigorous wiping.

Standard application consisted in a single wipe of surfaces from top to bottom.

Vigorous decontamination consisted in scrubbing surfaces energetically several times for a few seconds.

Decontamination was carried out using 10×10 cm wipes (Texwipe™ 3210, ITW Texwipe, Kenersville, NC, USA) moistened with 300 µL decontamination solution. Four different solutions were tested: Solution 1: 70% isopropanol (Klercide™, Ecolab, Farmham, UK); Solution 2: ethanol (91.6 mg/g) hydrogen peroxide (50.0 mg/g) (Anioxyspray™, Anios, Hellemmes, France); Solution 3: 10⁻² M sodium dodecyl sulfate:isopropanol 80:20 (home-made sterile solution) and Solution 4: 0.5% sodium hypochlorite, obtained by diluting a 3% marketed solution (Hänseler, Herisau, Switzerland) in UPW.

Residual contamination measurement and decontamination assessment

Residual contamination was measured by applying a liquid chromatography with tandem mass spectrometry detection method developed and validated specifically to estimate surface contamination by the tested contaminants [34–36].

Briefly, sampling was performed using a polyester swab (Texwipe™ 716, ITW Texwipe, Kernersville, NC, USA). Swabs were humidified with 100 µL of 75% IPA (50 µL/side) for wipe sampling [36], then introduced into 12-mL amber glass tubes containing 2-mL of desorption solution (10 mM acetic + 2% acetonitrile) and the 5 internal standards. The tube was then vortexed for a few seconds. The aqueous solution was analysed according to a previously published method [34,35].

Statistics

All experiments were repeated 6 times and were conducted by a single operator to limit inter-individual variability of the results.

Samples were considered contaminated if at least one contaminant was quantified. For a few samples in which contaminants were unquantifiable because the signal ranged between the LOD and the LOQ, the value was expanded to the LOQ.

The contamination rate was considered to be the proportion of contaminated samples. The ability of the decontamination solution to remove chemical contamination was estimated by decontamination efficiency, defined as follows [37]:

$$\text{Eff}_q = 1 - \frac{\text{quantity after decontamination procedure}}{\text{quantity before decontamination procedure}} \quad \text{for one contaminant} \quad \text{Eq 1}$$

$$\text{Eff}_Q = \frac{\sum \text{Eff}_q}{n} \quad \text{for all contaminants} \quad \text{Eq 2}$$

Efficiency results are presented as medians [min–max], expressed in %. Non-parametric tests were used because of the number of samples ($n < 30$) and normality was unverifiable. The comparison of the overall efficiency (Eff_Q) of the four solutions was performed with an ANOVA on ranks according to the Kruskal-Wallis method ($P < 0.05$). When this analysis revealed a significant p-value, contrasts were obtained with the Conover–Iman test on ranks to detect significant differences between couples of solution. The Bonferroni correction was applied on P-values to limit interpretation bias due to repeated tests. The two decontamination modalities were compared for each solution by a non parametric Mann-Whitney test ($p < 0.05$).

Statistics were performed in using XLSTAT[®] for Excel[®] (Addinsoft, Paris, France). Figures were drawn using Excel[®] (Microsoft, Paris, France).

Results

Standard decontamination

The highest overall efficiency was observed with sodium hypochlorite. The medians [min–max] of overall Eff_Q were 79.9% [69.3–100]; 86.5% [13–100]; 85.4% [56.5–100] and 100% [52.9–100] for solutions 1, 2, 3 and 4, respectively. The Eff_q for each antineoplastic drug by solution are summarize in [Table 1](#).

The lowest Eff_q were observed for etoposide (69.3%), docetaxel (44.8%), etoposide (56.5%) and docetaxel (52.9%) for solutions 1, 2, 3, and 4, respectively. The highest Eff_q was 100.0%, reached for 4, 6, 9 and 14 contaminants for solutions 1, 2, 3 and 4, respectively ([Fig 1](#)). The four solutions reached very good Eff_q for pemetrexed, daunorubicin, etoposide phosphate. As well as 100% efficiency, Eff_q reached at least 90% for 2 (ganciclovir and topotecan) and 6 contaminants (ganciclovir, raltitrexed, doxorubicin, idarubicin, topotecan and vincristine) for solutions 2 and 3, respectively. Inversely, solution 4 did not reach this threshold for 3 contaminants (cyclophosphamide, ifosfamide and docetaxel).

Vigorous decontamination

[Table 2](#) summarizes the Eff_q for each antineoplastic drug by decontamination solution. The highest overall efficiency was observed with sodium hypochlorite. The observed Eff_Q medians were 100%, 99.6%, 92.3% and 84.3% for S4, S3, S2 and S1, respectively. Residual contamination could be observed more frequently with solutions 1 and 2.

Eff_q comparison of the four solutions using the Kruskal-Wallis test revealed some significant differences for doxorubicin (S1 = 66.0%; S2 = 95.3% vs. S3 = 100%; S4 = 100%; $p = 0.011$), epirubicin (S1 = 74.4%; S2 = 83.2% vs. S3 = 100%; S4 = 100%; $p = 0.001$), ganciclovir (S1 = 84.6%; S2 = 93.6% vs. S3 = 100%; S4 = 100%; $p = 0.001$), irinotecan (S1 = 82.5%; S2 = 89.3% vs. S3 = 100%; S4 = 100%; $p = 0.001$), methotrexate (S1 = 76.4%; S2 = 86.2% vs. S4 = 97.7%; $p = 0.026$), raltitrexed (S1 = 94.5%; S2 = 92.0% vs. S3 = 100%; S4 = 100%; $p = 0.001$) and topotecan (S1 = 92.9%; S2 = 92.3% vs. S3 = 99.6%; S4 = 100%; $p = 0.004$). For these contaminants, the Conovar-Iman analysis revealed contrasts between couples of solutions: solutions 3 and 4 versus solutions 1 and 2. A significant difference was observed on etoposide between the four solutions (S1 = 86.3% vs. S2 = 96.6% vs. S3 = 91.2%; S4 = 100%; $p = 0.001$). Finally, solution 4 had a greater effect than other solutions on dacarbazine (S1 = 84.3%; S2 = 93.8%; S3 = 92.5% vs. S4 = 100%; $p = 0.002$).

Table 1. Efficiency of the decontamination solutions on the 23 tested antineoplastic drugs on stainless steel surfaces after standard single motion. Solution 1 (S1): 70% isopropanol; Solution 2 (S2): ethanol (91.6 mg/g) hydrogen peroxide (50.0 mg/g); Solution 3 (S3): 10^{-2} M sodium dodecyl sulfate:isopropanol 80:20; Solution 4 (S4): 0.5% sodium hypochlorite aqueous solution. 5FU: 5-fluorouracil; Cyta: cytarabine; Fluda: fludarabine; Ganci: ganciclovir; Gem: gemcitabine; Mtx: methotrexate; Peme: pemetrexed; Ralti: raltitrexed; Busu: busulfan; Cyc: cyclophosphamide; Ifos: ifosfamide; Dacar: dacarbazine; Dauno: daunorubicin; Doxo: doxorubicin; Ida: idarubicin; Epi: epirubicin; EtopoP: etoposide phosphate; Eto: etoposide; Dtx: docetaxel; Pcx: paclitaxel; Irino: irinotecan; Topo: topotecan; Vin: vincristine.

	Eff _q	5FU	Cyta	Fluda	Ganci	Gem	Mtx	Peme	Ralti
S1	m	79.9%	73.5%	86.1%	83.5%	73.6%	85.8%	100.0%	86.1%
	sd	36.1%	37.3%	26.8%	27.5%	37.4%	23.4%	0.0%	27.7%
S2	m	100.0%	76.1%	88.1%	94.0%	75.5%	85.7%	100.0%	100.0%
	sd	0.0%	25.3%	15.3%	14.6%	28.5%	13.4%	0.0%	0.0%
S3	m	100.0%	71.5%	81.0%	98.4%	71.0%	85.4%	100.0%	100.0%
	sd	0.0%	32.4%	31.4%	1.5%	39.4%	18.7%	0.0%	0.0%
S4	m	100.0%	98.5%	94.4%	100.0%	99.2%	100.0%	100.0%	100.0%
	sd	0.0%	2.6%	8.8%	0.0%	1.9%	0.0%	0.0%	0.0%
	Eff _q	Busu	Cyc	Ifos	Dacar	Dauno	Doxo	Ida	Epi
S1	m	73.0%	77.5%	76.9%	79.7%	100.0%	75.4%	82.9%	82.5%
	sd	52.4%	34.5%	35.6%	34.3%	0.0%	30.1%	20.7%	16.6%
S2	m	86.5%	84.4%	83.7%	81.3%	100.0%	88.0%	66.7%	49.1%
	sd	28.6%	21.1%	22.8%	27.9%	0.0%	15.5%	57.7%	76.7%
S3	m	77.5%	79.6%	77.1%	74.9%	100.0%	100.0%	100.0%	82.9%
	sd	52.0%	31.8%	38.1%	41.6%	0.0%	0.0%	0.0%	35.8%
S4	m	100.0%	81.7%	77.5%	99.5%	100.0%	100.0%	100.0%	100.0%
	sd	0.0%	26.8%	35.6%	1.3%	0.0%	0.0%	0.0%	0.0%
	Eff _q	EtopoP	Eto	Dtx	Pcx	Irino	Topo	Vin	
S1	m	100.0%	69.3%	81.1%	78.4%	78.8%	72.8%	100.0%	
	sd	0.0%	49.7%	31.4%	34.5%	35.8%	41.0%	0.0%	
S2	m	100.0%	87.1%	44.8%	64.7%	72.1%	92.6%	100.0%	
	sd	0.0%	19.5%	103.9%	42.3%	37.5%	16.6%	0.0%	
S3	m	100.0%	56.5%	85.8%	63.3%	78.0%	100.0%	100.0%	
	sd	0.0%	74.7%	21.0%	80.1%	38.7%	0.0%	0.0%	
S4	m	100.0%	100.0%	52.9%	86.7%	99.4%	100.0%	100.0%	
	sd	0.0%	0.0%	87.1%	18.7%	1.4%	0.0%	0.0%	

<https://doi.org/10.1371/journal.pone.0235131.t001>

In this context, decontamination efficiency also depended on the tested contaminant (Fig 2). The lowest Eff_q were observed for doxorubicin (66.0%), paclitaxel (68.7%), cytarabine (84.8% and 82.9%) for solutions 1, 2, 3, and 4, respectively. The highest Eff_q (i.e. 100%) was observed on 5, 1, 11 and 13 contaminants for solutions 1, 2, 3, and 4, respectively. An Eff_q of 100% was observed on etoposide for the four tested solutions. Solution 1 reached an Eff_q ≥ 90% for fludarabine, raltitrexed and topotecan. For solution 2, this threshold was attained except for cytarabine, gemcitabine, methotrexate, epirubicin, docetaxel and paclitaxel. The Eff_q for solution 3 was lower than 90% for cytarabine, methotrexate and idarubicin. In the case of solution 4, this threshold was not reached for cytarabine and gemcitabine, nor for docetaxel and paclitaxel. Both cyclophosphamide and ifosfamide were decontaminated with an Eff_q of 89.8 and 89.9%, respectively.

Comparison of decontamination modalities

After standard application of the solutions (Fig 3), a significant difference was observed for solution 4 compared to the others ($p < 0.001$) according to the Mann-Whitney test. After vigorous application, Eff_q were 84.3% [66.0–100]; 92.3% [68.7–100]; 99.6% [84.8–100] and 100%




Fig 1. Comparison of decontamination efficiency per contaminant of four decontamination or deactivation solutions after standard single motion. Values represented are median Eff_q for each drug (n = 6). 5FU: 5-fluorouracil; Cyta: cytarabine; Fluda: fludarabine; Ganci: ganciclovir; Gem: gemcitabine; Mtx: methotrexate; Peme: pemetrexed; Ralti: raltitrexed; Busu: busulfan; Cyc: cyclophosphamide; Ifos: ifosfamide; Dacar: dacarbazine; Dauno: daunorubicin; Doxo: doxorubicin; Ida: idarubicin; Epi: epirubicin; EtopoP: etoposide phosphate; Eto: etoposide; Dtx: docetaxel; Pcx: paclitaxel; Irino: irinotecan; Topo: topotecan; Vin: vincristine. Blue line/circles: 70% isopropanol; red line/triangles: admixture of ethanol-hydrogen peroxide (91.6–50.0 mg/g); green line/diamonds: admixture of 10⁻² M sodium dodecyl sulfate/isopropanol (80/20) and yellow line/squares: sodium hypochlorite. * Significant difference for solution 4 over other solutions; ** Significant difference between solutions 4 and 2; \$ significant difference between solutions 4 and 1 and between \$^s solutions 4 and 3. + significant difference for both solutions 3 and 4 compared to others.

<https://doi.org/10.1371/journal.pone.0235131.g001>

[82.9–100] for solutions 1, 2, 3 and 4, respectively (Fig 3) with a significant difference for both solutions 3 and 4 compared to solutions 1 and 2 ($p < 0.001$). Vigorous application significantly increased the Eff_Q of solution 3 compared to the standard application ($p = 0.007$). This result was not noted for the other solutions.

Discussion

Many products have been marketed for use in compounding units to remove chemical contamination by ICADs. Despite these, contamination is still present. [2,38]. In compounding units, both containment primary and secondary engineering controls (respectively C-PEC and C-SEC) are contaminated, which means that safety measures have to be improved. As mentioned in the <USP800> monograph, the removal of chemical contaminants may be obtained

Table 2. Efficiency of the decontamination solutions on the 23 tested antineoplastic drugs on stainless steel surfaces after vigorous decontamination. Solution 1 (S1): 70% isopropanol; Solution 2 (S2): ethanol (91.6 mg/g) hydrogen peroxide (50.0 mg/g); Solution 3 (S3): 10^{-2} M sodium dodecyl sulfate:isopropanol 80:20; Solution 4 (S4): 0.5% sodium hypochlorite aqueous solution. 5FU: 5-fluorouracil; Cyta: cytarabine; Fluda: fludarabine; Ganci: ganciclovir; Gem: gemcitabine; Mtx: methotrexate; Peme: pemetrexed; Ralti: raltitrexed; Busu: busulfan; Cyc: cyclophosphamide; Ifos: ifosfamide; Dacar: dacarbazine; Dauno: daunorubicin; Doxo: doxorubicin; Ida: idarubicin; Epi: epirubicin; EtopoP: etoposide phosphate; Eto: etoposide; Dtx: docetaxel; Pcx: paclitaxel; Irino: irinotecan; Topo: topotecan; Vin: vincristine.

	Eff _q	5FU	Cyta	Fluda	Ganci	Gem	Mtx	Peme	Ralti
S1	m	100.0%	74.9%	90.3%	84.6%	76.4%	76.4%	100.0%	94.5%
	sd	0.0%	13.5%	10.4%	9.6%	14.9%	29.6%	0.0%	4.2%
S2	m	95.9%	83.0%	98.2%	93.6%	85.2%	86.2%	92.3%	92.0%
	sd	9.1%	11.8%	4.1%	6.3%	8.8%	8.8%	4.9%	4.1%
S3	m	100.0%	84.8%	100.0%	100.0%	91.5%	87.9%	100.0%	100.0%
	sd	0.0%	8.1%	0.0%	0.0%	7.3%	11.4%	0.0%	0.0%
S4	m	100.0%	82.9%	91.4%	100.0%	88.2%	97.7%	100.0%	100.0%
	sd	0.0%	10.7%	13.7%	0.0%	7.0%	3.5%	0.0%	0.0%
	Eff _q	Busu	Cyc	Ifos	Dacar	Dauno	Doxo	Ida	Epi
S1	m	87.2%	80.6%	81.9%	84.3%	100.0%	66.0%	72.8%	74.4%
	sd	10.8%	13.3%	12.9%	13.6%	0.0%	26.0%	26.1%	14.3%
S2	m	91.5%	94.9%	93.5%	93.8%	97.8%	95.3%	75.8%	83.2%
	sd	8.6%	3.7%	4.5%	5.7%	5.4%	7.8%	12.6%	15.1%
S3	m	94.6%	93.7%	93.7%	92.5%	100.0%	100.0%	86.1%	100.0%
	sd	5.9%	3.5%	3.6%	4.2%	0.0%	0.0%	31.0%	0.0%
S4	m	94.8%	89.8%	89.9%	100.0%	100.0%	100.0%	91.0%	100.0%
	sd	5.9%	7.0%	7.7%	0.0%	0.0%	0.0%	20.1%	0.0%
	Eff _q	EtopoP	Eto	Dtx	Pcx	Irino	Topo	Vin	
S1	m	100.0%	86.3%	82.6%	69.9%	82.5%	92.9%	100.0%	
	sd	0.0%	9.4%	17.9%	31.8%	9.1%	9.4%	0.0%	
S2	m	100.0%	96.6%	74.0%	68.7%	89.3%	92.3%	95.9%	
	sd	0.0%	2.7%	20.3%	62.8%	8.9%	6.1%	10.1%	
S3	m	100.0%	91.2%	97.9%	99.0%	100.0%	99.6%	100.0%	
	sd	0.0%	6.4%	3.0%	2.5%	0.0%	0.5%	0.0%	
S4	m	100.0%	100.0%	85.9%	84.8%	100.0%	100.0%	100.0%	
	sd	0.0%	0.0%	16.1%	19.1%	0.0%	0.0%	0.0%	

The four solutions yielded the same decontamination efficiency for cytarabine, daunorubicin, vincristine and 5-fluorouracil. Oxazophosphorines were removed more efficiently, notably with solutions 2, 3 and 4. Solution 3 had better Eff_q on both docetaxel and paclitaxel (Fig 2).

<https://doi.org/10.1371/journal.pone.0235131.t002>

by two means: deactivation or decontamination [25]. Among the solutions tested in this study, sodium hypochlorite and hydrogen peroxide can deactivate conventional antineoplastic drugs as they are good oxidisers. As the oxidation of contaminants may generate degradation compounds, these were not detected by our assay and probably persist on the surface. Moreover, such oxidants may also act on the surfaces of isolators or BSC and may degrade them over time.

The originality of this study lies in the range of drugs studied whereas in literature the majority have been limited to few markers, sometimes up to 10 contaminants in the same contamination [26]. New data is therefore accessible even as regards oxidisers that have been studied thoroughly in the past.

Our study confirms that sodium hypochlorite has good overall efficiency compared to other solutions after standard application, especially on antimetabolites, anthracyclines, topo-I and topo-II inhibitors. These results are consistent with previously published results [27,29,33]. Indeed, 0.5% sodium hypochlorite was successfully tested on stainless steel surfaces




Fig 2. Comparison of the decontamination efficiency per contaminant of four decontamination or deactivation solutions after vigorous decontamination. Values represented are median Eff_Q for each drug ($n = 6$). 5FU: 5-fluorouracil; Cyta: cytarabine; Fluda: fludarabine; Ganci: ganciclovir; Gem: gemcitabine; Mtx: methotrexate; Peme: pemetrexed; Ralti: raltitrexed; Busu: busulfan; Cyc: cyclophosphamide; Ifos: ifosfamide; Dacar: dacarbazine; Dauno: daunorubicin; Doxo: doxorubicin; Ida: idarubicin; Epi: epirubicin; EtopoP: etoposide phosphate; Eto: etoposide; Dtx: docetaxel; Pcx: paclitaxel; Irino: irinotecan; Topo: topotecan; Vin: vincristine. Blue line/circles: 70% isopropanol; red line/triangles: admixture of ethanol-hydrogen peroxide (91.6–50.0 mg/g); green line/diamonds: admixture of 10^{-2} M sodium dodecyl sulfate/isopropanol (80/20) and yellow line/squares: sodium hypochlorite. + significant difference for both solutions 3 and 4 compared to others; @ significant differences between the 4 solutions; * significant difference of solution 4 over other solutions.

<https://doi.org/10.1371/journal.pone.0235131.g002>

on a miscellaneous of 10 antineoplastic drugs with an Eff_Q of 97.5% [33]. In the other studies, Adé et al., and Hon et al. have tested only cyclophosphamide, with an Eff_Q of 99.74% and 96.62%, respectively. Our result observed on paclitaxel shows that 0.5% sodium hypochlorite may be efficient, as it was previously demonstrated by Lee et al [31]. In this study, the lowest effect is observed on taxanes, with a marked effect on docetaxel, which explains the difference in the Eff_Q range comparatively to the study of Queruau-Lamerie et al.

This study also confirms that 70% isopropanol is a poor decontaminating agent, as has already been observed [33]. In a previous literature review, the Eff_Q of 70% isopropanol on stainless steel surfaces reached a mean of 80.6%, [26] compared with 79.9% in our study. Its decontamination effect on taxanes is higher than NaOCl, probably because it is an organic solvent capable of removing these contaminants by physicochemical affinity.

Two other decontamination solutions were tested. Marketed biocide combining an oxidiser and ethanol proved to be a poor chemical decontaminant. Indeed, if a threshold of 90% is




Fig 3. Comparison of the overall efficiency of four decontamination or deactivation solutions according to their application modalities. White boxes represent standard decontamination and striped boxes represent vigorous decontamination. The solutions compared are 70% isopropanol (IPA), an admixture of ethanol-hydrogen peroxide (91.6–50.0 mg/g), an admixture of 10^{-2} M sodium dodecyl sulfate/isopropanol (80/20) and sodium hypochlorite (NaOCl). Significant differences are observed for single standard motion between NaOCl and the other solutions and for vigorous application for both NaOCl and SDS/IPA compared to the two other solutions.

<https://doi.org/10.1371/journal.pone.0235131.g003>

supposed to define a decontaminant enough efficient [39], the solution 2 (admixture of EtOH/H₂O₂) reached such a value for only 8 contaminants. This solution was chosen because it is used in practice as a disinfectant in many compounding units. Solution 2 was effectively previously tested in a real-life study and compared to solution 3, showing a lower efficiency [40]. As the formulation of disinfectants can vary, our results should not be generalised. It is however advisable to test whether a disinfectant is a good chemical decontaminant before using it [39].

The admixture of SDS/IPA tested here was previously assessed [28,33] and tested on different drugs, demonstrating a difference in Eff_Q depending on contaminant as was also observed in our study. In the study of Queruau-Lamerie et al., such solution reached an Eff_Q of 87.5% on stainless steel surfaces [33], closed from our results (i.e. 85.4%) and also shown variable Eff_Q depending on the contaminant.

Vigorous application tended to increase the Eff_Q for all solutions, but the effect was statistically significant for only solution 3. Therefore, both disinfectant solutions proved to be the worst chemical decontaminants when compared to SDS/IPA and NaOCl in this particular case. These two solutions had comparable Eff_Q after vigorous application. The biggest difference between the two solutions was certainly due to the greater effect of SDS/IPA on the two taxanes, indicating the advantage of surfactant solutions in removing lipophilic and hydrophilic drugs in the same action. A significant effect of vigorous application was found only for SDS/IPA in this study, implying the importance of application modalities (i.e. the human

factor) in the effectiveness of the decontamination process. This seems logical as the contaminants to be removed have to be incorporated into the formed micelles and this requires a mechanical action which may indicate variability in decontamination efficacy between operators: variability in the strength and energy applied during the decontamination process. SDS/IPA solution was previously tested during two studies in real conditions [37,40]. Among the differences between these two studies, a human factor was suspected to explain the difference in the results, notably by the application modalities of the decontamination solution [26]. It is therefore important to consider the training of the operators involved in decontamination to limit this potential interindividual factor as much as possible.

The handling of ICADs necessitates both sterility [41] and safety in care facilities [21]. The <USP800> monograph distinguished two situations relative to duality in the context of sterile compounding: removing microorganisms with a cleaning or a disinfectant solution and removing chemical contaminants using a deactivation or a decontamination solution [25]. In medical wards, this problem has to be analysed in just the same way, as antimicrobial solutions are used there but their decontamination potential is not known. In a previous work, the performance of decontamination solutions was assessed according to three characteristics: overall efficiency (Eff_Q), the number of tested contaminants and the risk of using a decontamination solution [39]. For routine decontamination, two solutions among those tested here have been shown to be the best decontaminants against the whole range of contaminants tested. Therefore some critical points have to be taken into account like the oxidising character of NaOCl and the risk of depositing organic residue at the end of the decontamination process. All these points have previously been discussed [39]. The last important point with regard to our results is the human factor implicated in the whole decontamination process. Indeed, this study has highlighted that decontamination modalities can condition decontamination efficacy just as in other studies that have already explored factors liable to modify it [26]. This study also underlines the importance of operator training so as to standardise decontamination and establish the most efficient procedure which takes into account the duality of handling hazardous and sterile drugs. A new pedagogical tool could be formulated to this end [42].

Conclusion

This study, assessing four solutions, concludes that sodium hypochlorite is the best decontaminant after standard application although an admixture of surfactant agent and isopropanol gave the same results after vigorous application on 23 contaminants. Decontamination efficiency depends on the solution used but also on application modality. An SDS admixture seems to be a good option, notably for vigorous chemical decontamination without any hazard to either materials or workers.

Acknowledgments

The authors thank Alexandra Tavernier, M.A. (University of Glasgow U.K.), Professeur Agrégée (France) for her assistance in correcting and editing the text.

Author Contributions

Conceptualization: Nicolas Simon, Sandrine Fleury-Souverain.

Formal analysis: Nicolas Simon, Sandrine Fleury-Souverain.

Investigation: Nicolas Simon.

Methodology: Nicolas Guichard, Pascal Bonnabry, Sandrine Fleury-Souverain.

Project administration: Pascal Bonnabry.

Resources: Sandrine Fleury-Souverain.

Supervision: Sandrine Fleury-Souverain.

Writing – original draft: Nicolas Simon, Nicolas Guichard, Sandrine Fleury-Souverain.

Writing – review & editing: Pascal Odou, Bertrand Decaudin, Pascal Bonnabry, Sandrine Fleury-Souverain.

References

1. Sessink PJ, Friemel NS, Anzion RB, Bos RP. Biological and environmental monitoring of occupational exposure of pharmaceutical plant workers to methotrexate. *Int Arch Occup Environ Health* 1994; 65: 401–403. <https://doi.org/10.1007/BF00383251> PMID: 8034364
2. Sessink PJM, Trahan J, Coyne JW. Reduction insurface contamination with cyclophosphamide in 30 US Hospital pharmacies following implementation of Closed-System Drug Transfer Device. *Hosp Pharm* 2013; 48:204–212. <https://doi.org/10.1310/hpj4803-204> PMID: 24421463
3. Hedmer M, Tinnerberg H, Axmon A, Jönsson BAG. Environmental and biological monitoring of antineoplastic drugs in four workplaces in a Swedish hospital. *Int Arch Occup Environ Health* 2008; 81:899–911 <https://doi.org/10.1007/s00420-007-0284-y> PMID: 18066576
4. Böhlandt A, Sverdel Y, Schierl R. Antineoplastic drug residues inside homes of chemotherapy patients. *Int J Hyg Environ Health*. 2017; 220: 757–765. <https://doi.org/10.1016/j.ijheh.2017.03.005> PMID: 28372941
5. Alexander K, Northrup N, Clarke D, Lindell H, Laver T. Engineering controls in veterinary oncology: A survey of 148 ACVIM board-certified oncologists and environmental surveillance in 20 specialty hospitals. *Vet Comp Oncol*. 2018; 16: 385–391. <https://doi.org/10.1111/vco.12390> PMID: 29446222
6. Falck K, Gröhn P, Sorsa M, Vainio H, Heinonen E, Holsti LR. Mutagenicity in urine of nurses handling cytostatic drugs. *Lancet* 1979; 1:1250–1. [https://doi.org/10.1016/s0140-6736\(79\)91939-1](https://doi.org/10.1016/s0140-6736(79)91939-1) PMID: 87722
7. Valanis BG, Vollmer WM, Labuhn KT, Glass AG. Association of antineoplastic drug handling with acute adverse effects in pharmacy personnel. *Am J Hosp Pharm* 1993; 50:455–62. PMID: 8442461
8. Valanis BG, Vollmer WM, Labuhn KT, Glass AG. Acute symptoms associated with antineoplastic drug handling among nurses. *Canc Nurs* 1993; 16:288–95.
9. McDiarmid M, Egan T. Acute occupational exposure to antineoplastic agents. *J Occup Med* 1998; 30:984–7.
10. Connor TH, Lawson CC, Polovich M, McDiarmid M. Reproductive health risks associated with the occupational exposures to antineoplastic drugs in health care settings: a review of the evidence. *J Occup Environ Med* 2014; 56:901–10. <https://doi.org/10.1097/JOM.0000000000000249> PMID: 25153300
11. Villarini M, Gianfredi V, Levorato S, Vannini S, Salvatori T, Moretti M. Occupational exposure to cytostatic/antineoplastic drugs and cytogenetic damage measured using the lymphocyte cytokinesis-block micronucleus assay: A systematic review of the literature and meta-analysis. *Mutat Res* 2016; 770:35–45. <https://doi.org/10.1016/j.mrrev.2016.05.001> PMID: 27894689
12. Harrison BR. Developing guidelines for working with antineoplastic drugs. *Am J Hosp Pharm* 1981; 38:1686–1693. PMID: 7304620
13. Zellmer WA. Reducing occupational exposure to potential carcinogens in hospitals. *Am J Hosp Pharm* 1981; 38(11):1679. PMID: 7304618
14. Zimmerman PF, Larsen RK, Barkley EW, Gallelli JF. Recommendations for the safe handling of injectable antineoplastic drug products. *Am J Hosp Pharm* 1981; 38(11): 1693–1695. PMID: 7304621
15. Yodaiken RE, Bennett D. OSHA work-practice guidelines for personnel dealing with cytotoxic (antineoplastic) drugs. Occupational Safety and Health Administration. *Am J Hosp Pharm* 1986; 43(5):1193–1204. PMID: 3717176
16. Occupational Safety and Health Administration. Controlling occupational exposure to hazardous drugs. *Am J Health Syst Pharm* 1996; 53(14):1669–1685. <https://doi.org/10.1093/ajhp/53.14.1669> PMID: 8827233
17. Martin S, Larson E. Chemotherapy-handling practices of outpatient and office-based oncology nurses. *Oncol Nurs Forum* 2003; 30(4):575–581. <https://doi.org/10.1188/03.ONF.575-581> PMID: 12861318

18. Centers for Disease Control and Prevention. The National Institute for Occupational Safety and Health. Hazardous exposure in healthcare—Antineoplastic drugs. <https://www.cdc.gov/niosh/topics/hazdrug/antineoplastic.html>
19. American Society of Health-System Pharmacists. ASHP guidelines on handling hazardous drugs. *Am J Health-Syst Pharm* 2006; 63:1172–1193.
20. GERPAC-Europharmat workgroup. Preparation and administration of drugs at risk for both workers and environment. 2007 http://www.euro-pharmat.com/documents/dm_de_preparation/guiderecommandationdm.pdf
21. International Society of Oncology Pharmacy Practitioners Standards Committee. ISOPP standards of practice. Safe handling of cytotoxics. *J Oncol Pharm Pract* 2007; 13:1–81. <https://doi.org/10.1177/1078155207082350> PMID: 17933809
22. European parliament. Preventing occupational exposure to cytotoxic and other hazardous drugs. Available on www.europeanbiosafetynetwork.eu
23. Power LA, Coyne JW. ASHP guidelines on hazardous drugs. *Am J Health-syst Pharm* 2018; 75: 1996–2031. <https://doi.org/10.2146/ajhp180564> PMID: 30327293
24. Boiano JM, Steege AL, Sweeney MH. Adherence to Precautionary Guidelines for Compounding Antineoplastic Drugs: A Survey of Nurses and Pharmacy Practitioners. *J Occup Environ Hyg*. 2015; 12: 588–602. <https://doi.org/10.1080/15459624.2015.1029610> PMID: 25897702
25. US Pharmacopeia. USP General chapter <800> Hazardous drugs—handling in healthcare settings. Available on www.usp.org
26. Simon N, Odou P, Décaudin B, Bonnabry P, Fleury-Souverain S. Efficiency of degradation or desorption methods in antineoplastic drug decontamination: a critical review. *J Oncol Pharm Pract*. 2019; 25: 929–946. <https://doi.org/10.1177/1078155219831427> PMID: 30786823
27. Adé A, Chauchat L, Ouellete Frève JF, Gagné S, Caron N, Bussi eres JF. Comparison of d econtamination efficacy of cleaning solutions on a biological safety cabinet workbench contaminated by cyclophosphamide. 2017 *Can J Hosp Pharm*; 70: 407–414. <https://doi.org/10.4212/cjhp.v70i6.1708> PMID: 29298999
28. Bo hlandt A, Groeneveld S, Fischer E, Schierl R. Cleaning Efficiencies of Three Cleaning Agents on Four Different Surfaces after Contamination by Gemcitabine and 5-fluorouracil. *J Occup Environ Hyg* 2015; 12:384–392. <https://doi.org/10.1080/15459624.2015.1009985> PMID: 25751496
29. Hon CY, Chua PP, Danyluk Q, Astrakianakis G. Examining factors that influence the effectiveness of cleaning antineoplastic drugs from drug preparation surfaces: a pilot study. *J Oncol Pharm Pract* 2014; 20:210–16. <https://doi.org/10.1177/1078155213497070> PMID: 23929731
30. L e LMM, Jolivot PA, Sadou Yaye H, Rieutord A, Bellanger A, Pradeau D et al. Effectiveness of cleaning of workplace cytotoxic surface. *Int Arch Occup Environ Health* 2013; 86:333–41. <https://doi.org/10.1007/s00420-012-0769-1> PMID: 22526087
31. Lee SG, Ambados F, Tkaczuk M, Jankewicz G. Paclitaxel exposure and its effective decontamination. *J Pharm Pract Res* 2009; 39:181–5.
32. Touzin K, Bussi eres JF, Langlois E, Lefebvre M, M etra A. Pilot Study Comparing the efficacy of two cleaning techniques in reducing environmental contamination with cyclophosphamide. *Ann Occup Hyg* 2010; 54: 351–9. <https://doi.org/10.1093/annhyg/meq004> PMID: 20118195
33. Queruau-Lamerie T, Nussbaumer S, D ecaudin B, Fleury-Souverain S, Goossens JF, Bonnabry P. Evaluation of decontamination efficacy of cleaning solutions on stainless steel and glass surfaces contaminated by 10 antineoplastic agents. *Ann Occup Hyg* 2013; 57:456–69. <https://doi.org/10.1093/annhyg/mes087> PMID: 23223271
34. Guichard N., Fekete S, Guillaume D, Bonnabry P, Fleury-Souverain S. Computer-assisted UHPLC-MS method development and optimization for the determination of 24 antineoplastic drugs used in hospital pharmacy. *J Pharm Biomed Anal*. 2019; 164: 395–401. <https://doi.org/10.1016/j.jpba.2018.11.014> PMID: 30439666
35. Guichard N, Rudaz S, Bonnabry P, Fleury-Souverain S. Validation and uncertainty estimation for trace amounts determination of 25 drugs used in hospital chemotherapy compounding units. *J Pharm Biomed Anal*. 2019; 172: 139–148. <https://doi.org/10.1016/j.jpba.2019.04.042> PMID: 31035095
36. Guichard N, Boccard J, Rudaz S, Bonnabry P, Fleury-Souverain S. Wipe sampling procedure optimization for the determination of 23 antineoplastic drugs used in hospital pharmacy. *Eur J Hosp Pharm*. 2019; accepted for publication
37. Anastasi M, Rudaz S, Queruau-Lamerie T, Odou P, Bonnabry P, Fleury-Souverain S. Efficacy of Two Cleaning Solutions for the Decontamination of 10 Antineoplastic Agents in the Biosafety Cabinets of a Hospital Pharmacy *Ann Occup Hyg* 2015; 59(7):895–898. <https://doi.org/10.1093/annhyg/mev031> PMID: 25979920

38. Simon N, Vasseur M, Pinturaud M, Soichot M, Richeval C, Humbert L et al. Effectiveness of a Closed-System Transfer Device in Reducing Surface Contamination in a New Antineoplastic Drug-Compounding Unit: A Prospective, Controlled, Parallel Study. *PLoS One* 2016; 11:e0159052 <https://doi.org/10.1371/journal.pone.0159052> PMID: 27391697
39. Simon N, Odou P, Décaudin B, Bonnabry P, Fleury-Souverain S. Chemical decontamination of hazardous drugs: a comparison of solution performances. *Annals of Work Exposure Health* 2020; 64: 114–124.
40. Vasseur M, Simon N, Picher C, Richeval C, Soichot M, Humbert L, et al. A decontamination process adding a tensioactive agent and isopropanol to a closed-system drug transfer device for better control of isolator contamination. A prospective, parallel study. *PLoS One*. 2018; 13(8):e0201335. <https://doi.org/10.1371/journal.pone.0201335> PMID: 30089139
41. European Pharmacopeia. Edition 9.8. Monograph « Parenteralia ». Strasbourg: 2019.
42. Berthod F, Bouchoud L, Grossrieder F, Falaschi L, Senhaji S, Bonnabry P. Learning good manufacturing practices in an escape room: Validation of a new pedagogical tool. *J Oncol Pharm Pract* 2019; <https://doi.org/10.1177/1078155219875504> PMID: 31566110