



HAL
open science

Outcome of large B-cell lymphoma patients treated with tafasitamab plus lenalidomide either before or after CAR-T-cells.

V. Camus, R. Houot, G. Brisou, B. Tessoulin, S. Bailly, P. Sesques, J. Decroocq, D. Krzisch, L. Obéric, F. Lemonnier, et al.

► To cite this version:

V. Camus, R. Houot, G. Brisou, B. Tessoulin, S. Bailly, et al.. Outcome of large B-cell lymphoma patients treated with tafasitamab plus lenalidomide either before or after CAR-T-cells.. Blood Advances, 2024, 8 (20), pp.5371-5381. 10.1182/bloodadvances.2024013726 . hal-04708799

HAL Id: hal-04708799

<https://hal.univ-lille.fr/hal-04708799v1>

Submitted on 13 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Outcome of patients with large B-cell lymphoma treated with tafasitamab plus lenalidomide either before or after CAR T-cell therapy

Vincent Camus,¹ Roch Houot,² Gabriel Brisou,³ Benoit Tessoulin,⁴ Sébastien Bailly,⁵ Pierre Sesques,⁶ Justine Decroocq,⁷ Daphné Krzisch,⁸ Lucie Oberic,⁹ François Lemonnier,¹⁰ Krimo Bouabdallah,¹¹ Arnaud Campidelli,¹² Ledraa Tounes,¹³ Julie Abraham,¹⁴ Charles Herbaut,¹⁵ Franck Morschhauser,¹⁶ Gandhi Laurent Damaj,¹⁷ Stéphanie Guidez,¹⁸ Sylvain Carras,¹⁹ Luc-Matthieu Fornecker,²⁰ Sylvain Choquet,²¹ Olivier Hermine,²² Jérôme Paillassa,²³ Adrien Chauchet,²⁴ Olivier Casasnovas,²⁵ Laurianne Drieu La Rochelle,²⁶ Cristina Castilla-Llorente,²⁷ Magalie Joris,²⁸ Vivien Dupont,²⁹ Alexandra Marquet,²⁹ Steven Le Gouill,³⁰ and Fabrice Jardin¹

¹Department of Hematology, Centre Henri Becquerel, Rouen, France; ²Department of Hematology, Centre Hospitalier Universitaire Rennes, University of Rennes, INSERM U1236, Etablissement Français du Sang, Rennes, France; ³Department of Hematology, Institut Paoli-Calmettes, Marseille, France; ⁴Department of Hematology, Nantes University Hospital, Nantes, France; ⁵Department of Hematology, Centre Hospitalier Universitaire Estaing, Clermont-Ferrand, France; ⁶Department of Hematology, Lyon-Sud Hospital, Hospices Civils de Lyon, Claude Bernard Lyon 1 University, Pierre-Benite, France; ⁷Department of Hematology, Cochin University Hospital, Paris, France; ⁸Department of Hemato-oncology, Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Université de Paris, Paris, France; ⁹Department of Hematology, Institut Universitaire du Cancer, Toulouse-Oncopole, Toulouse, France; ¹⁰Department of Hematology, Henri Mondor University Hospital, Créteil, France; ¹¹Department of Hematology and Cellular Therapy, Bordeaux University Hospital, Bordeaux, France; ¹²Department of Hematology, Hôpital de Brabois, Nancy University Hospital, Nancy, France; ¹³Department of Hematology, Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Antoine, Université de Paris, Paris, France; ¹⁴Department of Hematology, Dupuytren Hospital, Limoges University Hospital, Limoges, France; ¹⁵Department of Hematology, Montpellier University Hospital, Montpellier, France; ¹⁶Department of Hematology, Claude Huriez Hospital, Lille University Hospital, Lille, France; ¹⁷Department of Hematology, Caen University Hospital, Caen, France; ¹⁸Department of Hematology, Poitiers University Hospital, Poitiers, France; ¹⁹Department of Hematology, Albert Michallon University Hospital, Grenoble, France; ²⁰Department of Hematology, Strasbourg University Hospital, Strasbourg, France; ²¹Department of Hematology, Assistance Publique-Hôpitaux de Paris, Hôpital Pitié-Salpêtrière, Université de Paris, Paris, France; ²²Department of Hematology, Assistance Publique-Hôpitaux de Paris, Hôpital Necker, Université de Paris, Paris, France; ²³Department of Hematology, Angers University Hospital, Angers, France; ²⁴Department of Hematology, Besançon University Hospital, Besançon, France; ²⁵Department of Hematology, Dijon University Hospital, Dijon, France; ²⁶Department of Hematology, Tours University Hospital, Tours, France; ²⁷Department of Hematology, Gustave Roussy, Villejuif, France; ²⁸Department of Hematology, Amiens University Hospital, Amiens, France; ²⁹Lymphoma Academic Research Organisation, Lyon-Sud Hospital, Pierre-Bénite, France; and ³⁰Department of Hematology, Institut Curie, Paris, France

Key Points

- TAFAs combined with LEN showed limited efficacy after CAR T-cell failure in a high-risk population.
- Outcomes were comparable between the TAFAs-LEN combination and other treatments for the first progression after CAR T-cell therapy.

Tafasitamab plus lenalidomide (TAFAs-LEN) treatment relevance pre- or post-anti-CD19 chimeric antigen receptor (CAR) T-cell therapy is debated. We analyzed patients with large B-cell lymphoma in the DESCAR-T registry treated with axi[1]cel or tisa-cel in ≥ 3 rd line and TAFAs-LEN before ($n = 15$, “TL-pre-CAR-T” set) or directly after ($n = 52$, “TL-post-CAR-T” set) CAR T-cell therapy. We compared TAFAs-LEN v. other treatments using inverse probability weighting in the TL-post-CAR[1]T set. In the TL-post-CAR-T set, the median progression-free survival (mPFS), overall survival (mOS), and duration of response (mDOR) since the first treatment for progression (mPFS2/mOS2/mDOR2) were 3, 4.7, and 8.1 months, respectively. The best overall response rate (bORR) and best complete response rate (bCRR) after TAFAs-LEN were 13.5% and 7.7%, respectively. Outcomes were better for patients who relapsed > 6 months after CAR T-cell therapy (mPFS2: 5.6 vs 2 months, $P = .0138$; mOS2: not reached vs 3.8 months, $P = .0034$). The bORR and bCRR between TAFAs-LEN and other treatments were 20.6% vs 24.9% and 11.6% vs 15.6%, respectively. Outcomes were similar between TAFAs-LEN and other treatments (mPFS2: 2.9 vs 2.4 months, $P = .91$; mOS2: 3.3 vs 5.5 months, $P = .06$). In an exploratory analysis of the TL-pre-CAR-T set, the median TAFAs-LEN treatment

Submitted 28 May 2024; accepted 31 July 2024; prepublished online on *Blood Advances* First Edition 20 August 2024; final version published online 11 October 2024. <https://doi.org/10.1182/bloodadvances.2024013726>.

Data from the DESCAR-T registry are subject to controlled access by the Lymphoma Academic Research Organization owing to privacy and legal requirements and proprietary reasons. Anonymized individual patient (or participant) data (IPD) requests will be promptly reviewed by the corresponding author, Vincent Camus

(vincent.camus@chb.unicancer.fr) and the scientific committee of the DESCAR-T registry.

The full-text version of this article contains a data supplement.

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

duration before CAR-T was 3.7 months with no patient becoming CD19 negative. The bORR, bCRR, 6-month PFS, and OS rates after CAR T-cell infusion were 45.5%, 36.4%, 20.1%, and 58.2%, respectively. Neither TAFa-LEN nor comparative salvage treatment improved outcomes for patients relapsing after CAR T-cell therapy.

Introduction

Chimeric antigen receptor (CAR) T-cell therapy has been the standard of care for relapsed/refractory large B-cell lymphomas (R/R LBCLs) since 2018.¹⁻³ Despite the efficacy of these treatments, durable responses and curative effects have been reported in <40% of patients.⁴⁻⁷ The most common mechanism of resistance to CAR T cells is the loss of the target antigen, which is observed in ~30% of patients with LBCL due to negative selection pressure.^{1,4,8-10} Commercial CAR T-cell therapies all target CD19, a 95-kDa transmembrane glycoprotein belonging to the immunoglobulin superfamily, which is found on the surface of both normal and tumor B lymphocytes.^{11,12} CD19 expression is commonly measured via immunohistochemistry (IHC) of formalin-fixed paraffin-embedded tumor biopsies or flow cytometry of live suspended cells. The most widely used clone is LE-CD19, which contains a cytosolic epitope (the intracellular domain of the protein). However, this technique does not determine whether CD19 integrity is maintained after anti-CD19 treatment.^{9,13} Commercially available CAR T-cell therapies currently used for the treatment of R/R LBCL include a single-chain variable fragment derived from a murine monoclonal antibody (mAb) against CD19 known as the FMC63 clone.^{14,15} However, despite the high cost of CAR T-cell therapy and the existence of simple techniques to detect the expression of CD19, health authorities and official labels do not require evidence of CD19 positivity before the administration of anti-CD19 CAR T-cell therapies. Responses were observed in CD19-negative patients after anti-CD19 CAR T-cell therapy in the ZUMA-1 clinical trial¹ but the overall efficacy of these therapies in CD19-negative patients remains largely uncertain.

Additionally, tafasitamab (TAFa) is a cytolytic anti-CD19 mAb approved by the US Food and Drug Administration in 2020 and by the European Medical Agency in 2021 for the treatment of adult patients with R/R LBCL in combination with lenalidomide (LEN). TAFa is produced from a humanized murine anti-CD19 antibody, specifically the 4G7 clone,¹⁶ and has a terminal elimination half-life of 17 days.¹⁷ Emerging data have suggested that administering TAFa before anti-CD19 CAR T-cell therapy could lead to a delay in CD19 re-expression or demasking in pre-CAR T-cell biopsy samples, potentially interfering with patient treatment plans in centers that restrict CD19-targeted CAR T-cell therapy to patients with preserved CD19 expression.¹⁸ Furthermore, using a TAFa-competing antibody to detect CD19 expression in samples from patients exposed to TAFa may result in a reduction in signal intensity and confusion between CD19 epitope masking and antigen loss.¹⁹ In vitro data have revealed direct competition for CD19 binding between TAFas and CD19-targeted CAR T-cell therapy in mantle cell lymphoma, LBCL, and B-cell acute lymphoblastic leukemia cell lines. There is a partial overlap between the 4G7 and FMC63 clones at an epitope centered on residue R144, without

compromising CAR T-cell activity in terms of antigen-specific killing, degranulation, cytokine production, or proliferation.²⁰ However, there is a scarcity of clinical data regarding the sequence of the 2 anti-CD19 therapies in patients with R/R LBCL. Initial data on a limited number of patients or xenograft models seem to indicate that the use of an anti-CD19 mAb followed by an anti-CD19 CAR T-cell therapy does not appear to be detrimental and does not seem to cause the CD19 target antigen to disappear.²¹⁻²³ Consequently, it remains uncertain whether exposure to one form of anti-CD19 therapy might compromise the effectiveness or safety of the other. Therefore, to better explore this topic, we aimed to assess outcomes in patients with R/R LBCL treated with TAFa-LEN as a pre- or post-anti-CD19 CAR T-cell therapy in real-life settings in France.

Patients and methods

Patients and data collection

This study is a retrospective analysis of data from adult patients with R/R LBCL collected before 1 September 2023 in the DESCAR-T registry who received TAFa-LEN as a pre- (previous treatment line or bridge: “TL-pre-CAR-T set”) or post-anti-CD19 CAR T-cell therapy (to treat the first progression after CAR T cell: “TL-post-CAR-T set”) between 2018 and 2023 (supplemental Methods). The analysis was restricted to patients who received CAR T-cell therapy with marketing authorization in France for treating LBCL in the third line or later (L3+). To contextualize the results observed in patients who received TAFa-LEN directly after CAR T-cell therapy, we performed an indirect comparison with data from other patients with LBCL from the DESCAR-T registry who were refractory to treatment or relapsed after CAR T-cell therapy was administered in L3+ and who did not receive TAFa-LEN for their first TL-post-CAR T-cell therapy (supplemental Methods). This study was approved by the Lymphoma Study Association (LYSA) (10 January 2023) and DESCAR-T (17 January 2023) Scientific Committees. Patients’ nonopposition to participation in this study was obtained through a written information note provided before inclusion in the registry. DESCAR-T is registered under the [ClinicalTrials.gov](https://clinicaltrials.gov) identifier NCT04328298, and the study was conducted in accordance with the Declaration of Helsinki.

Study end points

The coprimary end points were progression-free survival (PFS), overall survival (OS), and duration of response (DOR) after treatment for the first progression post-CAR T-cell therapy, defined as PFS2, OS2, and DOR2, respectively, in the TL-post-CAR-T set. Secondary end points in the TL-post-CAR-T set were CAR T-cell-related safety events, response rates, PFS, and OS after CAR T-cell infusion, outcome according to CD19 status on biopsy if available, and indirect comparisons of outcomes between TAFa-LEN and other treatments as the first treatment for progression

after CAR T-cell therapy. PFS2 and OS2 were compared between patients treated with TAFa-LEN and LEN (single agent) post hoc.

In an exploratory analysis, we evaluated the duration of the response to TAFa-LEN, washout time between the last TAFa-LEN administration and CAR T-cell infusion, CD19 status before TAFa-LEN, response rate, CAR T-cell-related safety events, and PFS/OS after CAR T-cell infusion in the TL-pre-CAR-T set.

Indirect comparison procedures

We conducted an inverse probability weighting (IPW) comparison of TAFa-LEN v. other treatments (chemotherapy, bispecific anti-CD3/anti-CD20 antibody, transplant, radiotherapy, kinase inhibitors, and other anticancer therapies, excluding anti-CD19 antibodies) and v. LEN (single agent) in the TL-post-CAR-T set. The patients included in the comparison were identified in the DESCAR-T registry, which included ~1500 patients treated with CAR T-cell therapy for LBCL in L3+ (details provided in supplemental Methods).

Statistical considerations

Quantitative variables are displayed as means, standard deviations, medians, ranges, and quartiles. Qualitative variables are expressed as frequencies and percentages. Response rates, including complete response (CR) and partial response rates, are expressed as percentages with 95% exact Clopper-Pearson confidence intervals (supplemental Methods). Survival curves were obtained using the Kaplan-Meier method. P values for survival analysis were calculated using the log-rank test unless otherwise specified (supplemental

Methods). The level of significance for each test was 5%. Statistical analyses were performed using SAS software version 9.4 and AdClin version 3.4.0.

Results

TL-post-CAR-T set

Patient characteristics and outcomes after CAR T-cell infusion and before TAFa-LEN initiation. Within the DESCAR-T registry, we identified 52 patients treated with TAFa-LEN for first progression after CAR T-cell infusion in 18 different French centers (Figure 1).

The patient characteristics are summarized in Table 1. Briefly, these characteristics were as follows: median age, 65 years; Eastern Cooperative Oncology Group (ECOG) score 0 to 1, 88%; age-adjusted international prognostic index (aaiPI) 2 to 3, 61.7%; median prior lines of therapy (PLT) at the time of CAR T-cell treatment, 2; and refractory disease (defined as no response to the last line of therapy before enrollment), 85.4%. For some patients, only 1 PLT was recorded within the DESCAR-T registry by investigators, but 31 (60%) and 21 (40%) patients received CAR T-cell therapy in the third line and fourth line, respectively.

The best overall response (bORR) rate, best complete response rate (bCRR), 6-month PFS, and OS after the first CAR T-cell infusion (Tisagenlecleucel [tisa-cel]: n = 21; Axicabtagene ciloleucel [axi-cel]: n = 31) were 63.5%, 34.6%, 21.2%, and 62.3%, respectively (supplemental Table 1; supplemental Figure 1). Failure




Figure 1. Study flowchart. DESCAR-T, Dispositif d'Enregistrement et Suivi des patients traités par CAR T cells.

Table 1. Patient characteristics in the TL-post-CAR-T set at the time of enrollment in the DESCAR-T registry

	First progression after CAR T-cell therapy	
	N = 52	
Age (y)		
Median (min; max)	65.5 (20; 81)	
Age ≥65 y	29	(55.8%)
Sex, male	30	(57.7%)
ECOG		
0-1	44	(88.0%)
>2	6	(12.0%)
Missing	2	
LDH > normal		
No	21	(41.2%)
Yes	30	(58.8%)
Ann Arbor stage		
I-II	6	(12.0%)
III-IV	44	(88.0%)
Missing	2	
aaIPI		
0	1	(2.1%)
1	17	(36.2%)
2	28	(59.6%)
3	1	(2.1%)
Missing	5	
Histology		
LBCL	39	(75.0%)
Transformed FL or other indolent	12	(23.1%)
PMBL	1	(1.9%)
HCT-CI-comorbidity evaluated	52	(100.0%)
At least 1 HCT-CI-comorbidity	25	(48.1%)
Nb of PLT before CAR T-cell therapy*		
Median (min; max)	2.0 (1; 6)†	
Prior autologous transplant	5	(9.6%)
Time from initial diagnosis to CAR T-cell therapy eligibility (y)		
Median (min; max)	1.46 (0.2; 21.0)	
Disease status before leukapheresis		
Refractory	41	(85.4%)
Relapse	7	(14.6%)
Missing	4	
BT	44	(84.6%)
Disease status before CAR T-cell infusion		
CR	3	(6.8%)
Partial response (PR)	11	(25.0%)
Stable disease	4	(9.1%)
Progressive disease	25	(56.8%)
Not evaluated	1	(2.3%)

Table 1 (continued)

	First progression after CAR T-cell therapy	
	N = 52	
CAR T product administered		
Tisa-cel	21	(40.4%)
Axi-cel	31	(59.6%)

ECOG, Eastern Cooperative Oncology Group; FL, follicular lymphoma; HCT-CI, hematopoietic stem cell transplant comorbidity index; LDH, lactate dehydrogenase; max, maximum; min, Minimum; PMBL, primary mediastinal B-cell lymphoma.

*Up to 10 treatment lines can be collected from the register.

†Ongoing query within the DESCAR-T registry for patients with only 1 line of prior treatment.

after CAR T-cell therapy occurred after a median of 2.8 months (range, 0.4-34.7). Overall, 42 patients experienced early (ie, ≤6 months) failure after CAR T-cell therapy, and 10 experienced late (>6 months) failure.

Safety was evaluated in all patients. The all-grade cytokine release syndrome (CRS) and neurotoxicity rates were 82.7% (grade 3-4: n = 2, 3.8%) and 40.4% (grade 3-4: n = 7, 13.5%), respectively. Prolonged cytopenia and grade 3 to 5 infections occurred in 38 (73.1%) and 24 (46.2%) patients, respectively. In patients with CAR T-cell-specific toxicity (n = 45/52), the rates of tocilizumab use and intensive care unit admission were 71.1% and 17.8%, respectively.

Response rates and outcomes after TAFALLEN initiation for first progression after CAR T-cell therapy.

The median (Q1; Q3) time between CAR T-cell infusion and TAFALLEN initiation was 2.84 (1.1; 5.7) months. The outcomes of 2 patients who received TAFALLEN without LEN according to the DESCAR-T registry were reported by the investigators but were not excluded from the present intention-to-treat analysis.

After a median follow-up duration of 7 (5.9-13) months, the median PFS and OS since the first treatment for progression (mPFS2/mOS2) were 3 (95% confidence interval, 1.9-4.2) and 4.7 (3-5.6) months, respectively; the median (Q1; Q3) DOR2 was 8.1 (1; NA) months (supplemental Figure 2). The mPFS2 increased when TAFALLEN was initiated >6 months after CAR T-cell infusion (late CAR T-cell failure subgroup): 5.6 (3.2-NA) v. 2 (1.6-3.1), P = .0138. In addition, mOS2 was not reached (5.6-NA) when TAFALLEN was introduced later than the 6-month cutoff (v. 3.8 [2.2-5] months) for patients in the ≤6-month subset (P = .0034) (Figure 2). The causes of death after the first treatment for progression were lymphoma (91.4%), concurrent illness (5.7%), and late toxicity (2.9%) (Table 2). Failure after TAFALLEN occurred after a median of 3.35 months (0.5-9.9). Three late-relapsing patients remained in CR at the last contact.

Outcomes according to CD19 status. CD19 expression before CAR T-cell infusion, after CAR T-cell infusion, and after TAFALLEN treatment is summarized in supplemental Table 3. In most patients, CD19 expression was not routinely assessed. The bCRR after

Downloaded from http://ashpublications.org/bloodadvances/article-pdf/8/20/5371/22247370/bloodadv_2024-013726-main.pdf by guest on 13 November 2024




Figure 2. Outcomes Based on Timing of Progression After CAR T-Cell Therapy. PFS (A) and OS (B) since treatment for the first progression (PFS2 and OS2) in the TL-post-CAR-T set according to early (≤ 6 months) vs late (> 6 months) TFA-LEN introduction. CI, confidence interval; NA, not available.

CAR T-cell therapy was numerically lower in patients who were negative for CD19 before CAR T-cell therapy than in patients who were positive for CD19 ($n = 2/7$, 28.6% v. $n = 5/12$, 41.7%; supplemental Table 4). The mOS after CAR T-cell infusion was lower in patients with negative vs positive CD19 expression before CAR T-cell infusion (6 [1.5; NA] vs 8 [2.9; NA] months), but the median PFS was similar (2.8 [0.9; 3.2] vs 2.7 [1; 7.8] months; supplemental Figure 3). mPFS2 and OS2 did not differ according to the CD19 status before CAR T-cell therapy (supplemental Figure 4). mPFS2 and OS2, according to the CD19 status after CAR T-cell therapy were also similar (supplemental Figure 5).

Comparison between TFA-LEN and other treatments as initial therapies for progression after CAR T-cell therapy.

After IPW, outcomes were compared between 43 patients treated with TFA-LEN (including 2 patients with TFA only) and 354 control patients treated with other agents (immunomodulatory drug, 47.2%; chemotherapy, 25.7%; CD3/CD20 bispecific antibodies, 9.9%; and others, 17.2%; supplemental Table 5). A diagram illustrating the constitution of the efficacy comparison cohort is provided

in supplemental Figure 6, including justification for the exclusion of patients due to missing values for weighting variables or other reasons. Patient characteristics were well balanced between the groups, with standardized mean differences < 0.15 for all weighting variables (data not shown), but the median (Q1; Q3) follow-up was longer in the control group (19.1 [10.9; 30.1] vs 13 [4.6; 16.2] months). The bORR and bCRR between the TFA-LEN and control groups were 20.6% v. 22.1% and 11.6% v. 13.8%, respectively (supplemental Table 6). PFS and OS were not significantly different between the 2 groups (median PFS, 2.9 [2-2.6] vs 2.2 [2-2.5] months, $P = .77$; median OS, 3.3 [1.8-6.4] vs 5.1 [4.1-6] months, $P = .1$; Figure 3). All the sensitivity analyses led to the same conclusions (supplemental Figure 8).

Comparison between TFA-LEN and LEN as initial therapies for progression after CAR T-cell therapy.

After IPW, outcomes were compared between 43 patients treated with TFA-LEN (including 2 patients treated with TFA only) and 52 control patients treated with LEN (supplemental Figure 7). Patient characteristics were well balanced between groups with standardized

Table 2. Causes of death since the first treatment for progression in the TL-post-CAR-T set

	TL-post-CAR-T set		Period of first progression ≤ 6 mo		Period of first progression > 6 mo	
	N = 52		n = 42		n = 10	
Event according to OS definition since first treatment for progression						
No	17	(32.7%)	8	(19.0%)	9	(90.0%)
Yes	35	(67.3%)	34	(81.0%)	1	(10.0%)
If yes, cause of death						
Progression	32	(91.4%)	31	(91.2%)	1	(100.0%)
Concurrent illness	2	(5.7%)	2	(5.9%)	0	(0.0%)
Late toxicity	1	(2.9%)	1	(2.9%)	0	(0.0%)




Figure 3. Comparison between TAFa-LEN and other treatments as initial therapies for progression after CAR T-cell therapy. PFS (A) and OS (B) since the first treatment for progression after CAR T-cell therapy according to IPW using stabilized weight (SW) method between TAFa-LEN and other treatments.

mean differences <0.15 for all weighting variables (data not shown), but the median follow-up was longer in the control group: 22.8 (8.5; 30.2) v. 6.2 (3.3; 13) months. The bORR and bCRR between the TAFa-LEN and LEN groups were 13.9% v. 16.4% ($P = .74$) and 6% v. 13.4% ($P = .2$), respectively (supplemental Table 7). PFS and OS were similar between the TAFa-LEN and LEN groups (median PFS: 3 [1.5-4.6] vs 1.8 [1-2.4] months, hazard ratio = 0.71 [0.44-1.13]; median OS 4.2 [2.4-6] vs 3.3 [2.3-3.9] months, hazard ratio = 0.88 [0.54-1.44]; Figure 4).

TL-pre-CAR-T set

Patient characteristics. Within the DESCART registry, we identified 15 patients who received TAFa-LEN at any time before CAR T-cell infusion, as a PLT or bridging therapy (BT), in 12

different French centers (Figure 1). The patients' characteristics were as follows: median (range) age, 71 (47-80) years; male sex, 66.7%; ECOG score 0 to 1, 86.7%; elevated lactate dehydrogenase, 50%; stage III to IV, 84.6%; aalPI 2 to 3, 41.7%; median PLT, 3 (2-7); refractory disease, 100%; and prior autologous transplant, 6.7% (Table 3). TAFa-LEN was given as a PLT before CAR T-cell therapy in 13 patients. Among them, only 2 patients received no treatment between TAFa-LEN and CAR T-cell infusion, and all the other patients received a different BT. In addition, TAFa-LEN was given as a bridge before CAR T-cell therapy in 3 patients (1 patient received TAFa-LEN for both PLT and BT) (supplemental Table 8).

For patients receiving TAFa-LEN before treatment ($n = 13$), the median (Q1; Q3) duration of treatment with TAFa-LEN was 3.7 (2; 7.4) months, and the median (Q1; Q3) washout time between the




Figure 4. Comparison between TAFa-LEN and LEN as initial therapies for progression after CAR T-cell therapy. PFS (A) and OS (B) since the first treatment for progression after CAR T-cell therapy according to IPW using SW between TAFa-LEN and LEN (single agent).

Table 3. Patient characteristics in the TL-pre-CAR-T set at the time of enrollment in the DESCAR-T registry

	TL-pre-CAR-T set	
	N = 15	
Age (y)		
Median (min; max)	71.0 (47; 80)	
Age ≥65 y	10	(66.7%)
Sex, male	10	(66.7%)
ECOG		
0-1	13	(86.7%)
>2	2	(13.3%)
LDH > normal		
No	7	(50.0%)
Yes	7	(50.0%)
Missing	1	
Ann Arbor stage		
I-II	2	(15.4%)
III-IV	11	(84.6%)
Missing	2	
aalPI		
1	7	(58.3%)
2	4	(33.3%)
3	1	(8.3%)
Missing	3	
Histology		
LBCL	11	(73.3%)
Transformed FL	3	(20.0%)
Transformed Hodgkin	1	(6.7%)
HCT-CI-comorbidity evaluated	15	(100.0%)
At least 1 HCT-CI-comorbidity	7	(46.7%)
Nb of PLT before CAR-T*		
Median (min; max)	3.0 (2; 7)	
Prior autologous transplant	1	(6.7%)
Time from the initial diagnosis to CAR T-cell eligibility (y)		
Median (min; max)	1.62 (0.5; 18.2)	
Disease status before leukapheresis		
Refractory	14	(100.0%)
Missing	1	
Disease status before CAR T-cell infusion		
CR	2	(13.3%)
PR	3	(20.0%)
Progressive disease	10	(66.7%)
CAR T product administered		
Tisa-cel	3	(20.0%)
Axi-cel	12	(80.0%)

Abbreviations are explained in Table 1.

*Up to 10 treatment lines may be collected from the DESCAR-T registry.

Table 4. Best response after CAR T-cell infusion in the TL-pre-CAR-T set

	Evaluable patients in the TL-pre-CAR-T set	
	n = 11/15	
Best response		
CR	4	(36.4%)
PR	1	(9.1%)
Stable disease	1	(9.1%)
Progressive disease	4	(36.4%)
Not evaluated	1	(9.1%)
bORR*		
bORR	5	(45.5%)
95% confidence interval	(16.7%-76.6%)	
Time to best response (months)		
N	5	
Missing	0	
Mean (SD)	0.97 (0.055)	
Median	0.99	
Q1; Q3	1.0; 1.0	
P1; P99	0.9; 1.0	

SD, standard deviation.

*Defined as (CR + PR).

last TFAA-LEN administration and CAR T-cell infusion was 2.7 (2.1; 3.4) months (supplemental Table 9). The bORR after TFAA-LEN as a PLT was not collected in the registry.

For patients receiving TFAA-LEN as BT (n = 3), the median (Q1; Q3) duration of bridging with TFAA-LEN was 23 (14; 58) days, with a median washout time of 20 (3; 31) days. The responses after bridging with TFAA-LEN were partial response (n = 2) and progressive disease (n = 1) (supplemental Table 10).

Description of CD19 status. CD19 status (IHC) before TFAA-LEN was positive in 4 patients (26.7%), negative in 1 (6.7%), and not tested/not done in 10 (66.7%). On subsequent biopsy (if available) after TFAA-LEN and before CAR T-cell infusion, CD19 status was re-evaluated as follows: positive (n = 3, 20%), negative (n = 1, 6.7%), or not tested/not done (n = 11, 73.3%). No patients were reported to become CD19 negative after TFAA-LEN. After CAR T-cell infusion, 1 additional patient became negative for CD19 (supplemental Table 3).

Outcomes after CAR T-cell infusion. With a median (Q1; Q3) follow-up duration of 2.8 (1; 6.3) months, the bORR, bCRR, 6-month PFS rate, and OS rate after CAR T-cell infusion (tisa-cel: n = 3; axi-cel: n = 12) were 45.5%, 36.4%, 20.1%, and 58.2%, respectively (Table 4; Figure 5; supplemental Figure 9). Notably, we did not analyze treatments administered in the event of progression after CAR T-cell infusion in this patient set (2 patients




Figure 5. Outcomes after CAR T-cell infusion. PFS (A) and OS (B) after CAR T-cell infusion in the TL-pre-CAR-T set.

received and responded to subsequent treatment for progression after CAR T-cell infusion).

The safety was evaluated in 14 patients. The all-grade CRS and neurotoxicity rates were 64.3% and 14.3%, respectively (supplemental Figure 9). No grade 3 to 4 CRS was observed. In patients with CAR T-cell-specific toxicity ($n = 10/14$), the rates of tocilizumab use and intensive care unit admission were 30% and 10%, respectively. The rates of prolonged cytopenia and infection were 50% and 14.3%, respectively. Causes of death were lymphoma ($n = 4$, 66.7%), concurrent illness ($n = 1$, 16.7%), and acute toxicity ($n = 1$, 16.7%; data not shown).

Discussion

In this retrospective study of the French DESCAR-T registry, we reported the largest cohort to date of patients treated sequentially (either before or directly after CAR T-cell infusion) with 2 therapies targeting CD19: commercial anti-CD19 CAR T-cell therapy and the TAFALLEN combination.

In the TL-post-CAR T-cell cohort, our results confirmed the poor prognosis of patients who relapsed after CAR T-cell therapy regardless of the treatment received. The median PFS was 3 months and the median OS was 4.7 months. This population displayed high-risk features, with 57.6% having aIPI 2 to 3 disease, and we observed a slight overrepresentation of patients treated with tisa-cel (40.4%), compared with 27% in the study by Tomas et al and 36% in the study by Di Blasi et al.^{24,25} Our results are consistent with those for other cohorts in the literature regarding the outcomes of post-CAR T-cell-treated patients who subsequently benefited from subsequent lines of treatment. Indeed, the median PFS and OS after the first CAR T-cell treatment were 1.9 months and 8.5 months, respectively, in the study by Tomas A et al, and 2.8 months and 5.2 months, respectively, in the study by Di Blasi et al.^{24,25} In the cohort of the US lymphoma CAR-T consortium in L3+, the outcome was also poor, with a mPFS2 of just 55 days.²⁶ L2+ patients who received treatment for the first progression after CAR T in the experimental arm of ZUMA-7 displayed also a poor outcome, with a

median PFS2 and OS2 of 1.7 and 8.1 months, respectively, after salvage chemotherapy.²⁷ Checkpoint inhibitors were also not an effective salvage strategy for most patients after CAR T-cell failure, with median PFS2 and OS2 of 54 and 159 days, respectively, as reported by Major et al.²⁸ Notably, Sesques et al recently described a cohort of 67 patients treated with glofitamab, a CD20 \times CD3 bispecific antibody, for first progression after CAR T-cell therapy in a recent clinical trial (NCT04703686). Although the mPFS was short (~4.5 months), the bORR (65.9%) and bCRR (36.4%) indicated antitumor efficacy, and the mOS was not reached with a follow-up of ~10 months.²⁹ Epcoritamab, another CD20 \times CD3 bispecific antibody, also demonstrated antitumor efficacy (bORR, 54.1%; bCRR, 34.4%; median PFS, 4.4 months; and median OS, not reached) in patients who relapsed after CAR T-cell therapy ($n = 61$) in a recent clinical trial (NCT03625037).³⁰ Therefore, targeting an epitope other than CD19 after CAR T-cell failure using a CD3 \times CD20 bispecific antibody seems to be an interesting approach. However, more in-depth subgroup analyses are needed to identify the patients who could benefit the most from this strategy.

In a recent study reported in the form of an abstract at the American Society of Hematology (ASH) conference in 2023, the response and survival rates after TAFALLEN in a real-life setting appeared lower than those reported in the pivotal Tafasitamab plus lenalidomide in relapsed or refractory diffuse large B-cell lymphoma (L-MIND) study.³¹ In that study, the median PFS was 2.9 months (including all causes of death in progression events) and the median OS was 8.9 months, which is quite similar to the results observed in the TL-post-CAR T-cell cohort. In patients who received TAFALLEN after CAR T-cell therapy in the study by Ruckdeschel et al ($n = 24$), the median PFS was 7.9 months, and the median OS was 9.2 months. Post-TAFALLEN bCRR was lower in patients who were already exposed to CAR T-cell therapy (9.7% v. 21%), similar to our observation (bCRR TL-post-CAR-T set, 7.7%). However, outcomes after anti-CD19 mAb given post-CD19 CAR T-cell therapy were superior in another recently published real-life study involving a cohort of 53 patients treated with loncastuximab tesirin or TAFALLEN.³² In that study, the bORR and bCRR were 27% and 10%, respectively (compared with 13.5% and 7.7%, respectively, in our

TL-post-CAR T set), and the median PFS and OS were 8 and 20.9 months, respectively (compared with 3 and 4.7 months, respectively, in our study). Nevertheless, in that study, patients received a median of 1 line of treatment (range 0-5) between CAR T-cell and anti-CD19 mAb, with a median time of 7.3 (1.2-38.2) months between the 2 rounds of anti-CD19 immunotherapy, whereas in our cohort, patients received TFAFA-LEN immediately after the first post-CAR T-cell therapy (median time between CAR T-cell therapy and the start of TFAFA-LEN: 2.84 months). In addition, our exploratory analysis revealed that the outcome was better in the late CAR T-cell failure subgroup than that in the early failure subgroup (>6 months after CAR T-cell infusion). We speculate that a washout period of several months, probably >6 months, is associated with a better efficacy of subsequent anti-CD19 treatment. However, this interpretation could be biased because patients who relapse <3 months after CAR T-cell therapy have a well-known worse prognosis^{24,25} and because we were not able to make an indirect comparison between TFAFA-LEN and other treatments in the subset of patients with late CAR T-cell failure (for whom the sample size was too low). We acknowledge the survival analysis because CAR T-cell infusion requires careful analysis because patients administered TFAFA-LEN were alive before TFAFA-LEN administration, creating an immortality bias. However, our main analyses focused on efficacy since the first progression/TFAFA-LEN initiation, and not after CAR T-cell therapy.

Regarding the TL-pre-CAR-T set (n = 15), the limited sample size and short follow-up period prevented definitive conclusions from being drawn and hindered the creation of a relevant matched control cohort; therefore, we could not determine the specific impact of TFAFA-LEN on post-CAR T-cell outcomes. In our opinion, TFAFA-LEN is poorly represented in the DESCAR-T registry as a PLT because physicians fear CD19 expression loss by providing patients with anti-CD19 therapy when they can potentially benefit from anti-CD19 CAR T-cell therapy at subsequent lines. Patients in the TL-pre-CAR T-cell cohort presented high-risk characteristics, with a median age of 71 years, refractory disease in all patients, and a median of 3 previous lines of treatment, probably partly explaining the poor outcomes of these patients after CAR T-cell therapy (median OS, 8.1 months; median PFS, 3 months). The DESCAR-T registry did not include data on the best response to treatment with TFAFA-LEN as a PLT (n = 13) or other treatments given before treatment with CAR T-cell therapy. However, the median duration of treatment with TFAFA-LEN as a PLT was short (2.7 months), and all patients were reported to have refractory disease before leukapheresis; therefore, we speculate that TFAFA-LEN was not effective in this selected subgroup. Notably, by definition, the TFAFA-LEN efficacy assessment was biased in this subgroup because the patients were selected because they received CAR T-cell therapy after TFAFA-LEN failure. It was recently reported that CAR T-cell therapies remained effective in patients with R/R LBCL after prior exposure to bispecific antibodies targeting different antigens³³ but there is a scarcity of data in the literature regarding the outcomes of patients who are followed by CAR T-cell therapy. In a case series of 8 patients treated with TFAFA-LEN before CAR T-cell therapy, 4 patients achieved subsequent CR after CAR T-cell therapy.³⁴ Another study reported the use of anti-CD19 treatment, but unlike TFAFA, loncastuximab tesirine (antibody-drug conjugated) was administered to 14 patients who subsequently received (median 120 days later) anti-CD19 CAR T-cell therapy. The response rate

was 50%, and no survival data were available, with a median follow-up of only 6 months.²² Preclinical data recently demonstrated that prior treatment with TFAFA increased the antitumor activity of CAR T cells in vitro and reduced the severity of CRS.²³ These data have not been confirmed in clinical trials or real-life studies of the TFAFA-LEN association. We could not compare adverse event rates in the TL-pre-CAR-T set with those in the TL-post-CAR-T cohort because the 2 populations were different, and we could not identify a cohort of comparable control patients (who would have received treatments other than TFAFA-LEN before CAR T-cell therapy) in the DESCAR-T registry. In addition, TFAFA efficacy in treating B-cell malignancies involves several mechanisms beyond its interaction with CD19, including antibody-dependent cellular cytotoxicity, antibody-dependent cellular phagocytosis, and apoptosis induction in B cells through signaling pathways that do not involve immune cell mediation.^{17,35}

Concerning the role of CD19 (IHC) as a potential biomarker of response to treatment with TFAFA or CAR T-cell therapy, we were not able to study it with sufficient power, given that this staining is largely not performed in practice (in almost 3-quarters of patients) in a real-life setting. In France, neither histologically proving relapse/progression using rebiopsy nor checking CD19 positivity is required to prescribe CAR T-cell therapy, which certainly contributed to the low amount of data concerning CD19 status that we were able to collect. Additionally, it is important to note that the DESCAR-T registry does not collect histological reports or material. There is also no consensus on the specific IHC antibody to use for quantifying CD19 expression, nor on the cutoff percentage of stained cells necessary to consider a result positive. Therefore, for all these reasons, providing data on CD19 status, including retrospectively, is challenging. However, we observed that 7 of 52 (13.5%) patients in the TL-post-CAR-T cohort were negative for CD19 before CAR T-cell therapy, which did not prevent them from receiving this therapy, and that this status was not checked by another biopsy before the introduction of TFAFA-LEN as the next line of therapy. Overall, CD19 status has rarely been studied in patients receiving anti-CD19 treatment, and this marker is rarely correlated with the response to treatment.^{31,34} In a study in which loncastuximab-tesirine was used before CAR-T cell therapy, 10 of 14 patients were CD19-positive after loncastuximab-tesirine.²² Further studies are required to better decipher the role of CD19 as a biomarker to tailor anti-CD19 therapies and to better understand the clinical relevance of targeting CD19 at different lines of therapy in patients with LBCL. We will strongly encourage the biopsy of patients and prospective evaluation of CD19 antigen loss in future studies, given its relevance to many available and investigational anti-CD19 therapies.

To conclude, this analysis confirmed the poor outcome of patients who relapsed after CAR T-cell therapy in L3+. The efficacy of TFAFA-LEN administered directly after CAR T-cell therapy in this high-risk population was modest and similar to that of other salvage treatments, but the outcome improved when progression occurred later than 6 months after CAR T-cell therapy.

Acknowledgments

The authors thank the patients whose data were collected from the DESCAR-T registry and their families, and all the investigators and their staff involved in the data collection and analyses. The authors

18. Fitzgerald KN, Quesada AE, Von Keudell G, et al. CD19 epitope masking by tafasitamab leads to delays in subsequent use of CD19 CAR T-cell therapy in two patients with aggressive mature B-cell lymphomas. *Leuk Lymphoma*. 2022;63(3):751-754.
19. Ilieva K, Eberl M, Jaehrling J, et al. Preclinical study of CD19 detection methods post tafasitamab treatment. *Front Immunol*. 2023;14:1274556.
20. Horvei P, Sakemura R, Cox MJ, et al. Targeting of CD19 By tafasitamab does not impair CD19 directed chimeric antigen receptor T cell activity in vitro. *Blood*. 2019;134(suppl 1):2859.
21. Tabbara N, Gaut D, Oliai C, Lewis T, de Vos S. Anti-CD19 CAR T-cell therapy remission despite prior anti-CD19 antibody tafasitamab in relapsed/refractory DLBCL. *Leuk Res Rep*. 2021;16:100260.
22. Thapa B, Caimi PF, Ardeshna KM, et al. CD19 antibody-drug conjugate therapy in DLBCL does not preclude subsequent responses to CD19-directed CAR T-cell therapy. *Blood Adv*. 2020;4(16):3850-3852.
23. Sakemura RL, Manriquez Roman C, Horvei P, et al. CD19 occupancy with tafasitamab increases therapeutic index of CART19 cell therapy and diminishes severity of CRS. *Blood*. 2024;143(3):258-271.
24. Di Blasi R, Le Gouill S, Bachy E, et al. Outcomes of patients with aggressive B-cell lymphoma after failure of anti-CD19 CAR T-cell therapy: a DESCAR-T analysis. *Blood*. 2022;140(24):2584-2593.
25. Alarcon Tomas A, Fein JA, Fried S, et al. Outcomes of first therapy after CD19-CAR-T treatment failure in large B-cell lymphoma. *Leukemia*. 2023;37(1):154-163.
26. Spiegel JY, Dahiya S, Jain MD, et al. Outcomes of patients with large B-cell lymphoma progressing after axicabtagene ciloleucel therapy. *Blood*. 2021;137(13):1832-1835.
27. Ghobadi A, Munoz J, Westin JR, et al. Outcomes of subsequent anti-lymphoma therapies after second-line axicabtagene ciloleucel or standard of care in ZUMA-7. *Blood Adv*. 2024;8:2023011532.
28. Major A, Yu J, Shukla N, et al. Efficacy of checkpoint inhibition after CAR-T failure in aggressive B-cell lymphomas: outcomes from 15 US institutions. *Blood Adv*. 2023;7(16):4528-4538.
29. Sesques P, Houot R, Al Tabaa Y, et al. Glofitamab monotherapy in patients with non-Hodgkin B-cell lymphoma after failing CAR T-cell infusion: primary analysis of the BICAR study, a phase II LYSA study. *Blood*. 2023;142(suppl 1):893.
30. Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a novel, subcutaneous CD3xCD20 bispecific T-cell-engaging antibody, in relapsed or refractory large B-cell lymphoma: dose expansion in a phase I/II trial. *J Clin Oncol*. 2023;41(12):2238-2247.
31. Ruckdeschel A, Sabrina K, Topp MS, et al. Tafasitamab and lenalidomide in relapsed/refractory B-cell lymphoma: a multicenter retrospective real-world-study of patients from Germany and Austria. *Blood*. 2023;142(suppl 1):1771.
32. Iqbal M, Jagadeesh D, Chavez J, et al. Efficacy of CD19 directed therapies in patients with relapsed or refractory large B-cell lymphoma relapsing after CD19 directed chimeric antigen receptor T-cell therapy. *Bone Marrow Transplant*. 2024;59(2):211-216.
33. Crochet G, Iacoboni G, Couturier A, et al. Efficacy of CAR T-cell therapy is not impaired by previous bispecific antibody treatment in large B-cell lymphoma. *Blood*. 2024;144(3):2024024526.
34. Veeraputhiran M, Mehta A, Alencar AJ, Modi D, Voorhees TJ, Narkhede M. Tafasitamab (Tafa) plus lenalidomide (LEN) prior to chimeric antigen receptor T-cell (CAR-T) therapy in relapsed/refractory (R/R) diffuse large B-cell lymphoma (DLBCL): case series of 8 patients. *Blood*. 2022;140(suppl 1):12089-12091.
35. Papageorgiou SG, Thomopoulos TP, Liaskas A, Vassilakopoulos TP. Monoclonal antibodies in the treatment of diffuse large B-cell lymphoma: moving beyond rituximab. *Cancers*. 2022;14(8):1917.