



HAL
open science

Examining the relationships between cognition and auditory hallucinations: A systematic review.

Adrienne Bell, W. L. Toh, P. Allen, M. Cella, Renaud Jardri, F. Larøi, P. Moseley, S. L. Rossell

► To cite this version:

Adrienne Bell, W. L. Toh, P. Allen, M. Cella, Renaud Jardri, et al.. Examining the relationships between cognition and auditory hallucinations: A systematic review.. Australian & New Zealand Journal of Psychiatry, 2024, Aust N Z J Psychiatry, 58 (6), pp.48674241235849. 10.1177/00048674241235849 . hal-04739445

HAL Id: hal-04739445

<https://hal.univ-lille.fr/hal-04739445v1>

Submitted on 16 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Examining the relationships between cognition and auditory hallucinations: A systematic review

Australian & New Zealand Journal of Psychiatry
2024, Vol. 58(6) 467–497
DOI: 10.1177/00048674241235849

© The Author(s) 2024



Article reuse guidelines:
sagepub.com/journals-permissions
journals.sagepub.com/home/anp



Adrienne Bell¹ , Wei Lin Toh^{1,2,3} , Paul Allen⁴,
Matteo Cella^{5,6}, Renaud Jardri⁷, Frank Larøi^{8,9},
Peter Moseley¹⁰ and Susan L Rossell^{1,2}

Abstract

Objective: Auditory hallucinations (hearing voices) have been associated with a range of altered cognitive functions, pertaining to signal detection, source-monitoring, memory, inhibition and language processes. Yet, empirical results are inconsistent. Despite this, several theoretical models of auditory hallucinations persist, alongside increasing emphasis on the utility of a multidimensional framework. Thus, clarification of current evidence across the broad scope of proposed mechanisms is warranted.

Method: A systematic search of the Web of Science, PubMed and Scopus databases was conducted. Records were screened to confirm the use of an objective behavioural cognitive task, and valid measurement of hallucinations specific to the auditory modality.

Results: Auditory hallucinations were primarily associated with difficulties in perceptual decision-making (i.e. reduced sensitivity/accuracy for signal-noise discrimination; liberal responding to ambiguity), source-monitoring (i.e. self–other and temporal context confusion), working memory and language function (i.e. reduced verbal fluency). Mixed or limited support was observed for perceptual feature discrimination, imagery vividness/illusion susceptibility, source-monitoring for stimulus form and spatial context, recognition and recall memory, executive functions (e.g. attention, inhibition), emotion processing and language comprehension/hemispheric organisation.

Conclusions: Findings were considered within predictive coding and self-monitoring frameworks. Of concern was the portion of studies which – despite offering auditory-hallucination-specific aims and inferences – employed modality-general measures, and/or diagnostic-based contrasts with psychologically healthy individuals. This review highlights disparities within the literature between theoretical conceptualisations of auditory hallucinations and the body of rigorous empirical evidence supporting such inferences. Future cognitive investigations, beyond the schizophrenia-spectrum, which explicitly define and measure the timeframe and sensory modality of hallucinations, are recommended.

Keywords

Auditory hallucinations, hearing voices, cognitive models, cognitive mechanisms

¹Centre for Mental Health and Brain Sciences, Swinburne University of Technology, Melbourne, VIC, Australia

²Psychiatry, St Vincent's Hospital, Melbourne, VIC, Australia

³Department of Psychology, Alfred Health, Melbourne, VIC, Australia

⁴Department of Neuroimaging, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

⁵South London and Maudsley NHS Foundation Trust, Maudsley Hospital, London, UK

⁶Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

⁷University of Lille, INSERM U-1172, Plasticity and Subjectivity Team, Lille Neuroscience and Cognition Research Centre, Fontan Hospital, CHU Lille, Lille, France

⁸Department of Psychology, University of Oslo, Oslo, Norway

⁹Psychology and Neuroscience of Cognition Research Unit, University of Liege, Liege, Belgium

¹⁰Department of Psychology, Northumbria University, Newcastle-Upon-Tyne, UK

Corresponding author:

Adrienne Bell, Centre for Mental Health and Brain Sciences, Swinburne University of Technology, John Street, Hawthorn, Melbourne, VIC 3122, Australia.

Email: adriennebell@swin.edu.au

Introduction

Auditory hallucinations (AHs), also known as hearing voices, are recognised as the perception of sound in the absence of corresponding external stimuli, often characterised by a sense of reality and lack of control (David, 2004). Theoretical models postulate that AHs reflect the externalisation of mental events, such as inner speech (Jones and Fernyhough, 2007) or intrusive/fragmented memories (Waters et al., 2006), and/or the exacerbation of environmental cues via increased hypervigilance or attributed salience (Dodgson and Gordon, 2009; Kapur, 2003). Similarly, broad frameworks of AHs suggest common difficulties in the recognition of self-generated mental events (Waters et al., 2012b), hemispheric organisation for language processing (Hugdahl et al., 2012) and the weighting of perceptual input against prior expectation (Corlett et al., 2019; Sterzer et al., 2018). A range of cognitive mechanisms have been implicated, including challenges in auditory signal detection and response biases (Bentall and Slade, 1985; McLachlan et al., 2013), source-monitoring (Bentall, 1990; Woodward et al., 2007), memory function (Brébion et al., 2020), attention modulation and inhibition (Badcock et al., 2005; Waters et al., 2003), language processes (Docherty, 2012; Green et al., 1994) and emotion detection (Alba-Ferrara et al., 2013; Rossell and Boundy, 2005; Shea et al., 2007). However, a notable portion of non-significant findings have also been reported (e.g. Brébion et al., 2008; Johns et al., 2001; McKay et al., 2000; McLachlan et al., 2013; Pinheiro et al., 2016; Rocca et al., 2006; Rossell et al., 2001; Schnakenberg Martin et al., 2018; Toh et al., 2020).

Previous meta-analyses, focusing on specific domains of cognition, have demonstrated moderate-to-large associations between hallucinations and a self-monitoring bias (Waters et al., 2012b), increased externalising source-monitoring errors (Brookwell et al., 2013; Damiani et al., 2022) and reduced left-hemispheric language lateralisation (Ocklenburg et al., 2013). Source-monitoring difficulties were observed in the context of auditory stimuli and imagined mental events (Damiani et al., 2022), and identified in spite of intact broader memory function (Waters et al., 2012b). While such reviews have provided valuable integration of findings, some methodological inconsistencies are evident. For example, the sensory modality (e.g. auditory and/or additional senses) and specified timeframe of hallucinations were not always identified. In addition, group contrasts based on diagnostic status (e.g. schizophrenia vs no history of mental illness) commonly contributed to findings. Thus, it is difficult to discern the extent to which findings are relevant to AHs specifically, as opposed to hallucinations of other forms, psychosis syndromes or broader diagnostic groups.

Numerous narrative and systematic reviews have summarised a breadth of cognitive functions in the context of proposed theoretical models (e.g. Seal et al., 2004; Tracy and Shergill, 2013; Uptegrove et al., 2016; Waters et al.,

2012a). These papers emphasise that competing theoretical accounts are not mutually exclusive, and that no single mechanism can sufficiently account for AH emergence. Thus, a multidimensional framework has been advanced, along with recommendations for transdiagnostic investigation at the symptom level. While inconsistencies regarding the definition and measurement of AHs have been acknowledged, limited consideration has been given to how this may affect the substantiation of corresponding theoretical models. Likewise, poor measurement practices relating to cognitive task development make interpretation of disparate findings difficult (Smailes et al., 2022). Subsequently, a lack of clarity and consensus remains when interpreting and integrating the abundance of cognitive studies at a conceptual level. Thus, a systematic review of the relationships between AHs and the range of proposed cognitive mechanisms is warranted.

Study aims

The current systematic review aimed to synthesise the literature examining cognition in relation to the presence and severity of AHs accompanied by a psychiatric diagnosis. We omitted hallucination proneness within psychologically healthy populations, to focus on cognitive profiles likely associated with a need for care. Cognition was considered across the broad range of mechanisms, with the aim of comparing the relative proportion and nature of evidence representing each. To address existing gaps, the sensory modality and explicit timeframe of AHs were carefully considered. Inferences regarding the alignment of empirical findings to theoretical frameworks were offered, where possible.

Method

Protocol pre-registration

A protocol for the systematic review was developed in line with the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P; Moher et al., 2015), and registered with the International Prospective Register of Systematic Reviews (PROSPERO; CRD42020148907).


Search strategy

A systematic search of the Web of Science, Scopus and PubMed databases was employed to identify peer-reviewed empirical records written in the English language, published from 1 January 1980 to 2 August 2022. Details of the search strategy, including full syntax, are reported in Supplemental Appendix A.

Study selection

The search yielded 22,391 records, after duplicate removal. A single reviewer (A.B.) screened the title, abstract and keywords against the eligibility criteria. Records were

Figure 1. PRISMA record screening.



Note: The following 11 studies met initial criteria but were omitted from synthesis as no AH-specific analyses were available (i.e. clinical AHs vs healthy control contrasts only) (Jacobsen et al., 2019; Løberg et al., 2004; Moseley et al., 2022; Olsson and Nielzen, 1999a, 1999b; Papageorgiou et al., 2003; Powers et al., 2017; Sun et al., 2021; Zhuo et al., 2020), populations fell outside the schizophrenia-spectrum (Strawson et al., 2021) and data pertained to phenomenological variations of AHs only (Stephane, 2019).

retained if they met the following criteria: (1) part of an original group design empirical investigation,¹ (2) an adult participant sample (i.e. 18–65 years), (3) AHs accompanied by a psychiatric diagnosis (i.e. individuals without a psychiatric diagnosis were omitted), (4) participant sample omitting organic illness or injury (e.g. neurological, substance-induced or other major medical conditions), (5) cognition investigated with reference to AHs, by means of an objective behavioural task and (6) the presence/severity of AHs was identified via appropriate clinical judgement and/or a valid measure specific to, or adapted for, the auditory modality (current and lifetime timeframes eligible). Independent reviewers (A.B., S.L.R.) screened the full-text articles retained at Stage 2. Discrepancies in eligibility were resolved during regular meetings, with mediation from a third reviewer (W.L.T.).

Following screening, 101 records met the inclusion criteria. Several records were omitted from data synthesis given that (1) no AH-specific data was available ($n=9$) (i.e. group contrasts between clinical AHs and healthy control

groups only), (2) as analyses pertained to phenomenological variations of AHs rather than presence or severity ($n=1$) or (3) AHs were examined in the context of borderline personality disorder ($n=1$). Although a transdiagnostic approach was initially intended, the latter instance was the only study to fall outside the schizophrenia-spectrum (i.e. schizophrenia or mood related diagnosis with psychotic features), and was therefore considered clinically distinct and unsuitable for synthesis. Subsequently, 90 studies remained for data extraction (see Figure 1).

Data extraction

The following data were extracted: record identifying information, participant groups (psychiatric diagnosis and/or AH presence/absence), AH frequency/severity, cognitive task(s) and associated outcome variables, analyses and data relevant to relationships between AHs and cognition (e.g. direction, effect sizes and significance coefficients, where available). Group sample characteristics (e.g. age, sex)

were extracted where possible. Given the extent of relevant data, no attempts were made to contact authors to obtain missing information. Data were summarised into three streams: group contrasts, correlations and other analyses (e.g. regression, mediation and discriminant function). Where possible, available effect sizes were converted to Cohen's d (Lenhard and Lenhard, 2016), with $d=0.2$ interpreted as small, $d=0.5$ as medium and $d=0.8$ as large in magnitude (Cohen, 1988). That said, these benchmarks are arbitrary, and small effect sizes can be equally important (Lakens, 2013). Furthermore, given typically limited sample sizes, effect confidence intervals are likely to be wide and should be interpreted with caution.

Quality assessment

The Newcastle Ottawa Scale (NOS; Wells et al., 2000) was used as a template for methodological quality assessment. Key factors were examined, and studies were assigned an overall quality score from 1 to 5. Categorical quality rankings were developed, based on the median and quartile split of total scores. These consisted of low (2.50–3.49), moderate (3.50–4.06), high (4.07–4.39) and very high (4.40–5.00). See Supplemental Appendix B for checklist items and anchors, alongside associated scores and categorical rankings.

Data analysis

To aid interpretation, studies were tabulated by broad cognitive domain: perceptual decision-making and imagery ($n=30$), source-monitoring and context memory ($n=28$), general memory ($n=29$), language processing, production and hemispheric organisation ($n=14$), executive function ($n=23$), and emotion processing and social cognition ($n=9$) (details in Supplemental Appendix C). Subsequently, refined cognitive sub-domains were formed by grouping results associated with similar outcome variables. These groupings accounted for all data in the results tables, except for the executive function domain, which had too much variation to meaningfully group. Hence, only key sub-domains were considered:² attention, inhibition and cognitive flexibility/set-shifting.

As outlined in the pre-registered protocol, a minimum of 10 studies per cognitive sub-domain was required to proceed with meta-analysis. After grouping studies by homogeneous task design, variable definition and analysis form, only one cognitive sub-domain met this criterion (i.e. 14 group contrasts for internal–external source misattributions). Given that the primary aim of the systematic review was to compare results across the range of proposed cognitive mechanisms, and that meta-analysis of source-monitoring data has been well documented (see Brookwell et al., 2013; Damiani et al., 2022; Waters et al., 2012b), performing a quantitative

synthesis for this single sub-domain was considered unsuitable. It is possible to conduct meta-analysis of broader, collapsed groups of data, with follow-up statistics to accommodate study deviations (e.g. meta-regression, subgroup and/or sensitivity analysis). However, caution has been drawn to implementing this approach to heterogeneous datasets, due to challenges in defining these points of variation (e.g. Imrey, 2020). Within this literature, we observed notable heterogeneity in the definition of constructs, the use of novel experimental paradigms and the interpretation of numerous outcome variables. Thus, we chose to proceed with a narrative synthesis to allow for consideration of study nuances, as documented comprehensively within the results tables. To facilitate the integration of a broad array of data, a counting method was adopted, to identify the number of studies with significant, mixed and non-significant findings pertaining to each cognitive sub-domain, as relevant to the three analysis streams (see Supplemental Appendix C for details).

Results

Sample characteristics

Across 81 unique samples,³ a total of 4933 individuals with a schizophrenia-spectrum diagnosis were represented in AH-specific groups/analyses, with a mean age of 34 years and gender ratio of 13 males:7 females.


AH status

Many studies explored cognition in the context of current AHs ($n=50$), and to a lesser extent, lifetime ($n=18$) and past AHs ($n=5$). A notable portion did not explicitly specify the timeframe ($n=23$). Only a small number of studies explored several AH status variations ($n=6$).

Cognitive domains

The full set of data extracted from the 90 studies, arranged by cognitive domain, can be seen in Tables 1–6; with the proportions of significant, mixed and non-significant findings displayed in Figure 2. Of the 21 identified sub-domains, in 9 cases, the number of investigations with any significant associations to AHs (i.e. significant or mixed results overall) outweighed the number of non-significant findings: spatial context attribution (3/3), language comprehension (5/7), signal detection response bias (7/10), temporal context attribution (4/6), working memory (9/13), verbal fluency/linguistic cohesion (5/8), signal-noise discrimination sensitivity (9/16), internal–external source attribution (11/21) and hemispheric language organisation (4/7). Below is a summary of the significant findings within each domain.

Figure 2. The proportion of non-significant, mixed and significant results per cognitive sub-domain, identified via group contrasts (AH presence), correlations (AH severity) and regression/other analyses.



Note: PD: perceptual decision-making and imagery; SM: source-monitoring and context memory; GM: general memory; LP: language processing, production and hemispheric organisation; EF: executive function; EP: emotion processing and social cognition. Given that a single study often explored multiple cognitive sub-domains, utilising a range of analyses for each, the number of investigations does not equal the number of applicable studies. Similarly, due to overlap, the number of studies relevant to each sub-domain does not sum to the number of studies constituting each overarching domain.

Perceptual decision-making and imagery. This domain encompassed data relevant to the detection of sensory input, primarily within the auditory modality, but also visual, and to a lesser extent tactile stimuli. Key variables reflected signal detection and/or noise discrimination sensitivity/accuracy, discrimination of perceptual qualities and response biases in the expectation of input. The intensity, integration and weighting of imagery in relation to these perceptual processes are also reflected here (see Table 1). Significant results suggested an association between AH presence/severity and reduced sensitivity/accuracy for signal-noise discrimination ($d=0.47-1.50$), liberal responding in the face of stimulus ambiguity ($d=0.42-2.42$), reduced stimulus feature discrimination capacity ($d=0.04-1.00$) and increased vividness/weighting of imagery or illusion susceptibility ($d=1.96-2.84$). Conversely, one study noted superior stimulus feature discrimination in participants with current, relative to past AHs, citing exaggerated environmental scanning as a possible contributor (Schneider and Wilson, 1983). In addition, AH severity was observed as a predictor of spatial detection sensitivity ($d=1.5$), and signal expectation bias ($d=0.33$); while signal detection also predicted the AH presence ($\chi^2=12.1$).

Source-monitoring and context memory. This domain reflected a complex set of processes involved in the monitoring of, and memory for, contextual aspects of presented stimuli and mental events. These included temporal order, visuospatial location, external form (e.g. picture vs word stimulus), internal form (e.g. thought vs said), and the differentiation of internal-external source (e.g. imagined/read vs heard; self vs other speech/action). Primary variables pertained to the number of source misattributions, as well as error rates and accuracy scores (see Table 2). In most instances the direction of internal-external source-monitoring was specified (e.g. externalising vs internalising misattributions), however, in some cases this was not discernible ($n=2$). Significant results suggested an association between AH presence/severity and greater challenges in distinguishing: temporal presentation order ($d=0.64-0.85$), spatial presentation location ($d=1.04-1.12$), internal-external stimulus source ($d=0.57-2.82$), internal mental event form ($d=0.49-1.12$) and external stimulus form ($d=1.01-1.12$). In addition, AH severity was observed as a predictor of misattribution for temporal context ($d=1.22$), spatial context ($\beta=0.53$) and internal-external source ($d=1.25$).

Table 1. Perceptual decision-making and imagery.

Author (year)	Groups (n)	Task type and sensory mode	Analyses	Results
Aleman et al. (2003)	SCZ (22)	Auditory & visual feature discrimination Auditory signal detection Imagery-perception interaction	Correlations ^a	↓ AH score ^b × ↑ auditory imagery-perception interaction: $p = 0.01$, $d = 1.96$ AH severity × auditory signal detection, auditory discrimination (imagery, perception) & visual discrimination (imagery, perception): all NS
Antonova et al. (2021)	SCZ-LAH (18) SCZ-NLAH (11)	Visual signal detection	Group contrasts	Accuracy, error rate, omissions: SCZ-LAH vs SCZ-NLAH (left/right-lateral, left/right-medial visual fields): all NS
Blakemore et al. (2000)	SCZ/SAD-CAH (15) SCZ/SAD-NCAH (23)	Tactile intensity detection	Group contrasts	Frequency of typical (external > internal) vs atypical (external < internal) responses: SCZ/SAD-CAH < SCZ/SAD-NCAH: $p < 0.05$, $d = 0.72$
Bristow et al. (2014)	SCZ (51)	Auditory & visual signal detection	Correlations ^a	↑ CAH severity × ↓ auditory perceptual bias ^d : $p < 0.05$, $d = 0.42$ ↑ CAH severity × ↓ visual perceptual bias ^d : $p < 0.05$, $d = 0.48$
Catts et al. (1980)	SCZ-LAH (12) SCZ-NLAH (12)	Auditory illusion/conditioning	Group contrasts Regression/other	Frequency of total & different illusions: SCZ-LAH vs SCZ-NLAH: both NS Chi-Square: Relationship between AH status (LAH/NLAH) & level of phonetic illusion-target connection (high/low): NS
Chandiramani and Varma (1987)	SCZ-CAH (20) SCZ-NCAH (20)	Visual imagery generation	Group contrasts	Imagery vividness score: SCZ-CAH vs SCZ-NCAH: NS
Chhabra et al. (2012)	SCZ-CAH (33) SCZ-NCAH (32)	Auditory feature discrimination	Group contrasts	Discrimination (pitch, resonance): SCZ-CAH vs SCZ-NCAH: NS
Chhabra et al. (2016)	SCZ-CAH (15)	Auditory signal detection	Correlations ^a	CAH severity × hit rate, discriminability, response bias & false alarm rate: all NS
Chhabra et al. (2022)	SCZ-AH (17)	Auditory signal detection	Correlations ^a	↑ AH severity × ↓ discrimination sensitivity: $p = 0.04$, $d = 1.15$
Daalman et al. (2012)	SCZ-CAH (40)	Auditory signal expectation	Correlations ^a	CAH severity × top-down expectation errors: NS
Evans et al. (2000)	SCZ-LAH (12) SCZ-NLAH (7)	Auditory illusion/conditioning Auditory imagery generation	Group contrasts	Imagery: letter-number string (with/without sound interference, with/without articulation interference), pitch, phoneme & rhyme judgements: SCZ-LAH vs SCZ-NLAH: all NS Illusion: verbal transformations: SCZ-LAH vs SCZ-NLAH: NS
He et al. (2022)	SCZ-CAH (47) SCZ-NLAH (33)	Visual-tactile illusion/conditioning	Group contrasts Correlations ^a	Proprioceptive drift: SCZ-CAH < SCZ-NLAH (synchronous): $p < 0.001$ ^e SCZ-CAH < SCZ-NLAH (asynchronous): $p < 0.015$ ^e CAH severity × proprioceptive drift: NS
Heilbrun et al. (1983)	LAH (16) NLAH (14)	Auditory & visual imagery generation Auditory spatial detection	Group contrasts	Auditory-visual imagery preference: LAH < NLAH: $p < 0.05$, $d = 0.81$ Voice spatial discrepancy: LAH (IA) ^g > LAH (EA) ^g : $p < 0.05$, $d = 0.40$ ^h LAH (IA) > NLAH (IA): $p < 0.05$, $d = 0.04$ ^h LAH (IA) > NLAH (EA): $p < 0.05$, $d = 0.34$ ^h Visual & auditory imagery vividness: LAH vs NLAH: both NS
Heilbrun and Blum (1984)	CAH (22) NCAH (22)	Auditory signal detection	Group contrasts	Number of false alarms: CAH (EA) ^g > CAH (IA) ^g : $p < 0.05$, $d = 1.62$ ^h CAH (EA) > NCAH (EA): $p < 0.05$, $d = 1.52$ ^h CAH (EA) > NCAH (IA): $p < 0.05$, $d = 1.77$ ^h Ambiguity tolerance, detection bias: CAH vs NCAH: NS

(Continued)

Table 1. (Continued)

Author (year)	Groups (n)	Task type and sensory mode	Analyses	Results
Hoffman (1999)	Study 1: SCZ-CAH (14) SCZ-NCAH (17) Study 2: SCZ-CAH (22) SCZ-NCAH (25)	Auditory masked speech tracking	Group contrasts	Correct words: Study 1 & 2. SCZ-CAH < SCZ-NCAH ^{ei} Study 1. Incorrect words: SCZ-CAH vs SCZ-NCAH: NS Study 2. Incorrect words: SCZ-CAH > SCZ-NCAH (low mask) ^{ei}
Hoffman et al. (1999)	SSD-CAH (24) SSD-NCAH (28)	Auditory masked speech tracking	Group contrasts Correlations Regression/other	Correct words: SSD-CAH < SSD-NCAH (low mask): $d = 0.47^{\text{hi}}$ SSD-CAH < SSD-NCAH (moderate mask): $d = 0.85^{\text{i}}$ Incorrect words: SSD-CAH > SSD-NCAH (low mask): $d = 0.78^{\text{hi}}$ SSD-CAH < SSD-NCAH (moderate mask): $d = 0.51^{\text{i}}$ ↑ CAH severity $x \rightarrow$ correct words: $p < 0.001$, $d = 1.07$ Discriminant function analysis: correct & incorrect word rate (moderate mask), correctly classified the CAH status of 82.2% of cases: $p < 0.002$, $\chi^2 = 12.1^{\text{i}}$
Ishigaki and Tanno (1999)	SCZ/SAD-CAH (11) SCZ/SAD-NCAH (11)	Visual signal detection	Group contrasts	Decision bias ^d : SCZ/SAD-CAH < SCZ/SAD-NCAH: $p = 0.027$, $d = 1.33$ Discriminating ability: SCZ/SAD-CAH vs SCZ/SAD-NCAH: NS
Kot and Serper (2002)	SCZ-CAH (15) SCZ-NCAH (15)	Auditory illusion/conditioning	Group contrasts Correlations	Frequency of illusions: SCZ-CAH > SCZ-NCAH (conditioning trials): $p = 0.01$, $d = 2.84$ SCZ-CAH > SCZ-NCAH (extinction trials): $p = 0.001$, $d = 2.35$ Latency & duration of illusion: SCZ-CAH vs SCZ-NCAH: both NS
Laloux et al. (2018)	SCZ (15)	Visual signal detection & imagery generation	Correlations ^a	CAH frequency ^k \times reaction time (visual stimuli, visual imagery) & % errors (visual stimuli, visual imagery): both NS
Lee et al. (2004)	SCZ-CAH (24) SCZ-NCAH (25)	Auditory masked speech tracking	Group contrasts	Correct words: SCZ-CAH < SCZ-NCAH (low mask): $p = 0.001$, $d = 1.50$ SCZ-CAH < SCZ-NCAH (moderate mask): $p = 0.004$, $d = 0.87$ SCZ-CAH vs SCZ-NCAH (high mask): NS
Li et al. (2002)	SCZ-CAH (28) SCZ-NLAH (16)	Auditory signal detection	Regression/other	CAH presence did not contribute to detection sensitivity, detection bias, or discrimination ability: NS
McKay et al. (2000)	SCZ/SAD-CAH (22) SCZ/SAD-NLAH (16)	Auditory signal detection & feature discrimination	Group contrasts	Monaural filtered speech score: SCZ/SAD-CAH < SCZ/SAD-NLAH (left ear): $p < 0.007$, $d = 0.93$ SCZ/SAD-CAH vs SCZ/SAD-NLAH (right ear): NS Binaural fusion, rapidly alternating speech perception, monaural speech discrimination (white noise + left/right ear) (competing speech + left/right ear), frequency tone pattern (verbal response + left/right ear) (hummed response + left/right ear) (hummed-verbal response), competing environmental sounds % errors (left/right ear): SCZ/SAD-CAH vs SCZ/SAD-NLAH: all NS
McKay et al. (1981)	LAH (10) NLAH (20)	LNNB subscales	Group contrasts	Rhythm reproduction: LAH > NLAH: $p < 0.05$, $d = 1.0^{\text{b}}$ Tactile sensitivity: LAH vs NLAH: NS
McLachlan et al. (2013)	SCZ/SAD-LAH (19) SCZ/SAD-NLAH (15)	Auditory feature discrimination	Group contrasts	Pitch discrimination sensitivity: SCZ/SAD-LAH < SCZ/SAD-NLAH (0, 1, 2.5, 5, 10, 20%): $p < 0.05^{\text{e}}$ Auditory streaming sensitivity: SCZ/SAD-LAH < SCZ/SAD-NLAH (easy): $p < 0.05^{\text{e}}$ Modulation discrimination sensitivity: SCZ/SAD-LAH vs SCZ/SAD-NLAH (2, 40, 120 Hz): NS SCZ/SAD-LAH vs SCZ/SAD-NLAH (difficult): NS

(Continued)

Table 1. (Continued)

Author (year)	Groups (n)	Task type and sensory mode	Analyses	Results
Perrin et al. (2018)	SCZ-AH (13)	Auditory spatial discrimination	Regression/other	AH severity contributed to spatial detection sensitivity at the para-central location: $p = 0.041$, $d = 1.5$, but not at the left and right lateral hemifield locations: NS
Schmakenberg Martin et al. (2018)	SCZ-CAH (20) SCZ-NCAH (24)	Auditory feature discrimination	Group contrasts Correlations	Percent correct for rhythm, frequency, duration, intensity & temporal discrimination: SCZ-CAH vs SCZ-NCAH: NS CAH intensity ^k × all auditory feature discrimination variables: NS
Schneider and Wilson (1983)	SCZ-CAH (12) SCZ-PAH (12) SCZ-NLAH (12)	Auditory & visual feature discrimination	Group contrasts	Error rate: SCZ-CAH < SCZ-PAH, SCZ-NLAH (auditory/visual + infrequent stimuli): $p < 0.05^e$ SCZ-CAH vs SCZ-PAH vs SCZ-NLAH (auditory/visual + frequent stimuli): NS Reaction time for correct responses: SCZ-CAH < SCZ-PAH, SCZ-NLAH (auditory): $p < 0.05^e$ SCZ-CAH vs SCZ-PAH vs SCZ-NLAH (visual): NS
Selvaraj et al. (2021)	SCZ-LAH (24)	Auditory signal detection	Correlations ^a	↑ LAH severity × ↑ false alarm rate: $p = 0.028$, $d = 1.01$ ↑ LAH severity × ↓ discriminability index: $p = 0.012$, $d = 1.17$ ↑ LAH severity × ↓ response bias ^d : $p = 0.009$, $d = 1.23$
Yang et al. (2021)	SCZ-CAH (32) SCZ-NLAH (31)	Auditory signal expectation	Group contrasts Correlations Regression/other	Detection sensitivity: SCZ-CAH > SCZ-NLAH (tone pattern-target consistent): $p = 0.009^e$ SCZ-CAH < SCZ-NLAH (tone pattern-target inconsistent): $p = 0.002^e$ SCZ-CAH vs SCZ-NLAH (preceding tone-target consistent & inconsistent): NS ↑ CAH severity × ↑ detection sensitivity (tone pattern-target consistent): $p < 0.001$, $d = 1.41$ ↑ CAH severity × ↓ detection sensitivity (tone pattern-target inconsistent): $p = 0.003$, $d = 1.19$ ↑ CAH severity × ↑ detection sensitivity (consistent vs inconsistent): $p < 0.001$, $d = 2.42$ CAH severity contributed to detection sensitivity (tone pattern-target consistent): $p = 0.001$, $d = 0.33$
Zenisek et al. (2015)	BPAD-LAH (17) BPAD-NLAH (13)	Auditory feature discrimination	Group contrasts	Discrimination sensitivity (pitch): BPAD-LAH vs BPAD-NLAH: NS

Diagnoses: BPAD: bipolar-affective disorder, SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizoprenia-spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Notes: Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Task conditions are listed in parentheses following relevant variables. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical AH vs healthy control only).

^bAH variable where higher scores indicate lower severity.

^cLower scores indicate a larger effect of auditory imagery over perceptual input, while higher scores indicate less weighting of imagery.

^dLower scores indicate a liberal pattern of responding, while higher scores indicate a more conservative pattern of responding.

^eEffect size values, means and standard deviations not available. Unable to calculate Cohen's d .

^fWhere positive scores indicate a preference for auditory imagery, while negative scores indicate a preference for visual imagery.

^gEA: predisposition to external attention; IA: predisposition to internal attention.

^hEffect size values not available. Cohen's d calculated from means and standard deviations.

ⁱSignificance value/level not reported, described as 'significant' by authors.

^jEffect size unable to be converted to Cohen's d .

^kAdditional results involving phenomenological AH variables not extracted.

^lCognitive variable where higher scores indicate poorer rhythm reproduction ability.

Table 2. Source-monitoring and context memory.

Author (year)	Groups (n)	Task type and sensory mode	Stimuli source options	Analyses	Results
Aleman et al. (2003)	SCZ (22)	Auditory source monitoring	Internal vs external (imagined, heard, new)	Correlations ^a	↑ AH score ^b × ↓ source attribution accuracy (immediate): $p = 0.01$, $d = 2.02$, (delayed): $p = 0.02$, $d = 1.81$
Allen et al. (2004)	SCZ-CAH (15) SCZ-NCH (13)	Auditory pre-recorded speech distortion	Internal vs external (own speech, other speech)	Group contrasts	Misattributions: SCZ-CAH > SCZ-NCH (distorted + self-speech): $p < 0.001$, $d = 1.59$ SCZ-CAH vs SCZ-NCH (distorted/non-distorted + other speech): NS Unsure responses: SCZ-CAH vs SCZ-NCH (distorted/non-distorted + self/other speech): NS
Allen et al. (2007)	SCZ-CAH (10) SCZ-NLAH (10)	Auditory pre-recorded speech distortion	Internal vs external (own speech, other speech)	Group contrasts	Misattributions: SCZ-CAH > SCZ-NLAH (self-speech): $p = 0.001$ ^c SCZ-CAH vs SCZ-NLAH (distorted/non-distorted + other-speech) (positive/negative/neutral valence): all NS
Bentall et al. (1991)	SCZ-CAH (22) SCZ-NLAH (16)	Auditory source monitoring	Internal vs external (said, heard, new)	Group contrasts	Misattributions: SCZ-CAH > SCZ-NLAH (self-generated + low-probability words): $p < 0.001$ ^c
Brébion et al. (2007a)	SCZ-AH (18) SCZ-NAH (23)	Auditory-visual free recall	Temporal context (list 1 or 2)	Group contrasts Regression/other	Temporal context misattributions: SCZ-AH > SCZ-NAH: $p < 0.015$, $d = 0.85$ Increased AH severity contributed to increased temporal context misattributions (overall): $p < 0.025$, $d = 1.22$, but not for (high/low frequency only): NS
Brébion et al. (2007b)	SCZ (24)	Visual spatial context memory	Spatial context (location)	Correlations	↑ AH severity × ↑ spatial context misattribution (black-white): $p < 0.025$, $d = 1.12$ AH severity × spatial context misattribution (colour): NS
Brébion et al. (2008)	SCZ-AH (17) SCZ-NAH (24)	Visual source monitoring	External form (picture, word)	Group contrasts Correlations	Misattributions: SCZ-AH vs SCZ-NAH = NS AH severity × misattributions (words as pictures): NS
Brébion et al. (2009)	SCZ-AH (12) SCZ-NAH (29)	Auditory free recall	Internal vs external (list, non-list) Temporal context (lists 1–4)	Group contrasts Correlations	Extra-list intrusions: SCZ-AH > SCZ-NAH: $p < 0.03$, $d = 0.79$ Intra-list intrusions: SCZ-AH vs SCZ-NAH: NS ↑ AH severity × ↑ extra-list intrusions: $p < 0.05$, $d = 0.75$ AH severity × intra-list intrusions: NS
Brébion et al. (2016)	SCZ (57)	Auditory-visual free recall & recognition	Internal vs external (list, non-list)	Regression/other ^a	Increased AH severity contributed to increased extra-list errors (male participants): $p < 0.01$, $d = 1.25$, but not female participants: NS
Brébion et al. (2019)	SCZ (59)	Visual spatial context memory	Spatial context (location)	Correlations ^a Regression/other	↑ AH severity × ↑ association misattributions (black-white + delayed + female participants): $p < 0.05$, $d = 1.04$ AH severity × association errors (colour + immediate/delayed + female participants) (black-white + immediate + female participants) (colour/black-white + immediate/delayed + male participants): all NS AH severity × response bias & location misattributions (immediate/delayed + colour/black-white + male/female participants): all NS Increased AH severity contributed to increased association misattributions (black-white + immediate + female participants): $p < 0.025$, $\beta = 0.53$ ^d , but not for (black-white + immediate + female participants) or (colour + immediate/delayed + female participants) or (colour/black-white + immediate/delayed + male participants): all NS AH severity did not contribute to location misattributions (colour/black-white + immediate/delayed + male/female participants): all NS

(Continued)

Table 2. (Continued)

Author (year)	Groups (n)	Task type and sensory mode	Stimuli source options	Analyses	Results
Brébin et al. (2020)	SCZ (43)	Visual temporal context memory	Temporal context (run 1 or 2)	Regression/other ^a	Increased AH severity contributed to increased temporal context errors (black-white); $p < 0.015$, $d = 1.01$, but not temporal context errors (colour): NS
Brunelin et al. (2006)	SCZ-CAH (30) SCZ-NCAH (31)	Auditory source monitoring	Internal form (said, imagined) Internal vs external (said, heard)	Group contrasts	Say-imagine misattributions: SCZ-CAH vs SCZ-NCAH: NS Hear-imagine externalising misattributions: SCZ-CAH > SCZ-NCAH: $p = 0.029$, $d = 0.57$ Hear-imagine internalising misattributions: SCZ-CAH vs SCZ-NCAH: NS
Chhabra et al. (2013)	SCZ-CAH (32) SCZ-NCAH (32)	Auditory source monitoring	External form (voice-word pair same or different)	Group contrasts Correlations	Voice-word binding: SCZ-CAH vs SCZ-NCAH: NS CAH frequency ^e × voice-word binding (novel voice, novel word): NS
Docherty (2012)	SCZ/SAD (110)	Auditory source monitoring	Internal form (thought, said, new) External form (man, woman, new)	Correlations ^a	↑ CAH severity × ↓ internal source memory accuracy: $p < 0.03$, $d = 0.49$ CAH severity × external source memory accuracy: NS
Franck et al. (2000)	SCZ-CAH (7) SCZ-PAH (5)	Auditory-visual source monitoring	Internal form (thought, said, new)	Group contrasts	Accuracy: SCZ-CAH < SCZ-PAH: $p = 0.009$, $d = 2.12$ Thought-said misattributions: SCZ-CAH vs SCZ-PAH: NS
Hubl et al. (2020)	SSD-AH (11) SSD-NLAH (9)	Auditory immediate speech feedback	Internal accuracy (voice delayed, not delayed)	Group contrasts	Accuracy, sensitivity & response bias: SSD-AH vs SSD-NLAH: all NS
Johns et al. (2001)	SCZ-AH (10) SCZ-NAH (8)	Auditory immediate speech feedback	Internal vs external (own speech, other speech, unsure)	Group contrasts	Overall errors & externalising misattributions: SCZ-AH vs SCZ-NAH: both NS
Johns et al. (2006)	SCZ/SAD-CAH (15) SCZ/SAD-PAH (15) SCZ/SAD-NLAH (15)	Auditory immediate speech feedback	Internal vs external (own speech, other speech, unsure)	Group contrasts	Overall errors: SCZ/SAD-CAH > SCZ/SAD-PAH (own voice + moderate distortion): $p = 0.01$, $d = 2.82$, (own voice + severe distortion): $p = 0.034$, $d = 0.81$, SCZ/SAD-CAH vs SCZ/SAD-NLAH (own voice + all distortions) (other voice + no distortion): all NS
Lee et al. (2013)	SCZ (26)	Visual picture evaluation	External form ('real', 'unreal')	Correlations ^a	Misattributions: SCZ/SAD-CAH > SCZ/SAD-PAH (own voice + moderate distortion): $p = 0.004$, $d = 1.23$ (own voice + severe distortion): $p = 0.03$, $d = 0.82$, SCZ/SAD-CAH > SCZ/SAD-NLAH (negative valence): $p = 0.03$, $d = 0.84$, SCZ/SAD-CAH vs SCZ/SAD-NLAH (own voice + all distortions) (other voice + no distortion): all NS
Morrison and Haddock (1997)	SCZ-CAH (15) SCZ-NCAH (15)	Auditory source monitoring	Internal vs external (heard, said, both)	Group contrasts	↑ AH severity × ↓ reality evaluation accuracy ('real' pictures): $p = 0.02$, $d = 1.12$, & ('un-real' pictures): $p = 0.03$, $d = 1.01$ Misattributions: SCZ-CAH vs SCZ-NCAH: NS
Pinheiro et al. (2016)	SCZ (17)	Auditory pre-recorded speech distortion	Internal vs external (own speech, other speech, unsure)	Correlations ^a	AH severity ^e × accuracy (self-speech + distorted/ undistorted + negative valence): NS

(Continued)

Table 2. (Continued)

Author (year)	Groups (n)	Task type and sensory mode	Stimuli source options	Analyses	Results
Seal et al. (1997)	SCZ-LAH (10) SCZ-CAH (6) SCZ-NLAH (11)	Auditory source monitoring	Internal vs external (heard, said, new)	Group contrasts	Accuracy & externalising misattributions: SCZ-CAH vs SCZ-LAH vs SCZ-NLAH (high/low cognitive & affective load): all NS
Smith and Gudjonsson (1995)	SSD-LAH (17) SSD-NLAH (15)	Auditory source monitoring	Internal vs external (heard, said, new)	Group contrasts	Accuracy & misattributions: SSD-LAH vs SSD-NLAH: both NS
Stephane et al. (2010a)	SCZ/SAD-LAH (26) SCZ/SAD-NLAH (6)	Auditory-visual source monitoring	Internal form (read aloud, read silently, new)	Group contrasts	Correct responses: SCZ/SAD-LAH vs SCZ/SAD-NLAH: NS Misattribution bias: (data not extracted) ^e
Stephane et al. (2010b)	SCZ/SAD-LAH (31) SCZ/SAD-NLAH (8)	Auditory-visual source monitoring	Internal vs external (heard, read, new)	Group contrasts	Misattribution bias: SCZ/SAD-LAH vs SCZ/SAD-NLAH: NS
Versmissen et al. (2007)	SCZ/SAD (41)	Auditory immediate speech feedback	Internal vs external (own speech, other speech, unsure)	Regression/other ^a	CAH severity did not contribute to number of misattributions: NS
Waters et al. (2006)	SCZ-CAH (19) SCZ.NCAH (24) SCZ-NLAH (4) ^f	Visual source monitoring & temporal context memory	Internal vs external (own/other action) Temporal context (run 1 or 2)	Group contrasts	% participants with source/temporal accuracy deficit: SCZ-CAH > SCZ-NCAH (+1 standard deviation from healthy control mean): $p < 0.05$, $d = 0.64$ SCZ-CAH vs SCZ-NCAH (+2 standard deviation from healthy control mean): NS
Woodward et al. (2007)	SCZ/SCAD-CAH (16) SCZ/SAD-NCAH (35)	Auditory-visual source monitoring	Internal vs external (heard, read, new) External form (experimenter, computer)	Group contrasts	Externalising misattributions: SCZ/SAD-CAH > SCZ/SAD-NCAH: $p < 0.05$, $d = 0.70$ Discrimination, internalising misattribution & external form misattributions: SCZ/SAD-CAH vs SCZ/SAD-NCAH: all NS

Diagnoses: SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizophrenia-spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Note: Task conditions are listed in parentheses following relevant variables. Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical AH vs healthy control only).

^bAH variable where higher scores indicate lower severity.

^cEffect size values, means and standard deviations not available. Unable to calculate Cohen's d .

^dEffect size unable to be converted to Cohen's d .

^eAdditional results involving phenomenological AH variables not extracted.

^fSCZ-NLAH did not show above-chance deficits of 1–2 standard deviation(s) from healthy control mean, thus scores were not entered into group contrasts.

Table 3. General memory.

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Badcock et al. (2005)	SCZ-CAH (23) SCZ-NCAH (20)	Inhibition of currently irrelevant memories	Group contrasts	Overall hits & run I false alarms: SCZ-CAH vs SCZ-NCAH: both NS
Brébion et al. (2007a)	SCZ-AH (18) SCZ-NAH (23)	Word list free recall	Group contrasts Correlations	Total number of recalled words: SCZ-AH vs SCZ-NAH: NS AH severity × total number of recalled words: NS
Brébion et al. (2007b)	SCZ (24)	Picture recognition	Correlations	↑ AH severity × ↓ recognition accuracy (colour): $p < 0.001$, $d = 1.58$ ↑ AH severity × ↑ response bias ^a (overall): $p < 0.05$, $d = 1.01$ ↑ AH severity × recognition accuracy (black-white), response bias (black-white/colour): all NS
Brébion et al. (2008)	SCZ-AH (17) SCZ-NAH (24)	Word & picture recognition	Group contrasts Correlations	Recognition accuracy: SCZ-AH vs SCZ-NAH: NS ↑ AH severity × ↓ response bias ^a (pictures): $p < 0.001$, $d = 1.76$ AH severity × response bias (words): NS
Brébion et al. (2009)	SCZ-AH (12) SCZ-NAH (29)	Word list free recall	Correlations	AH severity × correctly recalled words: NS
Brébion et al. (2011)	SCZ-AH (12) SCZ-NAH (29)	Word list free recall	Group contrasts Correlations	Serial & semantic encoding: SCZ-AH vs SCZ-NAH: both NS AH severity × semantic encoding: NS AH severity × serial encoding: NS
Brébion et al. (2015)	SCZ (50)	WAIS-III letter number span (LNS) Picture recognition	Correlations Regression/other	↑ AH severity × ↓ LNS working memory: $p < 0.005$, $d = 0.87$ Increased AH severity contributed to reduced picture recognition (colour + immediate): $p < 0.01$, $d = 1.01$, (high-complexity + colour + immediate): $p < 0.025$, $d = 0.77$ AH severity did not contribute to picture recognition (black-white + immediate/ delayed): NS
Brébion et al. (2016)	SCZ (57)	Word list recognition	Regression/other ^b	Increased AH severity contributed to increased response bias ^a (male participants + high-frequency words): $p < 0.025$, $d = 0.90$, (male participants + semantically organisable words): $p < 0.05$, $d = 0.98$ AH severity did not contribute to response bias (male participants + low-frequency words) (female participants + high/low frequency words) (female participants + semantically organisable words): all NS
Bruder et al. (2011)	SCZISAD-CAH (10) SCZISAD-NCAH (18)	Word serial position test	Group contrasts Correlations	Working memory, position 2: SCZISAD-CAH < SCZISAD-NCAH: $p < 0.05$, $d = 0.54$ Working memory, position 3: SCZISAD-CAH < SCZISAD-NCAH: $p < 0.05$, $d = 0.74$ ↑ CAH severity × ↓ overall working memory (in participants with 'normal' tone discrimination abilities): $p = 0.01$, $d = 0.75$
Brunelin et al. (2006)	SCZ-CAH (30) SCZ-NCAH (31)	Source monitoring task	Group contrasts	Distractor detection rate: SCZ-CAH vs SCZ-NCAH: NS

(Continued)

Table 3. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Franck et al. (2000)	SCZ-CAH (7) SCZ-PAH (5)	Source monitoring task	Group contrasts	Recognition errors: SCZ-CAH > SCZ-PAH: $p = 0.03$, $d = 1.45$
Gaudiot et al. (2019)	SCZ (31)	Free association task with familiarity rating BACS digit sequencing test (DST)	Regression/other ^b	Mediation analysis: Familiarity rating (pseudo-words) mediated the relationship between DST working memory and AH severity: $p < 0.05^c$
Gisselgård et al. (2014)	SSD (52)	WAIS-III digit span forward (DS-F) & backward (DS-B), letter-number sequencing (LNS) California verbal learning test (CVLT-II)	Correlations Regression/other	↑ CAH severity × ↓ DS-F auditory memory span: $p < 0.05$, $d = 0.58$ ↑ CAH severity × ↓ DS-B verbal working memory: $p < 0.01$, $d = 0.80$ ↑ CAH severity × ↓ LNS verbal working memory: $p < 0.005$, $d = 1.07$ CAH severity × CVLT-II verbal learning & memory: NS Reduced working memory score (mean z-score for DS-B & LNS) contributed to increased CAH severity: $p < 0.001$, $\beta = 0.61^c$
Guillem et al. (2001)	SCZ (27)	WMS-R digit span forward (DS-F) & backward (DS-B), paired-associates (words) learning test (PALT)	Correlations	↑ AH severity × ↓ PALT score (difficult pairs + immediate recall): $p = 0.031$, $d = 0.90$ AH severity × PALT score (easy pairs + immediate recall) (difficult/easy pairs + delayed recall), DS-F auditory memory span, DS-B auditory working memory: all NS
Haddock et al. (1996)	Study 1: SCZ-CAH (16) SCZ-NCAH (11) Study 2: SCZ-CAH (12) SCZ-NCAH (12)	Phonological encoding (PE) task Unattended speech effect (USE) task	Group contrasts	Study 1. PE % correct recall: SCZ-CAH vs SCZ-NCAH: NS Study 2. USE total & % correct serial position (speech present/absent): SCZ-CAH vs SCZ-NCAH: both NS
Jenkins et al. (2018)	SSD-CAH (64) SSD-NCAH (12)	MCCB working memory, verbal memory & visual memory subscales	Group contrasts Regression/other	Working Memory: SSD-CAH < SSD-NCAH: $p = 0.006$, $d = 0.90$ Verbal & visual memory: SSD-CAH vs SSD-NCAH: both NS Working memory contributed to CAH presence: $p = 0.009^d$, but verbal and visual memory did not: NS
Kayser et al. (2012)	SSD-CAH (26) SSD-NCAH (49)	Recognition memory paradigm Working memory paradigm	Group contrasts	Recognition sensitivity (words, faces): SSD-CAH vs SSD-NCAH: NS Working memory % correct (faces): SSD-CAH vs SSD-NCAH: NS Working memory % correct (words): SSD-CAH < SSD-NCAH: $p \leq 0.05^e$
Lee et al. (2004)	SCZ-CAH (24) SCZ-NCAH (25)	Sentence repetition test	Group contrasts	Longest sentence correct: SCZ-CAH vs SCZ-NCAH: NS
Lee et al. (2013)	SCZ (26)	Reality evaluation task	Correlations ^b	AH severity × recognition discriminability & response bias: both NS
McKay et al. (1981)	LAH (10) NLAH (20)	LNNB memory subscale	Group contrasts	LNNB memory: LAH vs NLAH: NS
Rocca et al. (2006)	SCZ (70)	California Verbal Learning Test (CVLT) The Corsi Test (CT)	Correlations	AH severity × CT spatial memory span, CVLT auditory recall & recognition: all NS

(Continued)

Table 3. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Siddi et al. (2016)	SSD-CAH (46) SSD-NCAH (44)	WAIS-III digit span forward & backward (DS-F & DS-B)	Group contrasts	DS-F auditory memory span & DS-B auditory working memory: SSD-CAH vs SSD-NCAH: NS
Siddi et al. (2017)	SSD-CAH (46) SSD-NCAH (44)	WAIS-III digit span forward & backward (DS-F & DS-B) Rey auditory verbal learning test (RAVLT)	Regression/other ^b	DS-F auditory memory span, DS-B auditory working memory, RAVLT short-term memory, verbal learning & delayed recall did not contribute to CAH presence: all NS
Smith and Gudjonsson (1995)	SSD-LAH (17) SSD-NLAH (15)	Short story free recall	Group contrasts	Delayed recall: SSD-LAH < SSD-NLAH: $p < 0.05$, $d = 0.79$ Immediate recall & false recall: SSD-LAH vs SSD-NLAH: both NS
Stephane et al. (2010b)	SCZ/SAD-LAH (31) SCZ/SAD-NLAH (8)	Source monitoring	Group contrasts	Recognition accuracy: SCZ/SAD-LAH vs SCZ/SAD-NLAH: NS
Thompson and Copolov (1998)	SSD-LAH (19) SSD-NLAH (13)	Sentence repetition test (SRT) Fable recall (FR) PALPA digit-span subtests	Group contrasts	SRT recall in grammatical context & FR recall of information units: SSD-LAH vs SSD-NLAH: all NS PALPA: digit span & matching digit span: SSD-LAH vs SSD-NLAH: both NS
Toh et al. (2020)	SSD-CAH (46) SSD-PAH (37) SSD-NLAH (40)	MCCB working memory subscale	Group contrasts	Working memory: SSD-CAH vs SSD-PAH vs SSD-NLAH: NS
Waters et al. (2003)	SCZ (42)	WAIS-III digit span forward (DS-F)	Correlations ^b	AH severity \times DS-F auditory memory span: NS
Woodward et al. (2007)	SCZ/SCAD-CAH (16) SCZ/SAD-NCAH (35)	Source monitoring	Group contrasts	Item recognition: SCZ/SAD-CAH vs SCZ/SAD-NCAH: NS

Diagnoses: SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizophrenia-spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Task abbreviations: BACS: Brief assessment of cognition in Schizophrenia, LNNB: Luria-Nebraska Neuropsychological Battery, MCCB: Matrics Consensus Cognitive Battery, PALPA: Psycholinguistic Assessment of Language Processing, WAIS-III: Wechsler Adult Intelligence Scales, third edition, WMS-R: Wechsler Memory Scales, Revised.

Note: Task conditions are listed in parentheses following relevant variables. Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aLower scores indicate a conservative pattern of responding, while higher scores indicate a more liberal pattern of responding.

^bAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical AH vs healthy control only).

^cEffect size unable to be converted to Cohen's d .

^dEffect size values, means and standard deviations not available. Unable to calculate Cohen's d .

^eAdditional results involving phenomenological AH variables not extracted.

Table 4. Language processing, production and hemispheric organisation.

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Docherty (2012)	SCZ/SAD (110)	Speech samples analysed	Correlations ^a	↑ CAH severity × ↑ frequency of missing referents: $p < 0.001$, $d = 0.77$
Gisselgård et al. (2014)	SSD (52)	Verbal letter fluency	Correlations	CAH severity × phonetic fluency: NS
Green et al. (1994)	SCZ/SAD-CAH (24) SCZ/SAD-NCAH (21)	Dichotic listening test	Group contrasts	Mean right ear advantage (REA): SCZ-CAH < SCZ-NCAH (non-forced attention): $p < 0.04$, $d = 0.66$ SCZ-CAH < SCZ-NCAH (forced-left attention): $p < 0.04$, $d = 0.65$ SCZ-CAH < SCZ-NCAH (forced-right attention): $p < 0.02$, $d = 0.76$
Hoffman et al. (1999)	SSD-CAH (28) SSD-NCAH (26)	Benton–Hamsher sentence repetition test	Group contrasts	Longest sentence accurately recited (serial linguistic expectation): SSD-CAH < SSD-NCAH: $p < 0.05$, $d = 1.05^b$
Levitan et al. (1999)	SCZ-CAH (11) SCZ-PAH (19)	Dichotic listening test	Group contrasts Correlations	Laterality index ^c : SCZ-CAH < SCZ-PAH: $p < 0.02$, $d = 0.91$ CAH severity × laterality index: NS
Løberg et al. (2015)	SSD-FAH (33) SSD-NFAH (17)	Dichotic listening test	Group contrasts	Laterality index: SSD-FAH T2 ^d vs SSD-NFAH T2 (left/right/non-forced attention): NS SSD-FAH T1 vs SSD-FAH T2 ^d (left/right/non-forced attention): NS SSD-NFAH T1 vs SSD-NFAH T2 (left/right/non-forced attention): NS Left ear correct responses: SSD-FAH T2 vs SSD-NFAH T2 (left/right/non-forced attention): NS Right ear correct responses: SSD-FAH T2 vs SSD-NFAH T2 (left/right/non-forced attention): NS Total correct responses: SSD-FAH T2 vs SSD-NFAH T2 (forced-right/non-forced attention): NS SSD-FAH T2 < SSD-NFAH T2 (forced-left attention): $p = 0.04$, $d = 0.64$
McKay et al. (2000)	SCZ/SAD-CAH (22) SCZ/SAD-NLAH (16)	Staggered spondaic word test (SSWT) Dichotic consonant–vowels test (DCVT)	Group contrasts	SSWT % errors (non-competing/competing mode + left/ right/left-right ears) (response bias + ear/order/reversal effects) & DCVT score (simultaneous/lagging presentation + left/ right/ left-right ears): SCZ/SAD-CAH vs SCZ/SAD-NLAH: all NS
McKay et al. (1981)	LAH (10) NLAH (20)	LNNB expressive and receptive language subscales	Group contrasts	Expressive language ^e : LAH > NLAH: $p < 0.01$, $d = 1.01^c$ Receptive language: LAH vs NLAH: NS
Rossell and Boundy (2005)	SCZ/SAD-CAH (20) SCZ/SAD-NLAH (20)	Dichotic listening task (DLT) Semantic fluency task (SFT)	Group contrasts	DLT number of correct responses: SCZ/SAD-CAH < SCZ/SAD-NLAH (right ear): $p < 0.05$, $d = 0.39^b$ SCZ/SAD-CAH vs SCZ/SAD-NLAH (left/both ears): both NS DLT total % correct: SCZ/SAD-CAH vs SCZ/SAD-NLAH (overall): NS SFT % correct: SCZ/SAD-CAH vs SCZ/SAD-NLAH: NS
Rossell et al. (2001)	SCZ-LAH (42) SCZ-NLAH (29)	Dichotic listening test (DLT) Finger tapping test (FTT)	Group contrasts	DLT laterality index ^c : SCZ-LAH vs SCZ-NLAH: NS FTT laterality index: SCZ-LAH vs SCZ-NLAH: NS

(Continued)

Table 4. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Siddi et al. (2016)	SSD-CAH (46) SSD-NCAH (44)	Semantic fluency test (SFT) Papagno Test (PT) metaphor & idiom processing subtests	Group contrasts Correlations Regression/other	PT metaphor processing: SSD-CAH < SSD-NCAH: $p < 0.05$, $d = 0.48^b$ PT idiom processing & SFT score: SSD-CAH vs SSD-NCAH: both NS ↑ CAH severity \times ↓ PT metaphor processing: $p < 0.05$, $d = 0.82$ CAH severity \times PT idiom processing: NS Decreased PT metaphor processing contributed to increased CAH severity: $p < 0.0001$, $d = 0.87$, but idiom processing did not: NS
Siddi et al. (2017)	SSD-CAH (46) SSD-NCAH (44)	Semantic & phonetic fluency task	Regression/other ^a	Reduced semantic fluency contributed to increased CAH presence: $p = 0.036$, $d = 0.49$, but phonetic fluency did not: NS
Thompson and Copolov (1998)	SSD-LAH (19) SSD-NLAH (13)	Speech sample analysis Semantic & phonetic fluency task PALPA subtests: minimal pairs (MP), lexical decision (LD), picture matching (PM), confrontation naming (CN), sentence-picture matching (SPM), reading- sentence-picture matching (RSPM)	Group contrasts	Monologue cohesion score: SSD-LAH > SSD-NLAH: $p < 0.05$, $d = 0.89^b$ Monologue word count, cartoon description, cohesion score, word count, semantic & phonetic fluency scores: SSD-LAH vs SSD-NLAH: all NS PALPA: MP phonological distinctions, LD real word distinctions, PM semantic knowledge, CN object labelling, SPM auditory comprehension, RSPM written comprehension: SSD-LAH vs SSD-NLAH: all NS
Titone and Levy (2004)	SCZISAD (18)	Lexical competition paradigm	Correlations ^a	↑ LAH severity \times ↑ misidentification bias (high-frequency words): $p < 0.05$, $d = 1.15$ LAH severity \times word identification accuracy: NS

Diagnoses: SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizophrenia spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). There was one study for which these definitions did not apply, with groups described as 'frequent' (FAH) and 'non-frequent' (NFAH) instead (Løberg et al., 2015). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Task abbreviations: LNNB: Luria-Nebraska Neuropsychological Battery, PALPA = Psycholinguistic Assessment of Language Processing.

Note: Task conditions are listed in parentheses following relevant variables. Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical AH vs healthy control only).

^bEffect size values not available. Cohen's d calculated from means and standard deviations.

^cLaterality index: positive scores indicate a right ear advantage (REA), while negative scores indicate a left ear advantage (LEA).

^dParticipants examined at baseline (T1) and 3 months post admission (T2).

^eCognitive variable where higher scores indicate poorer expressive language performance.

Table 5. Executive function.

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Badcock et al. (2005)	SCZ-CAH (23) SCZ-NCAH (20)	Inhibition of currently irrelevant memories	Group contrasts	Suppression, false alarms run 2-4: SCZ-CAH > SCZ-NCAH: $p = 0.027$, $d = 0.71$
Bhatia et al. (2009)	SCZ/SAD (172)	Trail making task (TMT) part A & B	Regression/ other ^a	Presence of LAHs contributed to slower TMT-A visual attention: $p = 0.01$, $d = 0.49$, and reduced TMT-B set-shifting abilities: $p = 0.01$, $d = 0.48$, but not to TMT B-A executive control: NS
Carter et al. (1996)	SCZ (23)	Global-local letter detection task	Correlations ^a	↑ AH severity × ↓ local target processing, visual attention: $p = 0.02$, $d = 1.42$
Chu et al. (2021)	SSD (35)	Learned irrelevance task	Correlations ^a	AH severity × learning score, cognitive flexibility: NS
Gisselgård et al. (2014)	SSD (52)	WAIS-III vocabulary, digit symbol coding (SC), block design (BD) & trail making task (TMT) A & B	Correlations	CAH severity × vocabulary, SC perceptual-motor speed, BD visuospatial abilities, TMT-A visual attention & TMT-B set-shifting: all NS
Guillem et al. (2001)	SCZ (27)	Wisconsin card sort test (WCST) Continuous performance test (CPT) Cancellation test of Zazzo (CTZ) Wickens' test for proactive interference and release (WPIR)	Correlations	↑ AH severity × ↑ CTZ visual scanning speed: $p = 0.021$, $d = 0.98$ AH severity × CTZ spatial attention, WCST executive function, CPT sustained attention & WPIR interference: all NS
Guillem et al. (2008)	SCZ (96)	Wickens' test for proactive interference and release	Regression/ other	Increased AH severity contributed to increased sensitivity to interfering information: $p = 0.02$, $d = 0.63$
Heilbrun and Blum (1984)	CAH (22) NCAH (22)	Unusual meanings vocabulary test	Group contrasts	Vocabulary score: CAH < NCAH: $p < 0.01$, $d = 0.86$
Henquet et al. (2020)	SCZ (1261)	Beads task	Regression/ other ^a	LAH severity × jumping to conclusions did not contribute to psychosis risk: NS
Hoffman (1999)	SCZ-CAH (22) SCZ-NCAH (25)	Auditory continuous performance test	Group contrasts	Omission & commission errors, auditory attention: SCZ-CAH > SCZ-NCAH ^{b,c}
Hoffman et al. (1999)	SSD-CAH (24) SSD-NCAH (21)	Auditory continuous performance test	Group contrasts	Hits & false alarms, auditory attention: SSD-CAH vs SSD-NCAH: both NS

(Continued)

Table 5. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Jenkins et al. (2018)	SSD-CAH (64) SSD-NCAH (12)	MCCB processing speed, attention & reasoning subscales	Group contrasts Regression/ other	Processing speed: SSD-CAH < SSD-NCAH: $p = 0.026$, $d = 0.71$ Attention and reasoning: SSD-CAH vs SSD-NCAH: both NS Processing speed, attention & reasoning did not contribute to CAH presence: all NS
Kråkvik et al. (2013)	SSD-CAH (31)	Connors' visual continuous performance test	Correlations	↑ CAH frequency ^d × ↑ detectability, visual attention ^e : $p < 0.05$, $d = 1.06$
Laloyaux et al. (2018)	SCZ (15)	Visual source flexibility task (attention)	Correlations ^a	CAH frequency × reaction time (stimuli to imagery switch, imagery to stimuli switch) & % errors (stimuli to imagery switch, imagery to stimuli switch): all NS
Løberg et al. (2015)	SSD-FAH (33) SSD-NFAH (17)	Dichotic listening test (attention paradigm)	Group contrasts	Laterality index: Forced-left vs forced-right vs non-forced (SSD-FAH): NS Forced-left < non-forced < forced-right (SSD-NFAH): $p < 0.03$ ^b
McKay et al. (1981)	LAH (10) NLAH (20)	LNNB subscales	Group contrasts	LNNB: motor speed/control, visual processing, motor writing/spelling, reading accuracy & arithmetic, intelligence subscales: LAH vs NLAH: all NS
Rocca et al. (2006)	SCZ (70)	Stroop test Trail making task (TMT) part A & B Wisconsin card sorting test (WCST) WAIS-R digit symbol coding (SC)	Correlations	AH severity ^c × Stroop inhibition, TMT-A visual attention, TMT-B set-shifting, WCST executive function, SC perceptual-motor speed: all NS
Siddi et al. (2016)	SSD-CAH (46) SSD-NCAH (44)	WAIS-III digit symbol substitution (DSS)	Group contrasts	DSS processing speed: SSD-CAH vs SSD-NCAH: NS
Siddi et al. (2017)	SSD-CAH (46) SSD-NCAH (44)	WAIS-III trail making test (TMT) part A & B, digit symbol substitution (DSS) Stroop test FAB motor programming, sensitivity to interference & Go-No-Go task (GNG)	Regression/ other ^a	Reduced TMT-B set-shifting contributed to CAH presence: $p = 0.040$, $d = 0.12$ Stroop cognitive flexibility, TMT-A visual attention, DSS processing speed, FAB motor programming, sensitivity to interference & GNG inhibitory control did not contribute to current CAH presence: all NS
Thompson and Copolov (1998)	SSD-LAH (19) SSD-NLAH (13)	National adult reading test (NART) Matrices of raven (MR) & Categorisation task (CT)	Group contrasts	NART premorbid intelligence, MR/CT abstract reasoning: SSD-LAH vs SSD-NLAH: all NS
Toh et al. (2020)	SSD-CAH (46) SSD-PAH (37) SSD-NLAH (40)	MCCB processing speed, attention/vigilance, verbal learning, visual learning, & reasoning/ problem solving Stroop test	Group contrasts Regression/other	Visual learning: SSD-CAH < SSD-NLAH: $p < 0.001$, $d = 1.24$ SSD-PAH < SSD-NLAH: $p < 0.001$, $d = 1.24$ SSD-CAH vs SSD-PAH: NS Processing speed, attention/vigilance, verbal learning, reasoning/problem solving & Stroop inhibition: SSD-CAH vs SSD-PAH vs SSD-NLAH: all NS Reduced visual learning ability contributed to LAH presence: $p = 0.007$, $d = 0.26$; Stroop inhibition did not contribute to CAH presence: NS

(Continued)

Table 5. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Waters et al. (2003)	SCZ (42)	Hayling sentence completion test (HSCT) Inhibition of currently irrelevant memories (ICIM)	Correlations ^a	<p>↑ AH severity × ↑ HSCT errors (high salience): $p < 0.01$, $d = 0.41$</p> <p>↑ AH severity × ↑ ICIM, false alarms inhibition: $p < 0.01$, $d = 0.93$</p> <p>AH severity × HSCT errors (low salience) & ICIM false alarms (new learning): both NS</p>
Waters et al. (2006)	SCZ-CAH (19) SCZ.NCAH (24) SCZ-NLAH (4) ^f	Hayling sentence completion test (HSCT) Inhibition of currently irrelevant memories (ICIM) Memory for context task (MCT) ^g	Group contrasts	<p>% participants with HSCT/ICIM inhibition deficit:</p> <p>SCZ-CAH > SCZ-NCAH (1 + standard deviation from HC mean): $p < 0.001$, $d = 1.45$</p> <p>SCZ-CAH > SCZ-NCAH (2 + standard deviation from HC mean): $p < 0.01$, $d = 1.07$</p> <p>% participants with HSCT/ICIM inhibition & MCT context memory deficit:</p> <p>SCZ-CAH > SCZ-NCAH (1 + standard deviation from HC mean): $p < 0.001$, $d = 0.62$</p> <p>SCZ-CAH > SCZ-NCAH (2 + standard deviation from HC mean): $p < 0.01$, $d = 0.82$</p>

Diagnoses: SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizophrenia-spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). There was one study for which these definitions did not apply, with groups described as 'frequent' (FAH) and 'non-frequent' (NFAH) instead (Løberg et al., 2015). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Task abbreviations: FAB: Frontal assessment battery, LNNB: Luria-Nebraska Neuropsychological Battery, MCCB: Matrices Consensus Cognitive Battery, WAIS-III & WAIS-R: Wechsler Adult Intelligence Scales, third edition & revised.

Note: Task conditions are listed in parentheses following relevant variables. Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical AH vs healthy control only).

^bEffect size values, means and standard deviations not available. Unable to calculate Cohen's d .

^cSignificance value/level not reported (identified as 'significant' by author).

^dAdditional results involving phenomenological AH variables not extracted.

^eCognitive variable where higher scores indicate poorer attentional performance.

^fSCZ-NLAH did not show above-chance deficits of 1–2 standard deviation(s) from healthy control mean, thus these scores were not entered into group contrasts.

^gPrimary data from the Memory for Context task were included in the source monitoring table. Task included here as relevant to data point examining the combined prevalence of inhibition and context memory deficits in the context of AHs (Waters et al., 2006).

Table 6. Emotion processing and social cognition.

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Bucci et al. (2008)	SCZ-CAH (28) SCZ-NCAH (32)	Gesture signal detection task	Group contrasts	Sensitivity & response bias: SCZ-CAH vs SCZ-NCAH: both NS
McLachlan et al. (2013)	SCZ/SAD-LAH (19) SCZ/SAD-NLAH (15)	Affective prosody task	Group contrasts	Sensitivity: SCZ/SAD-LAH vs SCZ/SAD-NLAH (happy/sad/scared/neutral): all NS
Rossell and Boundy (2005)	SCZ/SAD-CAH (20) SCZ/SAD-NLAH (20)	DANVA affective prosody & facial expression subtests Affective prosody & non-linguistic sound tasks	Group contrasts	DANVA, % correct: SCZ/SAD-CAH < SCZ/SAD-NLAH (prosody + overall): $p < 0.05$, $d = 0.46^a$ SCZ/SAD-CAH < SCZ/SAD-NLAH (prosody + sad): $p < 0.05$, $d = 0.70^a$ SCZ/SAD-CAH < SCZ/SAD-NLAH (prosody + fearful): $p < 0.05$, $d = 0.25^a$ SCZ/SAD-CAH vs SCZ/SAD-NLAH (prosody + happy/angry): both NS SCZ/SAD-CAH vs SCZ/SAD-NLAH (faces + all affects): NS DANVA % misidentified: SCZ/SAD-CAH > SCZ/SAD-NLAH (sad as happy, fearful as happy, fearful as angry): $p < 0.05^b$ Affective prosody, % correct: SCZ/SAD-CAH < SCZ/SAD-NLAH (angry): $p < 0.05$, $d = 0.53^a$ SCZ/SAD-CAH vs SCZ/SAD-NLAH (all other affects): all NS Affective prosody, intensity rating: SCZ/SAD-CAH vs SCZ/SAD-NLAH (all affects): all NS Non-linguistic sounds, % correct: SCZ/SAD-CAH < SCZ/SAD-NLAH (total): $p < 0.05$, $d = 0.50^a$ SCZ/SAD-CAH < SCZ/SAD-NLAH (happy): $p < 0.05$, $d = 0.32^a$ SCZ/SAD-CAH < SCZ/SAD-NLAH (fearful): $p < 0.05$, $d = 0.59^a$ SCZ/SAD-CAH vs SCZ/SAD-NLAH (sad/angry/neutral): all NS Non-linguistic sounds, % misidentified: SCZ/SAD-CAH > SCZ/SAD-NLAH (happy as fearful, fearful as sad): $p < 0.05^b$
Rossell et al. (2013)	SSD (108)	CATS affective prosody subtests	Correlations ^c	↑ CAH severity × ↓ % correct, affective tone identification: $p < 0.001$, $d = 0.65$ ↑ CAH severity × ↓ % correct, affective tone identification (conflicting tone-content): $p < 0.001$, $d = 0.70$ ↑ CAH severity × ↓ % correct, affective content identification (conflicting tone-content): $p < 0.001$, $d = 0.77$ CAH severity × % correct, affective tone discrimination: NS
Shah et al. (2018)	SCZ-CAH (23)	Emotional facial identification task	Correlations ^c	CAH severity × % accuracy, affect identification: NS

(Continued)

Table 6. (Continued)

Author (year)	Groups (n)	Cognitive task	Analyses	Results
Shea et al. (2007)	SCZ/SAD-CAH (38) SCZ/SAD-NCAH (29)	Prosodic processing task	Group contrasts	% Correct: SCZ/SAD-CAH < SCZ/ICZCAD-NCAH (overall): $p = 0.006$, $d = 0.63^a$ % Misidentified: SCZ/SAD-CAH < SCZ/ICZCAD-NCAH (sad as neutral): $p = 0.06$, $d = 0.64^a$ SCZ/SAD-CAH vs SCZ/ICZCAD-NCAH (happy/ neutral): NS
Toh et al. (2020)	SSD-CAH (46) SSD-PAH (37) SSD-NLAH (40)	MCCB social cognition subscale	Group contrasts Regression/other	Social cognition score: SSD-CAH vs SSD-PAH vs SSD-NLAH: NS Social cognition did not contribute to LAH presence: NS
Versmissen et al. (2008)	SCZ (40)	Hinting task	Regression/other	Reduced mentalising score contributed to CAHs: $p = 0.02$, $d = 0.93$
Zenisek et al. (2015)	BPAD-LAH (17) BPAD-NLAH (13)	Emotion recognition test	Group contrasts	Recognition accuracy (prosodic tone, facial expression): BPAD-LAH vs BPAD-NLAH: both NS

Diagnoses: BPAD: bipolar-affective disorder, SCZ: schizophrenia, SAD: schizo-affective disorder, SSD: schizo-affective disorder, SSD: schizophrenia-spectrum diagnosis. Where diagnostic information was not provided by authors, groups/variables were labelled according to AH status only.

Auditory hallucination status: CAH: current auditory hallucinations, LAH: lifetime auditory hallucinations (current status not measured), AH: auditory hallucinations present (timeframe not specified), NCAH: no current auditory hallucinations (lifetime history not measured), NLAH: no lifetime history of auditory hallucinations, PAH: past auditory hallucinations, NAH: no auditory hallucinations (timeframe not specified). Groups defined by general 'hallucination' status (i.e. sensory modality not specified) were not extracted. AH status was labelled according to the timeframes explicitly specified by authors in measurement descriptions. In the case of ambiguity, status inferences were not extrapolated from typical timeframes of administered measures, given that the implementation of assessment in its true form could not be confirmed.

Task abbreviations: CATS: Comprehensive Affective Testing System, DANVA: Diagnostic Analysis of Nonverbal Accuracy 2, MCCB: Matrix Consensus Cognitive Battery.

Note: Task conditions are listed in parentheses following relevant variables. Higher AH and cognition scores indicate increased severity and superior performance, respectively, unless otherwise specified. Significant findings ($p < 0.05$) are listed in bold font. NS: non-significant findings.

^aEffect size values not available. Cohen's d calculated from means and standard deviations.

^bEffect size values, means and standard deviations not available. Unable to calculate Cohen's d .

^cAdditional group contrasts omitted due to no AH-specific data available (i.e. clinical/AH vs healthy control only).

Table 7. Proportion of studies per cognitive domain within each quality range.

Cognitive domain	Number of studies	Low (%)	Moderate (%)	High (%)	Very high (%)
Overall	90	21	29	18	32
Perceptual decision-making and imagery	30	30	43	7	20
Source-monitoring and context memory	28	25	18	21	36
General memory	29	14	24	21	41
Language organisation, processing and production	14	14	7	29	50
Executive function	23	26	30.5	13	30.5
Emotion processing and social cognition	9	22	11	33.5	33.5

General memory. The general memory domain examined the retention of auditory and visual information, held across the short- and long-term, assessed via recall and recognition retrieval methods. Working memory paradigms were also included. In many instances, source-monitoring tasks involved a recognition component (e.g. identification of 'new' stimuli), and these results are reflected here. Key variables included the accuracy and error rate of recognised information, capacity of recall or working memory and composite memory scores reflecting overall performance on subtests (see Table 3). Significant results suggested an association between AH presence/severity and greater difficulties with: recall memory span ($d=0.58-0.90$), recognition capacity or bias ($d=1.01-1.45$), and working memory ($d=0.54-1.07$). In addition, AH severity was observed as a predictor of picture ($d=1.01-0.77$) and word list ($d=0.90-0.98$) recognition abilities, while working memory also predicted AH presence/severity ($\beta=0.61$).

Language processing, production and hemispheric organisation. This domain related to the processing and production of spoken language. The former is commonly examined via dichotic listening paradigms and considered within the context of a hemispheric organisation framework. A laterality index was often provided, reflecting the relative accuracy of detected input within the left and right ears. Language processing was also examined via broader measures of comprehension (e.g. higher order and integrative language processes). With respect the production of spoken language, key variables included verbal fluency and speech samples analysed for markers of coherence (see Table 4). Significant results suggested an association between AH presence/severity and a reduction in: left-hemispheric lateralisation for speech processing ($d=0.39-0.91$), verbal fluency and coherence ($d=0.77-1.01$), and

language comprehension capacity ($d=0.48-1.15$). Conversely, one study found increased verbal coherence in those with lifetime AHs, relative to those without, citing thought disorder within clinical control groups as a possible contributor (Thompson and Copolov, 1998). In addition, metaphor processing ability ($d=0.87$) and semantic fluency ($d=0.49$) were observed as predictors of AH severity and presence, respectively.

Executive function. More 'general' processes were grouped to form an executive function domain. This included processes such as inhibition, set-shifting, attention, processing speed and reasoning. These constructs were frequently measured as part of an overarching battery of neurocognitive tests, with each process represented by a particular subtest or a composite subscale score (see Table 5). Significant results suggested an association between AH presence/severity and reduced capacities across: inhibition ($d=0.71-1.45$) and attention ($d=1.06-1.42$). Additional isolated findings indicated associations between AH presence/severity and reduced: processing speed⁴ ($d=0.71-0.98$), vocabulary ($d=0.86$) and visual learning ($d=1.24$). AH presence was also observed as a predictor of visual attention ($d=0.49$) and set-shifting ($d=0.48$), while AH severity was seen to predict inhibition ($d=0.63$). Similarly, set-shifting ($d=0.12$) and visual learning ($d=0.26$) were observed as predictors of AH presence.

Emotion processing and social cognition. The emotion processing and social cognition domain related to the recognition of affect across multiple forms (e.g. facial expression, prosodic tone), and inference of others' state of mind or emotional perspective. Key variables included accuracy and sensitivity in the detection and differentiation of presented emotions, as well as composite scores for broad

social processing (see Table 6). Significant results suggested an association between AH presence/severity and greater difficulties in the accurate detection of emotional expression ($d=0.25-0.77$), with variation across specific emotions. In addition, ‘mentalising ability’ was observed as a predictor of AH presence ($d=0.93$).

Quality assessment

The proportion of quality ratings per cognitive domain are shown in TABLE7Table 7. For studies ranked the highest in quality, average checklist scores yielded similarly high rankings (4+) for all but the AH definition item (~3.5). For studies ranked the lowest in quality, average checklist scores were lowest for diagnostic status measurement, AH measurement and selection of controls (1–3). Regarding AH-specific language, 81% of studies demonstrated consistency across aims, methods and inferences, and these were relatively evenly spread across the domains. Of the studies which conducted group contrasts ($n=56$) and correlations ($n=35$), 36% and 69% of these reported at least 20 participants per group/analysis, respectively. A breakdown of studies reporting 20+ cases⁵ for group contrasts, per cognitive domain, is as follows: emotion processing and social cognition (67%), executive function (55%), language function (50%), perceptual decision-making and imagery (43%), general memory (32%) and source-monitoring/context memory (11%).

Discussion

Summary of findings

The current review offered a novel perspective by systematically summarising task-based measures of cognition across a broad range of domains, specifically relating to clinical AHs. Cognitive sub-domains with the highest frequency of reported significant findings suggested that AHs were associated with reduced sensitivity/accuracy for signal-noise discrimination and biases towards liberal responding for ambiguous stimuli, source-monitoring confusion for temporal order and self–other origin, reduced working memory capacity and reduced language coherency/fluency. Signal detection difficulties were noted for both auditory and visual stimuli and yielded medium-to-large effects. Challenges in distinguishing and remembering the temporal presentation order and self–other source of stimuli, along with reduced working memory capacity, yielded medium-to-large effects and were generally consistent with previous meta-analyses (Brookwell et al., 2013; Damiani et al., 2022; Waters et al., 2012b). In instances where the direction of self–other source attribution was distinguished, AHs were associated with a tendency to misattribute self-generated stimuli (e.g. thoughts, speech, actions) to an external source, but not vice versa.

Finally, medium-to-large effects were observed for reduced speech coherence, and to a lesser extent, verbal fluency. This was generally in line with prior integrative reviews, except for some sub-domains outlined below (e.g. Seal et al., 2004; Upthegrove et al., 2016; Waters et al., 2012a).

Primarily mixed findings, due to inconsistent results across outcome variables, were found for the monitoring of spatial context, language comprehension and hemispheric organisation. Notably, language comprehension demonstrated limited cohesion as a cognitive construct (e.g. metaphor processing, lexical competition, serial linguistic expectation; Hoffman et al., 1999; Siddi et al., 2016; Titone and Levy, 2004), while spatial context monitoring comprised only two studies. For hemispheric organisation of language function, variables extracted from dichotic listening tasks were interpreted inconsistently, thus undermining the strength of inferences made in some cases. While our findings offer less robust support for language lateralisation than previous reviews (Ocklenburg et al., 2013), our approach to screening AH measurement led to a narrowed selection of eligible data, likely accounting for discrepancies.

The sub-domains with predominantly non-significant associations to AHs were stimulus sensory feature discrimination, imagery vividness or illusion susceptibility, source-monitoring for internal events and external stimuli form, recognition and recall memory abilities, executive function, and processing of emotion and social cues. These results are consistent with previous accounts cautioning the relevance of mental imagery aberrations to AHs (Seal et al., 2004). While the past literature has suggested a possible role for sensory feature discrimination (McKay et al., 2000; McLachlan et al., 2013), and cognitive inhibition (Badcock and Hugdahl, 2014), these ideas were not well supported by the current empirical results. However, it is also noted that sub-components of these functions (e.g. automatic vs intentional and cognitive vs behavioural inhibition), which may yield distinct findings, were not examined here. Although broad summaries of executive function and AHs are yet to be provided, our observation of limited significant findings were mostly consistent with recent empirical conclusions (e.g. Toh et al., 2020). Given that a critical portion of literature was omitted from this review due to a lack of methodological specificity to AHs, the absence of substantial evidence noted in respect to the above cognitive sub-domains does not necessarily disconfirm the relevance of these processes, but rather highlights the need for further methodological rigour in validating such inferences. That said, it is important that the proportion of non-significant results, spanning many of the cognitive sub-domains (Figure 2), are not overlooked. Similarly, the issue of publication bias should be acknowledged, as non-significant findings may often not be retained/accepted for publication.

Quality assessment of studies indicated a breadth of rankings, with no clear differences across the cognitive domains. Studies of lower quality ranking exhibited the

most difficulty in measuring diagnostic status and AH experiences, along with employing appropriate selection methods for control groups. Studies ranked the highest in quality still demonstrated difficulty in AH definition, relative to the other methodological considerations. Notably, just over a third of studies employing group contrasts reported a minimum of 20+ cases per group,⁶ with difficulties in reaching this threshold most frequently observed in the source-monitoring literature.

Key take-home messages

Distinguishing AHs from diagnostic categories. A primary methodological concern was the lack of distinction between AHs and broader psychiatric diagnoses. Investigating cognitive profiles linked to clinical AHs with reference to a clinical (i.e. schizophrenia-spectrum diagnosis with no AH history) vs non-clinical (i.e. no major psychiatric history) control groups could reasonably be expected to yield distinct findings. Yet limited consideration for this issue has been offered in empirical studies. We chose to include only data which contrasted the presence and absence of AHs in the context of a psychiatric diagnosis. This was done with view that non-clinical control comparisons could not adequately distinguish AHs from broader diagnostic profiles. As a result, group contrast data for 25 retained studies were wholly excluded. Although these studies identified AHs as a variable of interest, appropriate methodological adjustments were not afforded. Subsequently, in some cases, differences can be seen between the original interpretations offered by authors, who drew on diagnostic-based contrasts to substantiate findings, and our interpretation of available secondary AH-specific data (e.g. follow-up symptom-based correlations). Furthermore, effectively isolating AHs from broader diagnoses is made increasingly difficult given the high level of comorbidity observed between schizophrenia and other conditions, such as depression or post-traumatic stress (Buckley et al., 2009). Thus, even if the presence/absence of AHs is examined exclusively within clinical samples, a range of different symptom profiles (e.g. mood state, trauma history, delusional ideation) may also interact with cognition. Yet, feasibly accounting for a large set of psychological factors presents quite a challenge. Our quality assessment found that approximately half of the included studies attempted to control for extraneous factor(s) as part of their analyses, while slightly over a third accounted for this by either matching, or examining the presence of statistical differences in, participant groups for key clinical variable(s). Considering the application of symptom dimensions on a broader scale, the relevance of transdiagnostic and continuum-based research is noted (e.g. De Leede-Smith and Barkus, 2013; Waters et al., 2012a). Notably, a transdiagnostic overview of cognitive research was not possible here, due

to a lack available studies examining AHs beyond the schizophrenia-spectrum; thus, demonstrating a notable gap within the literature.

Distinguishing AHs from related experiences. The measures employed to confirm or quantify the experience of AHs were vast, and in some instances, utilised in ways that failed to isolate AHs from broader psychosis symptomatology or cross-modality hallucination experiences. For example, gold-standard assessments, such as the Positive and Negative Syndrome Scale (PANSS; Kay et al., 1987) and Scale for the Assessment of Positive Symptoms (SAPS; Andreasen, 1984) were utilised frequently. While these measures offer good validity when employed in their intended form, they do not quantify the experience of hallucinations of an auditory nature specifically, unless relevant items are selected, or scoring is adapted accordingly. While many excluded studies provided aims and inferences in line with their employed methodology, a proportion offered interpretations in relation to AHs, despite a lack of methodological alignment. Notably, several of these studies represent seminal investigations which have been readily adopted by the field and informed theoretical models. A key driver may be the tendency to interchange the terms ‘hallucinations’ and ‘auditory hallucinations’ in academic language. Historically, AHs have received most attention, and as our recognition of hallucinations across the range of senses evolves (McCarthy-Jones et al., 2017), appropriate use of terminology and ensuing assessment measures, has lagged (Toh et al., 2019). While many studies exhibiting these discrepancies were screened out, ~20% of retained studies displayed inconsistencies in AH-specific terminology employed across aims, methodology and inferences.

Distinguishing variations in AH status. Many studies explored current, as opposed to lifetime or past AH experiences. Importantly, the presence of current/past AHs were often reported without consideration for, or measurement of, alternate timeframes. This highlights the potential for group overlap, and consequently difficulties in effectively isolating state from trait profiles. Similarly, there was much between-study variation in the definition of timeframes. For example, current AHs ranged from endorsement during testing, to confirmation over preceding weeks or even months. Similarly, while lifetime presence of AHs was sometimes guided by criteria, such as a specified number of episodes, this was not always the case. Furthermore, a notable portion of studies did not explicitly specify the time-period of interest when examining AHs. Longitudinal research has suggested that some cognitive profiles linked to schizophrenia may endure over time, while others demonstrate fluctuations (Heilbronner et al., 2016; Szöke et al., 2008). Yet, only six included studies directly compared variations in AH status, with source-monitoring and language lateralisation tentatively suggested as state markers.

Theoretical implications

Much debate remains with respect to the theoretical conceptualisation of AHs as a form of intrusive memory (Waters et al., 2006), inner speech (Jones and Fernyhough, 2007), or environmental cues attributed heightened salience/attention (Dodgson and Gordon, 2009; Kapur, 2003). However, given that no single cognitive mechanism solely informs each theoretical model, a multidimensional perspective is required. Hence, we found it useful to consider the current results in the context of broader cognitive frameworks, as discussed below.

Of the cognitive sub-domains identified within this review, those most substantially associated with AHs included reduced sensitivity/accuracy for signal-noise discrimination, and increased expectation biases for sensory input under conditions of ambiguity. These trends can be understood within a predictive coding framework, which emphasises the aberrant weighting of prior beliefs against incoming stimuli (Corlett et al., 2019; Sterzer et al., 2018). Similarly, excitatory–inhibitory models suggest that imbalances in the activation of bottom–up vs top–down neural networks may cause these two streams to reverberate and become misinterpreted as one another (i.e. circular inference), thus rendering an individual susceptible to anomalous perceptions (Jardri and Denève, 2013). Yet in the current review, task-based measures of candidate top–down functions (e.g. attentional control, inhibition, strong imagery) exhibited limited associations to AHs. However, much of the literature pertaining to inhibitory cognitive control, as measured by the dichotic listening task, was excluded from our synthesis due to non-AH-specific methodology.

Also associated with AHs in the current review were difficulties in distinguishing the self–other source and temporal presentation order of given events, along with reduced working memory capacity. These trends sit coherently within the self-monitoring framework, which proposes that individuals with AHs may exhibit greater difficulties in identifying with self-generated information (Waters et al., 2012b). Several functions have been hypothesised to underpin self-monitoring difficulties, including the ability to register the sensorimotor consequences of one’s actions at a neurological level (Ford and Mathalon, 2005), and the capacity to bind contextual cues with events in memory (Waters et al., 2004). It is possible that difficulties in the accurate encoding of relevant temporal cues may render the later recollection of events in memory as fragmented, and thus intrusive and foreign to oneself (Hardy, 2017; Steel, 2015). This notion applies to current cognitive perspectives of AHs in the context of trauma (McCarthy-Jones and Longden, 2016). It is worth acknowledging theoretical overlap, in that ‘low-level’ motor/sensory processing and ‘higher order’ expectations, posited to underpin the above frameworks, likely occur in conjunction. Consequently, due to its

integrative perspective, the predictive coding account has recently been employed as a means of reconciling these concepts (Griffin and Fletcher, 2017; Thakkar et al., 2021).

Clinical implications and developments

Psychological interventions for distressing AHs commonly focus on shifting individuals’ perceptions of, and relationships with, their voices; in addition to enhancing adaptive responses and coping strategies. Yet, there is a need to identify key processes of therapeutic change, rather than continuing to compare the efficacy of broad therapeutic frameworks (Thomas et al., 2014). Cognitive remediation programmes, which aim to improve functions such as memory, attention and verbal learning in individuals with schizophrenia-spectrum diagnoses (Wykes et al., 2011), have limited application to AHs, with mixed findings to date (Fiszdon et al., 2005; Thomas et al., 2018). Thus, ongoing cognitive research which considers the above pursuits is warranted; and several distinctions could be made to support clinical relevance. Namely, some processes have been hypothesised as implicated in the emergence of AHs (e.g. self-monitoring and context memory), while others may be more relevant to the phenomenology of such experiences (e.g. affective processing, perceptual expectation bias) (Waters et al., 2012a). Despite not being feasible for synthesis here, six out of the 101 studies meeting our initial inclusion criteria reported investigations between phenomenological characteristics of AHs and cognitive function. Thus, further examination of AH manifestations (e.g. valence, distress, controllability), beyond presence, could be of value. Similarly, differentiating between developmental and maintenance factors, as demonstrated within delusion literature (e.g. Freeman, 2016), may support the advancement of cognitive interventions relevant to the prevention, and alleviation, of distressing AHs, respectively. Future studies considering AH onset and chronicity would be well positioned to build on these ideas.

Given this review was restricted to AH presence/severity, it is acknowledged that cognitive processes associated with common therapeutic targets – such as levels of associated distress – have not been captured here. However, possible mechanisms of therapeutic change, aligning to the results of this review, could include targeting the accurate detection and anticipation of sensory input, identification with self-generated events, and integration of contextual memory cues. For example, imagery rescripting protocols, which aim to re-contextualise voice-related memories by meeting client needs and providing a corrective emotional experience, are hypothesised to target source-memory and context integration (Paulik et al., 2019). Evaluation of objective cognitive change markers, alongside the tracking of AH presentation and/or distress, pre- and post-intervention, could facilitate more direct links between cognitive mechanisms and key therapeutic outcomes.

Limitations and scope

There are several limitations within the current systematic review. Given the number of studies identified ($n=22,391$), stage one screening was completed by a single author, posing the possibility of missed records. Furthermore, the range of summarised cognitive mechanisms have, in many instances, been assessed with a breadth of novel behavioural paradigms. A key example is the operationalisation of internal–external source-monitoring, represented by memory for self (read/imagined) vs other (heard) words; differentiation of self vs other distorted speech; recognition of self-speech feedback delay; and recollection of learned words (other-presented) vs extra-list intrusions (self-originating errors). This rendered the definition of valid constructs, and development of broader cognitive themes, quite challenging, and overlap was inevitable. Given the scope of data, we chose to focus on methodological concerns regarding the measurement of AHs, cutting across studies, rather than providing a detailed critique of cognitive tasks. However, variation in task parameters can be seen in the results, and thus, some cognitive mechanisms may benefit from refinement. A range of stimulus presentation forms were also present. Conceptually, this assumes cognitive mechanisms to be modality-general in nature, and while we did not consider this in detail, further clarification is required to support such assumptions (Fernyhough, 2019). Importantly, the direction of relationships between cognition and AHs remains unconfirmed, and while difficulties are often inferred as mechanisms for AHs, they may also reflect by-products of such experiences. In addition, the status of antipsychotic medication use within study samples was not assessed during quality assessment or data extraction, due to inconsistent reporting of, and/or statistically controlling for, this information. However, antipsychotic medication use is recognised a possible confounding factor when examining cognitive performance (Haddad et al., 2023).

In contrasting the relative evidence supporting each cognitive mechanism, we focused on the proportion of significant results reported for each, as presented by the original authors. In some instances, a notable proportion of non-significant findings were also present (e.g. just under 50% internal–external source-monitoring studies), and thus, appropriate caution should be exercised when interpreting overall trends. It is also recognised that not all investigations can be attributed equal weighting, due to methodological concerns and sample size limitations. Nevertheless, this review offered a nuanced summary of a highly heterogeneous dataset, providing important insights into the methodological landscape of this field. Alternative approaches to synthesis (e.g. meta-analysis) could complement these findings in future. Finally, empirical evidence supporting theoretical AH models have emerged from various lines of investigation, extending beyond the

behavioural paradigms examined (e.g. electrophysiology, brain-imaging). Thus, while limited evidence was noted for some models, inspection of complimentary research streams may provide alternative conclusions.

Recommendations for future research

Recommendations for future research are focused on three areas: expanding, and matching for, the broader psychological context; measuring the timeframe of hallucinatory experiences; and specifying the sensory modality in which they emerge. Given that AHs emerge within a range of psychological conditions and the wider population (De Leede-Smith and Barkus, 2013; Sommer et al., 2010), the utility of transdiagnostic investigation is noted. With respect to expanding the psychological context of this research, cognitive investigations into the healthy AH continuum are progressing (e.g. Moseley et al., 2022), yet studies of this nature beyond schizophrenia-spectrum diagnoses are lacking. Phenomenological research examining hallucinations in the context of trauma are abundant and have begun to draw theoretical links with psychosis (McCarthy-Jones and Longden, 2016). However, at this stage, complementary task-based cognitive investigations warrant further attention, as do related conditions, such as borderline personality disorder. To this end, efforts should be made to match groups on psychological profiles where possible, with and without AHs, so that distinctions from related experiences can be made.

Explicitly defining and measuring the timeframe of AHs is also recommended, to allow for inferences regarding state vs trait cognitive markers. Importantly, links between proposed cognitive functions and current AHs do not, in isolation, confirm their role as a state marker. Rather, appropriate contrasts with past AHs are required, to identify the potential for cognitive change in conjunction with AH remission. Ideally, longitudinal investigation would be most revealing, although cross-sectional examination of AH status variations may provide a more feasible option. It is acknowledged that quantifying the status of complex phenomena, such as AHs, is difficult, given the need for arbitrary boundaries. Therefore, transparent reporting of AH measurement parameters is imperative in aiding cross-study comparisons. Going forward, the utilisation of measures which specifically examine hallucinations across each of the respective senses, rather than collapsing these diverse experiences into a single item, is advised. Subsequently, care must be taken when forming inferences regarding each sensory modality, so as to not form unfounded cross-modal conclusions. Appropriate measurement of these experiences will not only provide clarity within the AH literature, but also inform much needed investigations into the cognitive underpinnings of hallucinations beyond the auditory modality (Toh et al., 2019).

Conclusion

The current review highlighted the breadth and complexity of research examining cognition in the context of AHs. Difficulties in signal detection/expectation, self-other-source and temporal context monitoring, working memory, and language fluency, exhibited the most consistent links to AHs. Conceptually, these functions align with predictive coding and self-monitoring frameworks, and theoretical overlap is acknowledged. A lack of specificity regarding the sensory modality of hallucinatory experiences measured, as well as the common use of diagnostic-based comparisons, were of primary concern. Thus, we noted discrepancies between some theoretical AH models and corresponding empirical evidence. Finally, we recommended the extension of this research into broader transdiagnostic and healthy populations, in conjunction with relevant therapeutic intervention, to advance the field in a meaningful way.

Acknowledgements

The authors thank Associate Professor Vaughan Bell for his contributions to the protocol pre-registration and feedback on initial manuscript drafts. W.L.T. is supported by the Medical Research Council (NHMRC) New Investigator project grant [GNT1161609]; S.L.R. holds a Senior NHMRC Fellowship [GNT1154651]; and P.A. is supported by the UK Medical Research Council and University of Lille International Chair Award.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship and/or publication of this article.

ORCID iDs

Adrienne Bell  <https://orcid.org/0009-0007-8838-7463>

Wei Lin Toh  <https://orcid.org/0000-0002-4187-1182>

Susan L Rossell  <https://orcid.org/0000-0002-7415-8252>

Supplemental material

Supplemental material for this article is available online.

Notes

- Records were excluded if samples and analyses duplicated those reported in prior publications.
- Given its prevalence, working memory was considered separately in the ‘general memory’ table.
- Nine studies were not included in sample size/characteristic calculations due to sample duplication. In this case, studies with the largest auditory hallucination (AH)-specific samples were considered.
- While it is acknowledged that processing speed is not typically defined as an executive function, it is an important associated process and was therefore considered within this domain.
- While a cut-off of 20+ cases is ultimately an arbitrary representation of sample size adequacy, it is reflective of typical sample sizes within this body of literature and allowed for observation of sample size trends and limitations.
- A simple contrast (independent groups *t*-test) with two groups of 20 participants provides 80% power to detect effect sizes of $d=0.8$ or larger. It is reasonable to assume that associations between AH and cognition are likely to be smaller than this, so it is critical that future studies recruit larger sample sizes or justify their decision to detect only effect sizes of this magnitude.

References

- Alba-Ferrera L, De Erausquin GA, Hirnstein M, et al. (2013) Emotional prosody modulates attention in schizophrenia patients with hallucinations. *Frontiers in Human Neuroscience* 7: 59.
- Aleman A, Böcker KBE, Hijman R, et al. (2003) Cognitive basis of hallucinations in schizophrenia: Role of top-down information processing. *Schizophrenia Research* 64: 175–185.
- Allen PP, Amaro E, Fu CHY, et al. (2007) Neural correlates of the misattribution of speech in schizophrenia. *British Journal of Psychiatry* 190: 162–169.
- Allen PP, Johns LC, Fu CHY, et al. (2004) Misattribution of external speech in patients with hallucinations and delusions. *Schizophrenia Research* 69: 277–287.
- Andreasen NC (1984) *The Scale for the assessment of positive symptoms* (SAPS). Iowa City: The University of Iowa.
- Antonova I, van Swam C, Hubl D, et al. (2021) Altered visuospatial processing in schizophrenia: An event-related potential microstate analysis comparing patients with and without hallucinations with healthy controls. *Neuroscience* 479: 140–156.
- Badcock JC and Hugdahl K (2014) A synthesis of evidence on inhibitory control and auditory hallucinations based on the Research Domain Criteria (RDoC) framework. *Frontiers in Human Neuroscience* 8: 180.
- Badcock JC, Waters FAV, Maybery MT, et al. (2005) Auditory hallucinations: Failure to inhibit irrelevant memories. *Cognitive Neuropsychiatry* 10: 125–136.
- Bentall RP (1990) The illusion of reality: A review and integration of psychological research on hallucinations. *Psychological Bulletin* 107: 82–95.
- Bentall RP and Slade PD (1985) Reality testing and auditory hallucinations: A signal detection analysis. *British Journal of Clinical Psychology* 24: 159–169.
- Bentall RP, Baker GA and Havers S (1991) Reality monitoring and psychotic hallucinations. *British Journal of Clinical Psychology* 30: 213–222.
- Bhatia T, Garg K, Pogue-Geile M, et al. (2009) Executive functions and cognitive deficits in schizophrenia: Comparisons between probands, parents and controls in India. *Journal of Postgraduate Medicine* 55: 3–7.
- Blakemore SJ, Smith J, Steel R, et al. (2000) The perception of self-produced sensory stimuli in patients with auditory hallucinations and passivity experiences: Evidence for a breakdown in self-monitoring. *Psychological Medicine* 30: 1131–1139.
- Brébion G, David AS, Bressan RA, et al. (2009) Hallucinations and two types of free-recall intrusion in schizophrenia. *Psychological Medicine* 39: 917–926.

- Brébion G, David AS, Jones HM, et al. (2007a) Temporal context discrimination in patients with schizophrenia: Associations with auditory hallucinations and negative symptoms. *Neuropsychologia* 45: 817–823.
- Brébion G, David AS, Ohlsen R, et al. (2007b) Visual memory errors in schizophrenic patients with auditory and visual hallucinations. *Journal of the International Neuropsychological Society* 13: 832–838.
- Brébion G, Ohlsen RI, Pilowsky LS, et al. (2008) Visual hallucinations in schizophrenia: Confusion between imagination and perception. *Neuropsychology* 22: 383–389.
- Brébion G, Ohlsen RI, Pilowsky LS, et al. (2011) Serial and semantic encoding of lists of words in schizophrenia patients with visual hallucinations. *Psychiatry Research* 186: 5–10.
- Brébion G, Stephan-Otto C, Cuevas-Esteban J, et al. (2020) Impaired memory for temporal context in schizophrenia patients with hallucinations and thought disorganisation. *Schizophrenia Research* 220: 225–231.
- Brébion G, Stephan-Otto C, Ochoa S, et al. (2016) Impaired self-monitoring of inner speech in schizophrenia patients with verbal hallucinations and in non-clinical individuals prone to hallucinations. *Frontiers in Psychology* 7: 1381.
- Brébion G, Stephan-Otto C, Ochoa S, et al. (2019) Clinical and non-clinical hallucinations are similarly associated with source memory errors in a visual memory task. *Consciousness and Cognition* 76: 102823.
- Brébion G, Stephan-Otto C, Usall J, et al. (2015) Association of auditory-verbal and visual hallucinations with impaired and improved recognition of colored pictures. *Neuropsychology* 29: 667–674.
- Bristow E, Tabraham P, Smedley N, et al. (2014) Jumping to perceptions and to conclusions: Specificity to hallucinations and delusions. *Schizophrenia Research* 154: 68–72.
- Brookwell ML, Bentall RP and Varese F (2013) Externalizing biases and hallucinations in source-monitoring, self-monitoring and signal detection studies: A meta-analytic review. *Psychological Medicine* 43: 2465–2475.
- Bruder GE, Alschuler DM, Kropfmann CJ, et al. (2011) Heterogeneity of auditory verbal working memory in schizophrenia. *Journal of Abnormal Psychology* 120: 88–97.
- Brunelin J, Combris M, Poulet E, et al. (2006) Source monitoring deficits in hallucinating compared to non-hallucinating patients with schizophrenia. *European Psychiatry* 21: 259–261.
- Bucci S, Startup M, Wynn P, et al. (2008) Referential delusions of communication and reality discrimination deficits in psychosis. *British Journal of Clinical Psychology* 47: 323–334.
- Buckley PF, Miller BJ, Lehrer DS, et al. (2009) Psychiatric comorbidities and schizophrenia. *Schizophrenia Bulletin* 35: 383–402.
- Carter CS, Robertson LC, Nordahl TE, et al. (1996) Perceptual and attentional asymmetries in schizophrenia: Further evidence for a left hemisphere deficit. *Psychiatry Research* 62: 111–119.
- Catts SV, Armstrong MS, Norcross K, et al. (1980) Auditory hallucinations and the verbal transformation effect. *Psychological Medicine* 10: 139–144.
- Chandiramani K and Varma VK (1987) Imagery in schizophrenic patients compared with normal controls. *The British Journal of Medical Psychology* 60: 335–341.
- Chhabra H, Selvaraj S, Sreeraj VS, et al. (2022) Functional near-infrared spectroscopy in schizophrenia patients with auditory verbal hallucinations: Preliminary observations. *Asian Journal of Psychiatry* 73: 103127.
- Chhabra H, Sowmya S, Sreeraj VS, et al. (2016) Auditory false perception in schizophrenia: Development and validation of auditory signal detection task. *Asian Journal of Psychiatry* 24: 23–27.
- Chhabra S, Badcock JC and Maybery MT (2013) Memory binding in clinical and non-clinical psychotic experiences: How does the continuum model fare? *Cognitive Neuropsychiatry* 18: 304–325.
- Chhabra S, Badcock JC, Maybery MT, et al. (2012) Voice identity discrimination in schizophrenia. *Neuropsychologia* 50: 2730–2735.
- Chu RST, Ng CM, Chan KN, et al. (2021) Aberrant learned irrelevance in patients with first-episode schizophrenia-spectrum disorder. *Brain Sciences* 11: 1370.
- Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences*. New York: Routledge.
- Corlett PR, Horga G, Fletcher PC, et al. (2019) Hallucinations and strong priors. *Trends in Cognitive Sciences* 23: 114–127.
- Daalman K, Verkooijen S, Derks EM, et al. (2012) The influence of semantic top-down processing in auditory verbal hallucinations. *Schizophrenia Research* 139: 82–86.
- Damiani S, Donadeo A, Bassetti N, et al. (2022) Understanding source monitoring subtypes and their relation to psychosis: A systematic review and meta-analysis. *Psychiatry and Clinical Neurosciences* 76: 162–171.
- David AS (2004) The cognitive neuropsychiatry of auditory verbal hallucinations: An overview. *Cognitive Neuropsychiatry* 9: 107–123.
- De Leede-Smith S and Barkus E (2013) A comprehensive review of auditory verbal hallucinations: Lifetime prevalence, correlates and mechanisms in healthy and clinical individuals. *Frontiers in Human Neuroscience* 7: 367.
- Docherty NM (2012) Missing referents, psychotic symptoms, and discriminating the internal from the externalized. *Journal of Abnormal Psychology* 121: 416–423.
- Dodgson G and Gordon S (2009) Avoiding false negatives: Are some auditory hallucinations an evolved design flaw? *Behavioural and Cognitive Psychotherapy* 37: 325–334.
- Evans CL, McGuire PK and David AS (2000) Is auditory imagery defective in patients with auditory hallucinations? *Psychological Medicine* 30: 137–148.
- Fernyhough C (2019) Modality-general and modality-specific processes in hallucinations. *Psychological Medicine* 49: 2639–2645.
- Fiszdon JM, Whelahan H, Bryson GJ, et al. (2005) Cognitive training of verbal memory using a dichotic listening paradigm: Impact on symptoms and cognition. *Acta Psychiatrica Scandinavica* 112: 187–193.
- Ford JM and Mathalon DH (2005) Corollary discharge dysfunction in schizophrenia: Can it explain auditory hallucinations? *International Journal of Psychophysiology* 58: 179–189.
- Franck N, Rouby P, Daprati E, et al. (2000) Confusion between silent and overt reading in schizophrenia. *Schizophrenia Research* 41: 357–364.
- Freeman D (2016) Persecutory delusions: A cognitive perspective on understanding and treatment. *The Lancet Psychiatry* 3: 685–692.
- Gaudiot C, Du X, Summerfelt A, et al. (2019) A working memory related mechanism of auditory hallucinations. *Journal of Abnormal Psychology* 128: 423–430.
- Gisselgård J, Anda LG, Brønnick K, et al. (2014) Verbal working memory deficits predict levels of auditory hallucination in first-episode psychosis. *Schizophrenia Research* 153: 38–41.
- Green MF, Hugdahl K and Mitchell S (1994) Dichotic listening during auditory hallucinations in patients with schizophrenia. *American Journal of Psychiatry* 151: 357–362.
- Griffin JD and Fletcher PC (2017) Predictive processing, source monitoring, and psychosis. *Annual Review of Clinical Psychology* 13: 265–289.
- Guillem F, Bicu M, Bloom D, et al. (2001) Neuropsychological impairments in the syndromes of schizophrenia: A comparison between different dimensional models. *Brain and Cognition* 46: 153–159.
- Guillem F, Rinaldi M, Pampoulova T, et al. (2008) The complex relationships between executive functions and positive symptoms in schizophrenia. *Psychological Medicine* 38: 853–860.
- Haddad C, Salameh P, Sacre H, et al. (2023) Effects of antipsychotic and anticholinergic medications on cognition in chronic patients with schizophrenia. *BMC Psychiatry* 23: 61.
- Haddock G, Slade PD, Prasad R, et al. (1996) Functioning of the phonological loop in auditory hallucinations. *Personality and Individual Differences* 20: 753–760.

- Hardy A (2017) Pathways from trauma to psychotic experiences: A theoretically informed model of posttraumatic stress in psychosis. *Frontiers in Psychology* 8: 697.
- He J, Ren H, Li J, et al. (2022) Deficits in sense of body ownership, sensory processing, and temporal perception in schizophrenia patients with/without auditory verbal hallucinations. *Frontiers in Neuroscience* 16: 831714.
- Heilbronner U, Samara M, Leucht S, et al. (2016) The longitudinal course of schizophrenia across the lifespan: Clinical, cognitive, and neurobiological aspects. *Harvard Review of Psychiatry* 24: 118–128.
- Heilbrun AB and Blum NA (1984) Cognitive vulnerability to auditory hallucination: Impaired perception of meaning. *British Journal of Psychiatry* 144: 508–512.
- Heilbrun AB, Blum N and Haas M (1983) Cognitive vulnerability to auditory hallucination: Preferred imagery mode and spatial location of sounds. *British Journal of Psychiatry* 143: 294–299.
- Henquet C, van Os J, Pries LK, et al. (2020) A replication study of JTC bias, genetic liability for psychosis and delusional ideation. *Psychological Medicine* 52: 1777–1783.
- Hoffman RE (1999) New methods for studying hallucinated ‘voices’ in schizophrenia. *Acta Psychiatrica Scandinavica. Supplementum* 99: 89–94.
- Hoffman RE, Rapaport J, Mazure CM, et al. (1999) Selective speech perception alterations in schizophrenic patients reporting hallucinated ‘voices’. *American Journal of Psychiatry* 156: 393–399.
- Hubl D, Moor N, Kindler J, et al. (2020) Speed, accuracy, and efficiency of judgements of coincidence between speech acts and auditory perception during a delayed auditory feedback paradigm: A behavioral study in patients with schizophrenia. *Timing and Time Perception* 8: 177–191.
- Hugdahl K, Løberg EM, Falkenberg LE, et al. (2012) Auditory verbal hallucinations in schizophrenia as aberrant lateralized speech perception: Evidence from dichotic listening. *Schizophrenia Research* 140: 59–64.
- Imrey P (2020) Limitations of meta-analyses of studies with high heterogeneity. *JAMA Network Open* 3: e1919325.
- Ishigaki T and Tanno Y (1999) The signal detection ability of patients with auditory hallucination: Analysis using the continuous performance test. *Psychiatry and Clinical Neurosciences* 53: 471–476.
- Jacobsen P, Peters E, Ward T, et al. (2019) Overgeneral autobiographical memory bias in clinical and non-clinical voice hearers. *Psychological Medicine* 49: 113–120.
- Jardri R and Denève S (2013) Circular inferences in schizophrenia. *Brain* 136: 3227–3241.
- Jenkins LM, Bodapati AS, Sharma RP, et al. (2018) Working memory predicts presence of auditory verbal hallucinations in schizophrenia and bipolar disorder with psychosis. *Journal of Clinical and Experimental Neuropsychology* 40: 84–94.
- Johns LC, Gregg L, Allen P, et al. (2006) Impaired verbal self-monitoring in psychosis: Effects of state, trait and diagnosis. *Psychological Medicine* 36: 465–474.
- Johns LC, Rossell S, Frith C, et al. (2001) Verbal self-monitoring and auditory verbal hallucinations in patients with schizophrenia. *Psychological Medicine* 31: 705–715.
- Jones SR and Fernyhough C (2007) Thought as action: Inner speech, self-monitoring, and auditory verbal hallucinations. *Consciousness and Cognition* 16: 391–399.
- Kapur S (2003) Psychosis as a state of aberrant salience: A framework linking biology, phenomenology, and pharmacology in schizophrenia. *American Journal of Psychiatry* 160: 13–23.
- Kay SR, Fiszbein A and Opler LA (1987) The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin* 13: 261–276.
- Kayser J, Tenke CE, Kroppmann CJ, et al. (2012) A neurophysiological deficit in early visual processing in schizophrenia patients with auditory hallucinations. *Psychophysiology* 49: 1168–1178.
- Kot T and Serper M (2002) Increased susceptibility to auditory conditioning in hallucinating schizophrenic patients: A preliminary investigation. *Journal of Nervous and Mental Disease* 190: 282–288.
- Kråkvik B, Stiles T and Hugdahl K (2013) Experiencing malevolent voices is associated with attentional dysfunction in psychotic patients. *Scandinavian Journal of Psychology* 54: 72–77.
- Lakens D (2013) Calculating and reporting effect sizes to facilitate cumulative science: A practical primer for t-tests and ANOVAs. *Frontiers in Psychology* 4: 863.
- Laloyaux J, Della Libera C and Larøi F (2018) Source flexibility in schizophrenia: Specificity and role in auditory hallucinations. *Cognitive Neuropsychiatry* 23: 393–407.
- Lee JS, Chun JW, Lee SH, et al. (2013) Association of impaired reality processing with psychotic symptoms in schizophrenia. *Psychiatry Research* 210: 721–728.
- Lee SH, Chung YC, Yang JC, et al. (2004) Abnormal speech perception in schizophrenia with auditory hallucinations. *Acta Neuropsychiatrica* 16: 154–159.
- Lenhard W and Lenhard A (2016) Computation of effect sizes. Available at: www.psychometrica.de/effect_size.html (accessed 16 December 2022).
- Levitan C, Ward PB and Catts SV (1999) Superior temporal gyrus volumes and laterality correlates of auditory hallucinations in schizophrenia. *Biological Psychiatry* 46: 955–962.
- Li CSR, Chen MC, Yang YY, et al. (2002) Altered performance of schizophrenia patients in an auditory detection and discrimination task: Exploring the ‘self-monitoring’ model of hallucination. *Schizophrenia Research* 55: 115–128.
- Løberg EM, Jørgensen HA and Hugdahl K (2004) Dichotic listening in schizophrenic patients: Effects of previous vs. ongoing auditory hallucinations. *Psychiatry Research* 128: 167–174.
- Løberg EM, Jørgensen HA, Kroken RA, et al. (2015) Auditory verbal hallucinations reflect stable auditory attention deficits: A prospective study. *Cognitive Neuropsychiatry* 20: 81–94.
- McCarthy-Jones S and Longden E (2016) Auditory verbal hallucinations in schizophrenia and post-traumatic stress disorder: Common phenomenology, common cause, common interventions? *Frontiers in Psychology* 6: 1071.
- McCarthy-Jones S, Smailes D, Corvin A, et al. (2017) Occurrence and co-occurrence of hallucinations by modality in schizophrenia-spectrum disorders. *Psychiatry Research* 252: 154–160.
- McKay CM, Headlam DM and Copolov DL (2000) Central auditory processing in patients with auditory hallucinations. *American Journal of Psychiatry* 157: 759–766.
- McKay SE, Golden CJ and Scott M (1981) Neuropsychological correlates of auditory and visual hallucinations. *International Journal of Neuroscience* 15: 87–94.
- McLachlan NM, Phillips DS, Rossell SL, et al. (2013) Auditory processing and hallucinations in schizophrenia. *Schizophrenia Research* 150: 380–385.
- Moher D, Shamseer L, Clarke M, et al. (2015) Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 4: 1.
- Morrison AP and Haddock G (1997) Cognitive factors in source monitoring and auditory hallucinations. *Psychological Medicine* 27: 669–679.
- Moseley P, Alderson-Day B, Common S, et al. (2022) Continuities and discontinuities in the cognitive mechanisms associated with clinical and nonclinical auditory verbal hallucinations. *Clinical Psychological Science* 10: 752–766.
- Ocklenburg S, Westerhausen R, Hirnstein M, et al. (2013) Auditory hallucinations and reduced language lateralization in schizophrenia: A meta-analysis of dichotic listening studies. *Journal of the International Neuropsychological Society* 19: 410–418.
- Olsson O and Nielzen S (1999a) Contralateral induction by frequency spectrum in hallucinating schizophrenics. *Psychiatry Research* 87: 65–75.

- Olsson O and Nielzén S (1999b) Perceptual restoration of missing sounds in a group of hallucinating schizophrenics. *Perceptual and Motor Skills* 89: 315–326.
- Papageorgiou C, Oulis P, Vasios C, et al. (2003) Do atypical antipsychotics fail to exert cognitive sparing effects? *NeuroReport* 14: 505–509.
- Paulik G, Steel C and Arntz A (2019) Imagery rescripting for the treatment of trauma in voice hearers: A case series. *Behavioural and Cognitive Psychotherapy* 47: 709–725.
- Perrin MA, Kantowitz JT, Silipo G, et al. (2018) Mismatch negativity (MMN) to spatial deviants and behavioral spatial discrimination ability in the etiology of auditory verbal hallucinations and thought disorder in schizophrenia. *Schizophrenia Research* 191: 140–147.
- Pinheiro AP, Rezaii N, Rauber A, et al. (2016) Is this my voice or yours? The role of emotion and acoustic quality in self-other voice discrimination in schizophrenia. *Cognitive Neuropsychiatry* 21: 335–353.
- Powers AR, Mathys C and Corlett PR (2017) Pavlovian conditioning-induced hallucinations result from overweighting of perceptual priors. *Science* 357: 596–600.
- Rocca P, Castagna F, Marchiaro L, et al. (2006) Neuropsychological correlates of reality distortion in schizophrenic patients. *Psychiatry Research* 145: 49–60.
- Rossell SL and Boundy CL (2005) Are auditory-verbal hallucinations associated with auditory affective processing deficits? *Schizophrenia Research* 78: 95–106.
- Rossell SL, Shapleske J, Fukuda R, et al. (2001) Corpus callosum area and functioning in schizophrenic patients with auditory-verbal hallucinations. *Schizophrenia Research* 50: 9–17.
- Rossell SL, Van Rheenen TE, Groot C, et al. (2013) Investigating affective prosody in psychosis: A study using the Comprehensive Affective Testing System. *Psychiatry Research* 210: 896–900.
- Schnakenberg Martin AM, Bartolomeo L, Howell J, et al. (2018) Auditory feature perception and auditory hallucinatory experiences in schizophrenia spectrum disorder. *European Archives of Psychiatry and Clinical Neuroscience* 268: 653–661.
- Schneider SJ and Wilson CR (1983) Perceptual discrimination and reaction time in hallucinatory schizophrenics. *Psychiatry Research* 9: 243–253.
- Seal ML, Aleman A and McGuire PK (2004) Compelling imagery, unanticipated speech and deceptive memory: Neurocognitive models of auditory verbal hallucinations in schizophrenia. *Cognitive Neuropsychiatry* 9: 43–72.
- Seal ML, Crowe SF and Cheung P (1997) Deficits in source monitoring in subjects with auditory hallucinations may be due to differences in verbal intelligence and verbal memory. *Cognitive Neuropsychiatry* 2: 273–290.
- Selvaraj S, Chhabra H, Dinakaran D, et al. (2021) Auditory signal detection in schizophrenia: Correlates with auditory verbal hallucinations & effect of single session transcranial direct current stimulation (tDCS). *Psychiatry Research* 297: 113704.
- Shah D, Knott V, Baddeley A, et al. (2018) Impairments of emotional face processing in schizophrenia patients: Evidence from P100, N170 and P300 ERP components in a sample of auditory hallucinators. *International Journal of Psychophysiology* 134: 120–134.
- Shea TL, Sergejew AA, Burnham D, et al. (2007) Emotional prosodic processing in auditory hallucinations. *Schizophrenia Research* 90: 214–220.
- Siddi S, Petretto DR, Burrai C, et al. (2017) The role of set-shifting in auditory verbal hallucinations. *Comprehensive Psychiatry* 74: 162–172.
- Siddi S, Petretto DR, Scanu R, et al. (2016) Deficits in metaphor but not in idiomatic processing are related to verbal hallucinations in patients with psychosis. *Psychiatry Research* 246: 101–112.
- Smailes D, Alderson-Day B, Hazell C, et al. (2022) Measurement practices in hallucinations research. *Cognitive Neuropsychiatry* 27: 183–198.
- Smith P and Gudjonsson G (1995) The relationship of mental disorder to suggestibility and confabulation among forensic inpatients. *Journal of Forensic Psychiatry* 6: 499–515.
- Sommer IE, Daalman K, Rietkerk T, et al. (2010) Healthy individuals with auditory verbal hallucinations; Who are they? psychiatric assessments of a selected sample of 103 subjects. *Schizophrenia Bulletin* 36: 633–641.
- Steel C (2015) Hallucinations as a trauma-based memory: Implications for psychological interventions. *Frontiers in Psychology* 6: 1262.
- Stephane M (2019) The self, agency and spatial externalizations of inner verbal thoughts, and auditory verbal hallucinations. *Frontiers in Psychiatry* 10: 668.
- Stephane M, Kuskowski M, McClannahan K, et al. (2010a) Evaluation of inner-outer space distinction and verbal hallucinations in schizophrenia. *Cognitive Neuropsychiatry* 15: 441–450.
- Stephane M, Kuskowski M, McClannahan K, et al. (2010b) Evaluation of speech misattribution bias in schizophrenia. *Psychological Medicine* 40: 741–748.
- Sterzer P, Adams RA, Fletcher P, et al. (2018) The predictive coding account of psychosis. *Biological Psychiatry* 84: 634–643.
- Strawson WH, Wang HT, Quadt L, et al. (2021) Voice hearing in Borderline Personality Disorder across perceptual, subjective and neural dimensions. *International Journal of Neuropsychopharmacology* 25: 375–386.
- Sun Q, Fang Y, Shi Y, et al. (2021) Inhibitory top-down control deficits in schizophrenia with auditory verbal hallucinations: A Go/NoGo task. *Frontiers in Psychiatry* 12: 544746.
- Szöke A, Trandafir A, Dupont ME, et al. (2008) Longitudinal studies of cognition in schizophrenia: Meta-analysis. *The British Journal of Psychiatry* 192: 248–257.
- Thakkar KN, Mathalon DH and Ford JM (2021) Reconciling competing mechanisms posited to underlie auditory verbal hallucinations. *Philosophical Transactions of the Royal Society B-Biological Sciences* 376: 20190702.
- Thomas ML, Bismark AW, Joshi YB, et al. (2018) Targeted cognitive training improves auditory and verbal outcomes among treatment refractory schizophrenia patients mandated to residential care. *Schizophrenia Research* 202: 378–384.
- Thomas N, Hayward M, Peters E, et al. (2014) Psychological therapies for auditory hallucinations (voices): Current status and key directions for future research. *Schizophrenia Bulletin* 40: S202–S212.
- Thompson IM and Copolov DL (1998) The psycholinguistics of auditory hallucinations. *Aphasiology* 12: 919–932.
- Titone D and Levy DL (2004) Lexical competition and spoken word identification in schizophrenia. *Schizophrenia Research* 68: 75–85.
- Toh WL, McCarthy-Jones S, Copolov D, et al. (2019) Have we overlooked the significance of multinodal hallucinations in schizophrenia? *Psychiatry Research* 279: 358–360.
- Toh WL, Tan EJ, Neill E, et al. (2020) Identifying the cognitive underpinnings of voice-hearing by comparing never, past and current voice-hearers. *Acta Psychiatrica Scandinavica* 141: 553–562.
- Tracy DK and Shergill SS (2013) Mechanisms underlying auditory hallucinations – Understanding perception without stimulus. *Brain Sciences* 3: 642–669.
- Uptegrove R, Broome MR, Caldwell K, et al. (2016) Understanding auditory verbal hallucinations: A systematic review of current evidence. *Acta Psychiatrica Scandinavica* 133: 352–367.
- Versmissen D, Janssen I, Johns L, et al. (2007) Verbal self-monitoring in psychosis: A non-replication. *Psychological Medicine* 37: 569–576.
- Versmissen D, Janssen I, Myin-Germeys I, et al. (2008) Evidence for a relationship between mentalising deficits and paranoia over the psychosis continuum. *Schizophrenia Research* 99: 103–110.
- Waters F, Allen P, Aleman A, et al. (2012a) Auditory hallucinations in schizophrenia and nonschizophrenia populations: A review and

- integrated model of cognitive mechanisms. *Schizophrenia Bulletin* 38: 683–693.
- Waters F, Badcock JC, Maybery MT, et al. (2003) Inhibition in schizophrenia: Association with auditory hallucinations. *Schizophrenia Research* 62: 275–280.
- Waters F, Badcock JC, Michie PT, et al. (2006) Auditory hallucinations in schizophrenia: Intrusive thoughts and forgotten memories. *Cognitive Neuropsychiatry* 11: 65–83.
- Waters F, Maybery MT, Badcock JC, et al. (2004) Context memory and binding in schizophrenia. *Schizophrenia Research* 68: 119–125.
- Waters F, Woodward T, Allen P, et al. (2012b) Self-recognition deficits in schizophrenia patients with auditory hallucinations: A meta-analysis of the literature. *Schizophrenia Bulletin* 38: 741–750.
- Wells G, Shea B, O'Connell D, et al. (2000) The Newcastle–Ottawa Scale (NOS) for assessing the quality of non-randomized studies in meta-analysis. Available at: www.ohri.ca/programs/clinical_epidemiology/oxford.asp (accessed 27 June 2019).
- Woodward TS, Menon M and Whitman JC (2007) Source monitoring biases and auditory hallucinations. *Cognitive Neuropsychiatry* 12: 477–494.
- Wykes T, Huddy V, Cellard C, et al. (2011) A meta-analysis of cognitive remediation for schizophrenia: Methodology and effect sizes. *American Journal of Psychiatry* 168: 472–485.
- Yang F, Zhu H, Yu L, et al. (2021) Deficits in multi-scale top-down processes distorting auditory perception in schizophrenia. *Behavioural Brain Research* 412: 113411.
- Zenisek R, Thaler NS, Sutton GP, et al. (2015) Auditory processing deficits in bipolar disorder with and without a history of psychotic features. *Bipolar Disorders* 17: 769–780.
- Zhuo C, Chen M, Xu Y, et al. (2020) Reciprocal deterioration of visual and auditory hallucinations in schizophrenia presents V-shaped cognition impairment and widespread reduction in brain gray matter-A pilot study. *Journal of Clinical Neuroscience* 79: 154–159.